

INCA™

PCS 04457

A suspension concentrate formulation containing 125 g/l (12.1% w/w) epoxiconazole.
Also contains solvent naphtha and alkoxyated fatty alcohol.

A broad-spectrum systemic fungicide for control of *Septoria*, yellow and brown rust, eyespot, net blotch, *Rhynchosporium*, powdery mildew, *Fusarium* and sooty moulds in barley, wheat, triticale, rye and oats.



	<p>SAFETY INFORMATION</p> <p>Danger</p> <p>Harmful if inhaled. Suspected of causing cancer. May damage the unborn child. Suspected of damaging fertility. Very toxic to aquatic life. Toxic to aquatic life with long lasting effects. Repeated exposure may cause skin dryness or cracking.</p> <p>Keep out of reach of children. Obtain special instructions before use. Avoid breathing vapours or spray. Wear protective gloves/protective clothing and eye protection/face protection. IF INHALED: Remove person to fresh air and keep comfortable for breathing. Dispose of contents/container to a licensed waste disposal contractor or collection site except for triple rinsed, empty containers which can be disposed of as non-hazardous waste. To protect aquatic organisms respect an unsprayed buffer zone of 5 m to surface water bodies.</p> <p>To avoid risks to human health and the environment, comply with the instructions for use.</p> <p>PCS 04457</p>
--	--

FR R3.2.3 AUG18

5 litres e

To access the Safety Data Sheet for this product scan the QR code or use the weblink: <https://bit.ly/2KOuili>
- Alternatively, contact your supplier.



Scan for Safety Data Sheet

ADAMA

97050970B_201806

INCA™

PCS 04457

Adama Agricultural Solutions UK Ltd

Unit 15, Thatcham Business Village, Colthrop Way, Thatcham, Berkshire RG19 4LW

Telephone: +44 (0)1635 860555 | Technical Helpline: +44 (0)1635 876622

www.adama.com | ukenquiries@adama.com

PROTECT FROM FROST
SHAKE WELL BEFORE USE

**For advice on medical emergencies, fires or major spills telephone
the UK National Chemical Emergency Centre on +44 (0)1865 407333**

DIRECTIONS FOR USE:
full instructions in this leaflet.

Batch No.: see packaging

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crop	Maximum individual dose L/ha	Maximum number of applications	Maximum total dose L/ha	Latest time of application
Winter wheat, spring wheat, triticale and rye	1.0	2 per crop	2.0	Before end of flowering (GS 69)
Winter barley, spring barley, winter oats, spring oats	1.0	2 per crop	2.0	Before end of heading (GS 59)

Other specific restrictions:

A minimum interval of 21 days must be observed between applications.

To protect aquatic organisms respect an unsprayed buffer zone of 5 m to surface water bodies.

This leaflet is part of the approved label.

For advice on medical emergencies, fires or major spills telephone
the UK National Chemical Emergency Centre on +44 (0)1865 407333

A suspension concentrate formulation containing 125 g/l (12.1% w/w) epoxiconazole.

Also contains solvent naphtha and alkoxyated fatty alcohol. A broad-spectrum systemic fungicide for control of *Septoria*, yellow and brown rust, eyespot, net blotch, *Rhynchosporium*, powdery mildew, *Fusarium* and sooty moulds in barley, wheat, triticale, rye and oats.



SAFETY INFORMATION

Danger

Harmful if inhaled.
Suspected of causing cancer.
May damage the unborn child.
Suspected of damaging fertility.
Very toxic to aquatic life.
Toxic to aquatic life with long lasting effects.
Repeated exposure may cause skin dryness or cracking.

Keep out of reach of children.
Obtain special instructions before use.
Avoid breathing vapours or spray.
Wear protective gloves/protective clothing and eye protection/face protection.
IF INHALED: Remove person to fresh air and keep comfortable for breathing.

Dispose of contents/container to a licensed waste disposal contractor or collection site except for triple rinsed, empty containers which can be disposed of as non-hazardous waste.

To protect aquatic organisms respect an unsprayed buffer zone of 5 m to surface water bodies.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS 04457

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crop	Maximum individual dose L/ha	Maximum number of applications	Maximum total dose L/ha	Latest time of application
Winter wheat, spring wheat, triticale and rye	1.0	2 per crop	2.0	Before end of flowering (GS 69)
Winter barley, spring barley, winter oats, spring oats	1.0	2 per crop	2.0	Before end of heading (GS 59)

Other specific restrictions:

A minimum interval of 21 days must be observed between applications.

To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

RESTRICTIONS

Avoid drift onto neighbouring crops, as damage may occur, particularly to broad-leaved plants.

Malting barley

INCA™ is accepted by the Brewers and Licensed Retailers Association to be used on malting barley. Application should be before the start of ear emergence.

Avoid overlapping spray swaths.

Crops should not be re-entered until spray residues are dry.

RESISTANCE

INCA contains a DMI fungicide. Resistance to some DMI fungicides has been identified in *Septoria* leaf blotch (*Zymoseptoria tritici*) which may seriously affect the performance of some products. For further advice on resistance management in DMIs, contact your agronomist or specialist advisor.

The efficacy of this product may be affected if strains of disease pathogens resistant or less sensitive occur at any time. As such occurrences cannot be predicted no responsibility can be accepted for the results obtained.

Where strains of wheat powdery mildew with decreased sensitivity to triazoles occur or develop, INCA may not give satisfactory control.

DISEASE CONTROL

Specific diseases controlled					
	Wheat	Barley	Oats	Triticale	Rye
Septoria leaf blotch	MC	-	-	MC	-*
Septoria glume blotch	MC	-	-	MC	-*
Powdery mildew	-	-	C	-	C
Yellow rust	C	C	-	C	C
Brown rust	C	C	-	C	C
Leaf scald <i>Rhynchosporium</i>	-	MC	-	-	MC
Net blotch	-	R	-	-	-
<i>Fusarium</i> ear blight	R	-	-	-	-
Sooty moulds (<i>Cladosporium</i>, <i>Botrytis</i>)	R	-	-	-	-

C = Control MC = Moderate control R = Reduction * No claim for control on Rye

CROP SPECIFIC INFORMATION

INCA can be safely used on all varieties of wheat, barley, oats, rye and triticale.

For best disease control, apply as soon as disease is seen. Some specific recommendations can be seen below.

Disease	Specific recommendations
<i>Fusarium</i> ear blight and sooty moulds	Apply during ear emergence.
Mildew	For better mildew control when disease is established, apply with a specific mildew fungicide.
<i>Zymoseptoria tritici</i> and <i>nodorum</i>	Apply when disease is present, or conditions favour disease appearing. This is generally after the 3rd node detectable stage. A follow up treatment may be required.
Eyespot	For optimum control apply between leaf sheath erect stage and 2nd node detectable.

RATE OF USE

Apply 1.0 L/ha in 150-300 L water/ha. Use the higher volume of water for dense crops, or for the second application.

Apply as a medium spray as defined by BCPC.

MIXING AND APPLICATION

For use by tractor mounted/trailed sprayer only.

Half-fill the spray tank with clean water and begin agitation. Add the required quantity of INCA and add the remainder of the water.

Continue agitation until spraying is completed.

On emptying the container, RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely.

Wash out sprayer thoroughly after use.

COMPATIBILITY

For information on tank-mixes, consult Adama Agricultural Solutions UK Ltd, your agronomist or your distributor.

DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

Authorisation Holder and Marketed by

Adama Agricultural Solutions UK Ltd
Unit 15, Thatcham Business Village
Colthrop Way, Thatcham, Berkshire RG19 4LW
Telephone: 00 44 1635 860555
Technical Helpline: 00 44 1635 876622
www.adama.com
Email: ukenquiries@adama.com

INCA™ is a trademark of a company of the Adama Group.
© Adama Agricultural Solutions UK Ltd