

LEGION[®]

MAPP 16249 PCS 05179

A residual herbicide for the control of broad-leaved weeds and annual meadowgrass in winter oilseed rape.

A suspension concentrate formulation containing 375 g/l (32.9% w/w) metazachlor and 125 g/l (10.9% w/w) quinmerac.

Contains metazachlor and 1,2-benzisothiazol-3(2H)-one.

May produce an allergic reaction.



ADAMA



Scan for Safety Data Sheet or use weblink:
<https://bit.ly/2qU8flx>

FOR PROFESSIONAL USE ONLY



FR R3.2.6 APR20

10Le

Warning

Suspected of causing cancer.

Very toxic to aquatic life with long lasting effects.

Contains metazachlor and 1,2-benzisothiazol-3(2H)-one.

May produce an allergic reaction.

Keep out of reach of children.

Do not handle until all safety precautions have been read and understood.

Wear protective gloves.

If exposed or concerned: get medical advice/attention.

Store locked up.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed, empty containers which can be disposed of as non-hazardous waste.

Ireland only: To protect aquatic organisms respect an unsprayed buffer zone of 5 m to surface water bodies.

To avoid risks to human health and the environment, comply with the instructions for use.

MAPP 16249 PCS 05179

NA7384_02

LEGION[®]

MAPP 16249/PCS 05179

SHAKE WELL BEFORE USE

DIRECTIONS FOR USE:

full instructions in this leaflet.

Batch No.: see packaging

Adama Agricultural Solutions UK Ltd

Third Floor East, 1410 Arlington Business Park, Theale, Reading RG7 4SA

Telephone: +44 (0)1635 860555 | Technical Helpline: +44 (0)1635 876622

www.adama.com | ukenquiries@adama.com

For advice on medical emergencies, fires or major spills telephone the UK National Chemical Emergency Centre on +44 (0)1865 407333

UK only - The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

OPEN HERE ►

SAFETY PRECAUTIONS

Operator Protection

UK only: Engineering control of operator exposure must be used where reasonably practicable, in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate or contaminated surfaces.

UK only: However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection. WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

Environmental Protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

UK only: DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing waterbody, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application.

UK only: This product qualifies for inclusion within the Local Environmental Risk Assessment for Pesticides (LERAP) scheme.



Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with HSE's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for inspection for three years.

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area

Storage and Disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place. WASH OUT CONTAINER THOROUGHLY empty washings into spray tank and dispose of safely.

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL HERBICIDE

Crop	Maximum individual dose (L product/ha)	Maximum number of applications	Latest time of application
Winter oilseed rape	2.0	1	Up to 4 leaf stage (BBCH 14) and before end of January in the year of harvest

Other specific restrictions:

A maximum total dose of not more than 1000 g metazachlor/hectare may be applied in a three year period on the same field.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

BK R3.2.6 APR20

S7384B_02



This leaflet is part of the approved label.

For advice on medical emergencies, fires or major spills telephone the UK National Chemical Emergency Centre on +44 (0)1865 407333

A residual herbicide for the control of broad-leaved weeds and annual meadowgrass in winter oilseed rape.

A suspension concentrate formulation containing 375 g/l (32.9% w/w) metazachlor and 125 g/l (10.9% w/w) quinmerac. Contains metazachlor and 1,2-benzisothiazol-3(2H)-one. May produce an allergic reaction.



Warning

Suspected of causing cancer.
Very toxic to aquatic life with long lasting effects.
Contains metazachlor and 1,2-benzisothiazol-3(2H)-one.
May produce an allergic reaction.

Keep out of reach of children.

Do not handle until all safety precautions have been read and understood.

Wear protective gloves.

If exposed or concerned: get medical advice/attention.

Store locked up.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed, empty containers which can be disposed of as non-hazardous waste.

Ireland only: To protect aquatic organisms respect an unsprayed buffer zone of 5 m to surface water bodies.

To avoid risks to human health and the environment, comply with the instructions for use.



MAPP 16249 PCS 05179

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL HERBICIDE

Crop	Maximum individual dose (L product/ha)	Maximum number of applications	Latest time of application
Winter oilseed rape	2.0	1	Up to 4 leaf stage (BBCH 14) and before end of January in the year of harvest

Other specific restrictions:

A maximum total dose of not more than 1000 g metazachlor/hectare may be applied in a three year period on the same field.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

SAFETY PRECAUTIONS

Operator Protection

UK only: Engineering control of operator exposure must be used where reasonably practicable, in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate or contaminated surfaces.

UK only: However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

Environmental Protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

UK only: DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing waterbody, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application.



UK only: This product qualifies for inclusion within the Local Environmental Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with HSE's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for inspection for three years.

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area

Storage and Disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

LEGION® is a suspension concentrate formulation containing metazachlor and quinmerac for use as a residual herbicide in all varieties of winter oilseed rape. It is primarily taken up by the roots but also has some leaf uptake.

LEGION is recommended as a pre- and post-emergence herbicide.

Post-emergence application is advised where it is not possible to make pre-emergence applications due to light or stony soils, those with organic matter content higher than 10%, adverse weather conditions, poor seedbed preparation or missed timing of crop.

RESTRICTIONS OR WARNINGS

When LEGION is used pre-emergence it is important to ensure the herbicide does not come into contact with the crop seed by drilling the seed to a depth of 15 mm and ensuring it is well covered by soil before spraying.

Do not use pre-emergence on broadcast crops.

Loose or puffy seedbeds should be consolidated before treatment. Seedbeds should be moist, firm and free from clods.

Adequate moisture is required for good weed control.

Do not apply pre-emergence shortly before heavy rainfall is forecast since this can reduce crop vigour, weed control and possibly plant stand. This will normally be outgrown.

Do not apply under adverse weather conditions.

Some damage may occur, especially on brashy or stony soils. This will normally be outgrown.

Do not treat crops which are suffering from stress.

Do not apply to crops with a compaction layer or to poorly drained soils since damage can occur under waterlogged conditions.

Do not overlap spray swaths.

Avoid drift onto neighbouring crops.

LEGION is suitable for use on all soil types as defined by Soil Texture (85) System, except sands, very light soils and soils containing more than 10% organic matter.

Under frosty conditions a transient scorch may occur. On clay soils create a fine, consolidated seedbed to slow the downward movement of water.

Do not apply LEGION to dry soils. Moist soils have fewer and smaller cracks.

Do not apply LEGION if heavy rain is forecast, wait until after the event.

Crops should not be re-entered until spray residues are dry.

RESISTANCE

LEGION contains two active ingredients with different modes of action and therefore the risk of resistance building up is reduced. LEGION contains metazachlor, a Group K3 herbicide and quinmerac a Group O based on the mode of action classification system of the Herbicide Resistance Action Committee.

Strains of some annual grasses, e.g. blackgrass, wild oats, and Italian ryegrass, have developed resistance to herbicides which may lead to poor control. A strategy for preventing and managing such resistance should be adopted. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the HGCA, CPA, your distributor, crop adviser or product manufacturer.

WEEDS CONTROLLED

Pre-emergence

Susceptible

Annual meadowgrass
Blackgrass
Cleavers
Common chickweed

Common poppy
Field forget-me-not
Common field-speedwell
Common groundsel

Mayweed spp
Parsley-piert
Red dead-nettle
Shepherd's purse

Moderately resistant

Cut-leaved Crane's-bill

Resistant

Field pansy

Weeds germinating from depth may not be controlled.

Post-emergence – max growth stage

Annual meadowgrass
Blackgrass
Cleavers
Common chickweed
Common poppy
Cut-leaved Crane's-bill
Field forget-me-not

Pre-em
Pre-em
2 whorls
Cotyledon
2 leaves
-
Pre-em

Common field-speedwell
Common groundsel
Mayweed spp
Field pansy
Parsley-piert
Red dead-nettle
Shepherd's Purse

2 leaves
Pre-em
4 leaves
-
Pre-em
Cotyledon (MS)
Pre-em

CROP SPECIFIC INFORMATION

LEGION can be used on all varieties of winter oilseed rape.

Timing

May be applied pre or post-emergence of the crop up to 4 leaf stage (BBCH 14) and before end of January in year of harvest. Best weed control is achieved when applied pre-emergence of the weeds. The soil should be moist at time of application or rainfall should fall shortly afterwards.

Pre-crop emergence

Applications should be made before the crop seed chits, under warm moist conditions this may occur within 48 hours of drilling.

Post-crop emergence

Applications can be made from when the majority of the crop has two fully expanded cotyledons until 4 leaves (BBCH 14) and before the end of January in year of harvest.

RATE OF USE

Apply at 2.0 L/ha in 150-250 L water/ha.

FOLLOWING CROPS

Any crop can be planted after normal harvesting of a crop treated with LEGION. Ploughing is not required.

Where the crop fails in the autumn or winter, oilseed rape may be re-drilled immediately, providing the soil is ploughed, otherwise a minimum interval of 30 days is required after minimum cultivations.

Cereals may be drilled after ploughing providing 2 months have elapsed, otherwise 3 months should elapse if the field has minimum cultivations.

In the spring following crop failure, oilseed rape, cereals, sugar beet, maize, peas, potatoes and other crops including forage crops can be sown.

MIXING AND SPRAYING

Shake well before use.

Half-fill the spray tank with clean water and begin agitation.

Add the required quantity of LEGION and add the remainder of the water.

On emptying the container, RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely.

Continue agitation until spraying is completed.

Wash out sprayer thoroughly after use.

Apply as a MEDIUM spray as defined by BCPC.

COMPANY ADVISORY INFORMATION

– UK only

This section is not part of the Product Label under Regulation (EC) 1107/2009 and provides additional advice on product use.



OSR Herbicides?
Think Water

This product contains **metazachlor** plus **quinmerac** and is therefore included in the 'OSR Herbicides? **Think Water**' stewardship campaign. Particular care is needed when using in **Surface Water Drinking Water Safeguard Zones**:-

- **Do not use after** 30 September on drained fields, including mole-drained.

In other areas:-

- Avoid use after 15 October on drained fields, including mole-drained.

For further information, see www.adama.com and www.voluntaryinitiative.org.uk or contact Adama on 01635 876622.

DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

Marketed by

Adama Agricultural Solutions UK Ltd
Third Floor East, 1410 Arlington Business Park
Theale, Reading RG7 4SA
Telephone: + 44 (0)1635 860555
Technical Helpline: + 44 (0)1635 876622
www.adama.com
Email: ukenquiries@adama.com

LEGION® is a registered trademark of a company of the Adama Group.

© Adama Agricultural Solutions UK Ltd