

INSECTICIDE

# ALIAS<sup>®</sup> 240 SC FOLIAR APPLICATION

Control your worst cereal, potato, soybean, blueberry, vegetable and tree fruit insects with a long-lasting foliar application.



## ACTIVE INGREDIENT

Imidacloprid

## CHEMISTRY GROUP

Group 4

## APPLICATION RATES AND PACKAGING

- 70–190 ml/ac or 54–20 ac/3.785 L jug
- 4 x 3.785 L jugs/case

## REGISTERED CROPS:

- Apples
- Brussels sprouts
- Caneberries (Subgroup 13A)
- Cherries<sup>1</sup>
- Cole (Group 5)
- Eggplant
- Highbush blueberries<sup>2</sup>
- Lettuce<sup>3</sup>
- Nectarines
- Peaches
- Potatoes
- Tomatoes

<sup>1</sup>BC, ON only

<sup>2</sup>BC

<sup>3</sup>BC, ON, PQ, PEI, NS only

## KEY INSECTS CONTROLLED:

- Aphids
- Black cherry fruit fly
- Colorado potato beetle
- Leaf hoppers
- Mullein bug
- Potato flea beetles
- Soybean aphid
- Tentiform leafminer
- Western cherry fruit fly

## HOW IT WORKS:

Neonicotinoid chemistry provides control of insect pests through a combination of contact and ingestion.

## APPLICATION TIMING AND CROP STAGING:

The need and timing of application should be based on the presence of pests at vulnerable developmental stages and significant populations as determined by local monitoring. Consult label for specific crop and insect timing.

## WATER VOLUME:

Apply by ground application in a water volume that provides good coverage of foliage and insect pests.

## RAINFASTNESS:

Avoid application when heavy rain is forecast.

# ALIAS<sup>®</sup> 240 SC FOLIAR APPLICATION

## SUPPORTED TANK MIXES:

None registered.

## MIXING INSTRUCTIONS:

1. Shake well before using.
2. Add a portion of the required amount of water to the spray tank with agitation.
3. Add required amount of Alias<sup>®</sup> 240 SC.
4. Complete filling the tank with balance of water.
5. Maintain sufficient agitation during mixing and application.

## CROP ROTATIONS:

Acceptable plant-back intervals for:

- Cereal grains (wheat, barley, oats): minimum 30 days
- Peas and beans: 9 months
- All other food and feed crops: 12 months

## PRE-HARVEST INTERVALS:

- Caneberries (Subgroup 13A): 4 days
- Cherries: 10 days
- Highbush blueberries: 14 days
- Potatoes, field lettuce, cole (Group 5), apples, peaches, nectarines, brussels sprouts and eggplant: 7 days

## GRAZING RESTRICTIONS

Do not graze or harvest cover crops for food or feed.

## STORAGE

Store in cool, dry place.

### QUICK TIPS:

For optimal control, good coverage of the foliage is needed.