

NIMITZ®

Nematicide
EPA REG. NO. 66222-243

SUPPLEMENTAL LABELING

This label expires on April 4, 2023 and must not be distributed or used after that date.

READ THE ENTIRE LABEL FOR NIMITZ® BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for NIMITZ® and this supplement.

ACTIVE INGREDIENT:	% BY WT.
Fluensulfone: 5-chloro-2 (3,4,4-trifluoro-but-3-ene-1-sulfonyl)-thiazole	40.0%
OTHER INGREDIENTS*:	<u>60.0%</u>
TOTAL:	100.0%

Contains 4 pounds fluensulfone per gallon in an Emulsifiable Concentrate formulation.

*Contains petroleum distillates

Root Vegetables (Except Sugar Beet) Subgroup 1B; Tuberous and Corm Vegetables (Subgroup 1C)

KEEP OUT OF REACH OF CHILDREN

CAUTION/PRECAUCION

*Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)*

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This supplemental labeling must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for NIMITZ® before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.
- These directions can be found on the currently registered EPA Stamped Label.

<p>ROOT VEGETABLES (EXCEPT SUGAR BEET) Subgroup 1B</p> <p>Carrot, Radish, Turnip and other root vegetables including:</p> <p>Beet, garden; burdock, edible; celeriac; chervil, turnip-rooted; chicory; ginseng; horseradish; parsley, turnip-rooted; parsnip; radish, oriental; rutabaga; salsify; salsify, black; salsify, Spanish; skirret;</p>	<p>3.5 to 7 pints/ treated acre (56 to 112 fl. oz./treated acre)</p>	<p>Application must be made 7 or more days before planting for seeded crops.</p> <p>Do not apply more than one application per crop and no more than 3.5 lb fluensulfone per acre per calendar year.</p> <p>Turnip and Japanese white radish (daikon)* Transplanting: Apply at a minimum of 30 days prior to transplanting.</p>
---	--	---

		Direct seeding: apply NIMITZ a minimum of 30 days before seeding and make 2 supplemental irrigations prior to planting as per Chart 1.
<p>Tuberous and Corm Vegetables (Subgroup 1C)</p> <p>Potato, sweet potato, yam and other root vegetables including:</p> <p>arracacha; arrowroot; artichoke, Chinese; artichoke, Jerusalem; canna, edible; cassava, bitter and sweet; chayote (root); chufa; dasheen (taro); ginger; leren; tanier; turmeric; yam bean; yam, true</p>	3.5 to 7 pints/ treated acre (56 to 112 fl. oz./treated acre)	<p>Application can be made at planting or pre-planting.</p> <p>Do not apply more than one application per crop and no more than 3.5 lb fluensulfone per acre per calendar year.</p>

Product name is a registered trademark of an ADAMA Group Company.

Copyright language in size 8 font in order of competitor products listed in order of company alphabetically.

EPA Reg. No.: 66222-243