

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Legacy[®] MA

Herbicide

ACTIVE CONSTITUENTS:

250 g/L MCPA

present as the ETHYL HEXYL ESTER

25 g/L DIFLUFENICAN

SOLVENT: 150 g/L N-METHYL-2-PYRROLIDONE

GROUP

FII

HERBICIDE

For the control of certain broadleaf weeds in winter cereals and clover as specified in the Directions for Use table

Formulation type

Emulsifiable Concentrate



ADAMA

adama.com

CONTENTS: 5 L, 10 L, 20 L, 110 L, 200 L, 1000 L

DIRECTIONS FOR USE

Restrictions: DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.
DO NOT apply to crops under stress due to disease or insect damage.
DO NOT apply to frost-affected crops or if frosts are imminent.
DO NOT apply when heavy rain is expected within 4 hours.

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/ha	CRITICAL COMMENTS
Cereals Wheat, Barley, Oats, Triticale, Cereal Rye (including Cereals undersown with Clover) Pasture Newly sown and established Clover-based Pasture, Clover for hay and seed production	Wild Radish	Up to the 2 leaf stage and not more than 60 mm in diameter	WA only	250 mL	CROP STAGE Cereals Up to 750 mL (3 leaf to fully tillered stage - Z13 to 30). Over 750 mL (5 leaf to fully tillered stage - Z15 to 30). Optimum results are achieved when sprayed at 3 - 5 leaf crop stage (generally 4-8 weeks post sowing). WA only: DO NOT apply to Barley or Kulin Wheat before the 5 leaf stage (Z15). Warning: LEGACY [®] MA may cause transient crop yellowing of cereals. Some varieties of oats have not been tested. (Refer to "CROP TOLERANCE" section of GENERAL INSTRUCTIONS). Clover Application is recommended prior to the eighth trifoliate leaf stage, however, applications prior to the third leaf stage may result in crop damage especially under stressed conditions and in sandy soils. DO NOT apply to Annual Medics or lucerne. Warning: LEGACY [®] MA may cause transient crop yellowing of clover, and may affect growth and seed set of some varieties of clover. (Refer to "CROP TOLERANCE" section of GENERAL INSTRUCTIONS).
		Up to the 4 leaf stage and not more than 120 mm in diameter	All States	500 mL	
		Up to the 6 leaf stage and not more than 150 mm in diameter		750 mL	
		Up to the 8 leaf stage and not more than 180 mm in diameter		1.0 L	
	Charlock, Hedge Mustard, Indian Hedge Mustard, Shepherd's Purse, Turnip Weed, Wild Turnip	Up to the 2 leaf stage and not more than 60 mm in diameter		500 mL	
		Up to the 4 leaf stage and not more than 120 mm in diameter		750 mL	
Up to the 6 leaf stage and not more than 150 mm in diameter			1.0 L		

continued overleaf

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/ha	CRITICAL COMMENTS
<p>Cereals Wheat, Barley, Oats, Triticale, Cereal Rye (including Cereals undersown with Clover)</p> <p>Pasture Newly sown and established Clover-based Pasture, Clover for hay and seed production</p>	London Rocket	Up to the 5 leaf stage and not more than 120 mm in diameter	Qld only	750 mL	<p><i>From previous page</i></p> <p>WEED STAGE Apply when weeds are actively growing. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as:</p> <ul style="list-style-type: none"> • high crop and weed density • late season germinations • abnormal weed growth (including early flowering), higher rates of product (up to the maximum rate of application specified for that weed) may be required. <p>LEGACY® MA will not effectively control:</p> <ul style="list-style-type: none"> • regrowth of suppressed weeds; • transplanted weeds; • regrowth from rhizomes or roots; • weeds growing under stress from previous herbicide applications. <p>GRAZING Efficacy on larger weeds will be improved by grazing with normal levels of stock after the 7 day withholding period. Refer to 'PROTECTION OF LIVESTOCK' for grazing precautions.</p> <p>APPLICATION Activity of this product will be reduced if weeds are stressed. Optimum results will be obtained if good soil moisture exists at and after application. Where crop or weed density is high, water volume should be increased.</p> <p>WILD RADISH LEGACY® MA will provide residual control of Wild Radish for up to 4 weeks after application. Effective residual activity of this product may be reduced where:</p> <ul style="list-style-type: none"> • rates lower than 1.0 L/ha are used; • dry conditions prevail; • poor coverage of the soil surface is achieved; • crop is planted in non-wetting sand; • soils contain a high content of organic matter. <p>Optimum results will be obtained if good soil moisture exists at and after application.</p>
	Ward's Weed		SA only		
	Capeweed	Up to the 2 leaf stage and not more than 60 mm in diameter	All States	500 mL	
				1.0 L	
	Crassula	Up to the 2 leaf stage		500 mL	
				750 mL	
	Prickly Lettuce	Up to the 2 leaf stage		500 mL	
				750 mL	
				1.0 L	
	Dense-flower Fumitory	Up to the 2 leaf stage		750 mL	
				1.0 L	
	Corn Gromwell, Saffron Thistle, Toad Rush				
	Deadnettle		NSW, Vic, SA only		
	Sorrel	Up to the 2 leaf stage	Vic only		
Canola (rape)	Up to the 4 leaf stage	All States	500 mL		
Purple Goosefoot	Up to the 6 leaf stage	Qld only			
Turnip Weed, Wild Turnip	Cotyledon to 2 leaf stage	NSW only (West of Newell Hwy.) SA only (Eyre peninsula north of the line between Venus Bay and Cowell)	350 mL		
Cereals Wheat, Barley, Oats, Triticale, Cereal Rye	Fumitory	2 - 6 leaf stage	All States	500 mL + 200 mL terbutryn (500 g/L)	
SUPPRESSION OF THE FOLLOWING WEEDS					
<p>Cereals Wheat, Barley, Oats, Triticale, Cereal Rye (including cereals undersown with clover)</p> <p>Pasture Newly sown and established Clover based Pasture, Clover for hay and seed production</p>	Saffron Thistle	Up to the 6 leaf stage	All States	1.0 L	
	Chickweed, Fireweed, Hexham Scent (King Island Melilot), Iceplant, Mouseeared Chickweed, Nightscented Stock, Paterson's Curse, Peppergrass, Skeleton Weed, Long Storksbill, Volunteer Lupins	Up to the 4 leaf stage			
	Wireweed (Hogweed)			750 mL	
	Common Sowthistle (Milk Thistle), Cowvine, Dock, Doublegee (Spiny Emex), Fat Hen, Horehound, Hyssop Loosestrife, Marshmallow, Rough Poppy, Scarlet Pimpernel, Stemless Thistle, Tree Hogweed, Variegated Thistle, Vetch (Tares)	Up to the 2 leaf stage		1.0 L	

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/Ha	CRITICAL COMMENTS
Cereals Wheat, Barley, Oats, Triticale, Cereal Rye	Wild Radish	Up to the 4 leaf stage and not more than 120 mm in diameter	All States	350 mL + 200 mL MCPA LVE (500 g/L)	<p>Refer also to all Critical Comments relating to weed stage, grazing, application and Wild Radish above.</p> <ul style="list-style-type: none"> • Reduced efficacy (suppression only) may be achieved on Wild Radish larger than 8 leaf or greater than 180 mm in diameter. <p>DO NOT use this tank-mix if cereals are undersown with lucerne or annual medics.</p> <p>Crop Stage LEGACY® MA 350 mL + MCPA LVE 200 mL: Apply from 3 leaf to fully tillered (Zadok's Z13 to Z30). LEGACY® MA 500 mL + MCPA LVE 200 mL: Apply from 3 leaf to fully tillered (Zadok's Z13 to Z30). LEGACY® MA 500 mL + MCPA LVE 400 mL: Apply from 5 leaf stage to fully tillered (Zadok's Z15 to Z30).</p> <p>Optimum results are achieved when sprayed at 3 - 5 leaf crop stage (generally 4 - 8 weeks post sowing).</p> <p>WA only: DO NOT apply to Barley or Kulin Wheat before the 5 leaf stage (Z15).</p> <p>Warning: LEGACY® MA may cause transient crop yellowing of cereals. Some varieties of oats have not been tested. (Refer to "CROP TOLERANCE" section of GENERAL INSTRUCTIONS). Observe instructions also on MCPA LVE product label.</p>
		Up to the 6 leaf stage and not more than 150 mm in diameter		500 mL + 200 mL MCPA LVE (500 g/L)	
		Up to the 8 leaf stage and not more than 180 mm in diameter*		500 mL + 400 mL MCPA LVE (500 g/L)	

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

CROP HARVEST:

ALL CROPS:

NOT REQUIRED WHEN USED AS DIRECTED.

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

WEED LIST

Common name	Scientific name	Common name	Scientific name
Canola (Rape)	<i>Brassica napus</i>	Mouse-Eared Chickweed	<i>Cerastium glomeratum</i>
Capeweed	<i>Arctotheca calendula</i>	Night-Scented Stock	<i>Matthiola longipetala</i>
Charlock	<i>Sinapis arvensis</i>	Paterson's Curse	<i>Echium plantagineum</i>
Chickweed	<i>Stellaria media</i>	Peppergrass	<i>Lepidium spp.</i>
Common Sowthistle (Milk Thistle)	<i>Sonchus oleraceus</i>	Prickly Lettuce	<i>Lactuca serriola</i>
Corn Gromwell	<i>Buglossoides arvensis</i>	Purple Goosefoot	<i>Scleroblitum atriplicinum</i>
Cowvine	<i>Ipomoea lonchophylla</i>	Rough Poppy	<i>Papaver hybridum</i>
Crassula	<i>Crassula spp.</i>	Saffron Thistle	<i>Carthamus lanatus</i>
Deadnettle	<i>Lamium amplexicaule</i>	Scarlet Pimpernel	<i>Anagallis arvensis</i>
Dense-Flower Fumitory	<i>Fumaria densiflora</i>	Shepherd's Purse	<i>Capsella bursa-pastoris</i>
Dock	<i>Rumex spp.</i>	Skeleton Weed	<i>Chondrilla juncea</i>
Doublegee (Spiny Emex)	<i>Emex australis</i>	Sorrel	<i>Rumex spp.</i>
Fat Hen	<i>Chenopodium album</i>	Stemless Thistle	<i>Onopordum acaulon</i>
Fireweed	<i>Senecio spp.</i>	Toad Rush	<i>Juncus bufonius</i>
Fumitory	<i>Fumaria spp.</i>	Tree Hogweed	<i>Polygonum patulum</i>
Hedge Mustard	<i>Sisymbrium officinale</i>	Turnip Weed	<i>Rapistrum rugosum</i>
Hexham Scent (King Island Melilot)	<i>Melilotus indicus</i>	Variagated Thistle	<i>Silybum marianum</i>
Horehound	<i>Marrubium vulgare</i>	Vetch (Tares)	<i>Vicia sativa</i>
Hyssop Loosestrife	<i>Lythrum hyssopifolia</i>	Volunteer Lupins	<i>Lupinus spp.</i>
Iceplant	<i>Mesembryanthemum spp.</i>	Ward's Weed	<i>Carrichtera annua</i>
Indian Hedge Mustard	<i>Sisymbrium orientale</i>	Wild Radish	<i>Raphanus raphanistrum</i>
London Rocket	<i>Sisymbrium irio</i>	Wild Turnip	<i>Brassica tournefortii</i>
Long Storksbill	<i>Erodium botrys</i>	Wireweed (Hogweed)	<i>Polygonum aviculare</i>
Marshmallow	<i>Malva parviflora</i>		

GENERAL INSTRUCTIONS

RESISTANT WEEDS WARNING

LEGACY® MA Herbicide is a member of the phenoxy and nicotinilide groups of herbicides and acts by inhibiting carotenoid biosynthesis at the phytoene desaturase step (PDS inhibitors) and disrupting plant cell growth. For weed resistance management LEGACY® MA is both a Group F and a Group I herbicide. Some naturally occurring weed biotypes resistant to LEGACY® MA and other Group F and I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by LEGACY® MA or other Group F or Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Adama Australia accepts no liability for any losses that may result from the failure of LEGACY® MA to control resistant weeds.

TOLERANCE

Some pre-emergence herbicides, such as atrazine, can cause stress to certain crops resulting in an increase in crop damage when using this product. Subterranean clover is particularly sensitive.

CEREALS

After application some transient crop yellowing may occur. This usually appears as yellow or white banding on leaves. Provided the crop is not under stress from pre-emergent herbicide, root disease, insect damage, frost, dry or excessively moist conditions, the development of the crop and subsequent growth will be unaffected.

WARNING (Oats)

The tolerance of oat varieties Esk and Nile (the two main varieties grown in Tasmania) to LEGACY® MA has not been tested. Test a small area of crop before using LEGACY® MA over large areas. Consult your local Adama representative for advice on specific varieties.

PASTURE

The tolerance of clover varieties to LEGACY® MA can vary with rate of application, soil type, crop health, stage of growth and degree of moisture and temperature stress.

WARNING

LEGACY® MA may result in transient crop yellowing and suppression of growth with a resultant initial reduction in dry matter, particularly at rates in excess of 500 mL/ha and in areas of double spray. For this reason we recommend application prior to the 8 trifoliate leaf stage. However, at the lower rates (500 mL/ha and less) and under normal growing conditions, subsequent growth and seed yield should not be affected. Under normal growing conditions, the following varieties have shown acceptable levels of foliage tolerance to LEGACY® MA applied at 500 mL/ha:

ArrowLeaf: Zulu

Balansa: Paradana

Berseem: Sacromonte

Persian: Kyambro, Lupers, Maral

White: Haifa

Subterranean Clover: Daliak, Dalkeith, Denmark, Esperance, Geraldton, Goulburn, Karridale, Larissa, Leura, Mt.Barker, Nungarin, Rosedale, Seaton Park, Trikkala and Woogenellup.

The effects of LEGACY® MA on clover seed yield have been tested on the following varieties. Under normal growing conditions they show acceptable levels of tolerance to LEGACY® MA applied at 500 mL/ha:

Subterranean Clover: Esperance, Goulburn, Larissa, Seaton Park and Trikkala.

WARNING

Rose and Strawberry clover have shown increased sensitivity to LEGACY® MA. LEGACY® MA may affect the seed yield of Subterranean Clover variety Woogenellup. Some pasture grasses, including Phalaris and Cocksfoot, may show some initial reduction in vegetative growth after application of LEGACY® MA. Care should be exercised if sensitive clover varieties or grasses are included in the pasture sward. Varieties not listed should be tested before using LEGACY® MA over large areas. Consult your local Adama representative for advice on specific varieties.

SUBSEQUENT CROPS

To reduce effect on subsequent susceptible crops (e.g. canola), ensure thorough cultivation of soil prior to the sowing of these crops.

MIXING

To ensure even mixing, half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly then add the remainder of the water. Agitate again before spraying commences. Reseal part-used product container immediately after use. Spray mixtures containing LEGACY® MA should not be left to stand overnight. Prolonged periods of exposure to cold temperatures could result in settling out of the product in the mixture.

WARNING

The rubber components present in some spraying units may be affected by exposure to the solvents in LEGACY® MA and some other agricultural products. To reduce this risk it is recommended that the spray unit be thoroughly washed with a boom cleaner and fresh water after use. Contact the spray unit manufacturer to determine the suitability of the rubber components for use with agricultural products.

APPLICATION

Boom Sprayer: A minimum of 50 L of water per hectare should be used, however, for optimum results water rates of 70-100 L/ha are recommended. Increase the water volume if weed infestation is heavy or crop cover is dense. Complete coverage of weeds is essential.

Aircraft (NSW, Vic, SA only): Apply in a minimum of 30 L water per hectare. Effective weed control will only be achieved where good coverage of leaf surface is achieved.

COMPATIBILITY

The following products are physically compatible with LEGACY® MA as a two-way mixture in the spray tank but should only be used for the crops specified:

Crop	LEGACY® MA	Compatible Product
Wheat, Triticale and Cereal Rye only	Up to 750 mL/ha	Rhino® (also barley), fenoxaprop-p-ethyl (Wild Oats only),
Cereals (including undersown)	All rates	Chlorpyrifos (500 g/L)
Cereals (not undersown)	Up to 500 mL/ha	Lynx® WG, Tackle® WG, MCPA LVE, Lonestar® 750WG
	All rates	Bronco® 200, 2,4-D Amine 500 Herbicide, Victory®, Enforcer® 242, Cutlass® (up to 115 g only), Eclipse*
Wheat, Barley, Triticale, and Cereal Rye only (not undersown)		Pentagon®
Wheat only (not undersown)		Mandate®
Clover	Up to 750 mL/ha	Leopard®, Resilience®
Subterranean Clover		Simazine (500 g/L), Simazine (500 g/L) + paraquat (200 g/L) mixture
	Up to 1.0 L/ha	2,4-DB amine (500 g/L)

When mixing with other herbicides, crop yellowing may be enhanced. When mixing with Rhino® or fenoxaprop-p-ethyl some reduction in the efficacy and speed of action of these products may occur. If the crop is stressed, the application of the herbicide tank-mixtures may cause yield reduction. When mixing with Cutlass® a temporary wilting may be evident in some crops after application. Growers should seek advice before spraying recently released cereal varieties.

Use the recommended rates for both herbicides in the tank-mixture as well as the surfactant recommendation of the grass herbicide. If another herbicide is applied as a tank mix, observe the plantback restrictions on that label. DO NOT add surfactant when mixing LEGACY® MA and Lynx.

Simazine: Refer to the simazine label for correct application rates, especially with regard to soil types. This product may be mixed in the spray tank with one of the following insecticides according to the directions for use on this product: Alpha-Scud® Elite, Ballistic® Elite, Exonerate®, lambda-cyhalothrin and Venom®.

WARNING

DO NOT use crop oils with LEGACY® MA or LEGACY® MA tank mixtures with other products in cereals. As formulations of other manufacturer's products are beyond the control of Adama, all mixtures should be tested prior to mixing commercial quantities.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

PROTECTION OF LIVESTOCK

Grazing Precaution: Sprayed weeds may become more palatable to stock and a higher intake of some weeds may result in stock poisoning and death from causes such as nitrate poisoning. Care should be taken especially where Capeweed, Paterson's Curse and variegated thistles predominate in the pasture. Avoid grazing with young or breeding stock. DO NOT graze horses or pigs on Paterson's Curse. If in doubt, contact your nearest Department of Agriculture.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT
Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

5 L, 10 L, 20 L, 200 L: Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or a designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

Envirodrum Micro Matic Valve (110 L)

Store the original sealed Envirodrum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the Envirodrum to the point of purchase. The Envirodrum remains the property of Adama Australia.

1000 L

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs Adama Australia should be advised immediately. This minibulk container is reusable and remains the property of Adama Australia. DO NOT rinse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return to Adama Australia for cleaning, relabelling and refilling.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Will irritate the skin. Avoid contact with eyes and skin. DO NOT inhale vapour. When opening the container, preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, DO NOT induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for LEGACY® MA is available from adama.com or call Customer Service on 1800 423 262.

CONDITIONS OF SALE: The use of LEGACY® MA Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Adama Australia regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Adama Australia accepts no responsibility for any consequence whatsoever resulting from the use of this product.

® Registered trademarks of an Adama Group Company

* Other trademarks

NOT A DANGEROUS GOOD ACCORDING TO THE AUSTRALIAN DANGEROUS GOODS (ADG) CODE.

10/2018 10173