

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

## Quali-Pro Negate®

### Herbicide

ACTIVE CONSTITUENTS: **167 g/kg RIMSULFURON**  
**200 g/kg METSULFURON-METHYL**

**GROUP B HERBICIDE**

**Situation:** Turf; Couch, Couch hybrids, Zoysia, Qld Blue Couch.

**Controls:** Winter grass, Ryegrass, Onion grass and certain broadleaf weeds as specified in the Directions for Use table.

Formulation type  
**Water Dispersible  
Granule**

**WG**



**ADAMA**

adama.com

NET CONTENTS: 110 g

#### **DIRECTIONS FOR USE**

##### **RESTRAINTS**

For PROFESSIONAL use only.

DO NOT apply product more than 5 (five) times per year to child play areas (e.g. lawns and ovals).

DO NOT apply by aircraft.

DO NOT apply through any type of irrigation equipment.

##### **SPRAY DRIFT RESTRAINTS**

Specific definitions for terms used in this section of the label can be found at [apvma.gov.au/spraydrift](http://apvma.gov.au/spraydrift).

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

- Spray droplets are not smaller than a MEDIUM spray droplet size category.
- Minimum distances between the application site and downwind sensitive areas are observed (see 'Mandatory downwind buffer zones' section of the following table titled 'Buffer zones for boom sprayers').

## Buffer Zones for Boom Sprayers

Application Rate	Boom Height Above the Target Canopy	Mandatory Downwind Buffer Zones				
		Bystander Areas	Natural Aquatic Areas	Pollinator Areas	Vegetation Areas	Livestock Areas
Up to maximum label rate	0.5 m or lower	Not required	100 metres	Not required	40 metres	Not required
	1.0 m or lower	Not required	230 metres	Not required	100 metres	Not required

SITUATION	WEEDS CONTROLLED	RATE / Ha	CRITICAL COMMENTS
Turf  Common Couch, Hybrid Couch, Zoysia, Qld Blue Couch	Winter Grass ( <i>Poa annua</i> )	110 g/ha in 200-500 L of water per hectare	Apply to actively growing weeds and not to weeds under stress.  Avoid exceeding application rates through overlapping sprayed areas.  Safety screening may not have been conducted on all new couch varieties and it is therefore recommended that small areas be tested for turf safety before largescale application occurs.
	Ryegrass ( <i>Lolium perenne</i> )		
	Onion Grass ( <i>Romulea rosea</i> )		
	Broadleaf Weeds including; Bindii ( <i>Soliva sessilis</i> ), Caustic Weed ( <i>Chamaesyce prostrata</i> ), White Clover ( <i>Trifolium repens</i> ), Cobblers Peg ( <i>Bidens pilosa</i> ), Creeping Indigo ( <i>Indigofera spicata</i> ), Creeping Oxalis ( <i>Oxalis</i> spp.), Plantain ( <i>Plantago lanceolata</i> ), Tridax Daisy ( <i>Tridax procumbens</i> )		

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

### WITHHOLDING PERIOD

DO NOT graze treated turf or lawn; or feed turf or lawn clippings from any treated area to poultry or livestock.

### GENERAL INSTRUCTIONS

QUALI-PRO NEGATE® contains Rimsulfuron and Metsulfuron-Methyl herbicides which belong to the sulfonyleurea chemical family of herbicides. QUALI-PRO NEGATE® is absorbed through foliage of plants, rapidly inhibiting the growth of susceptible weeds. One to three weeks after post emergence application to weeds, leaves of susceptible plants appear chlorotic, and the growing point subsequently dies. In warm, moist conditions, the expression of herbicide symptoms is accelerated; in cold, dry conditions, expression of herbicide symptoms is delayed. It is necessary for the spray material to dry on the foliage to be sufficiently absorbed. QUALI-PRO NEGATE is rainfast in 4 hours.

### Mixing

QUALI-PRO NEGATE® is a physical blend of rimsulfuron and metsulfuron-methyl herbicide. One 110 g container treats 1 ha. It is recommended to use the entire contents of the container

- 1) Half fill the spray tank with water
- 2) While agitating, add the required amount of QUALI-PRO NEGATE®
- 3) Continue agitation until the QUALI-PRO NEGATE® is fully dispersed, at least 5 minutes
- 4) Maintain agitation and continue filling tank with water.

## RESISTANT WEEDS WARNING

QUALI-PRO NEGATE® Herbicide is a member of the Sulfonylurea group of herbicides. QUALI-PRO NEGATE® has the inhibitor of the enzyme acetolactate synthase (ALS) mode of action. For weed resistance management QUALI-PRO NEGATE® is a Group B herbicide. Some naturally occurring weed biotypes resistant to QUALI-PRO NEGATE® and other Group B herbicides are known to exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by QUALI-PRO NEGATE® or other Group B herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Adama Australia accepts no liability for any losses that may result from the failure of QUALI-PRO NEGATE® to control resistant weeds.

GROUP	<b>B</b>	HERBICIDE
-------	----------	-----------

## RE-ENTRY

DO NOT enter treated areas until spray has dried. If prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

This product may cause injury to or loss of trees and other vegetation, including aquatic plants. DO NOT apply or drain or flush equipment on or near native or non-target trees or other plants or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

## STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool well ventilated area out of direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

## SAFETY DIRECTIONS

Harmful if inhaled. Will irritate the eyes. When opening the container and preparing the spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow length chemical resistant gloves, goggles and disposable dust mask covering mouth and nose. Wash hands after use. After each day's use wash gloves, goggles and contaminated clothing.

## FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131 126; New Zealand 0800 764 766.

## WARNING

CONTAINS FORMALDEHYDE

## SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for QUALI-PRO NEGATE® is available from [adama.com](http://adama.com) or call Customer Service on 1800 423 262.

## CONDITIONS OF SALE

The use of QUALI-PRO NEGATE® HERBICIDE being beyond the control of the manufacturer, no warranty expressed or implied is given by Adama Australia regarding its suitability, fitness or efficiency for any purposes for which it is used by the buyer, whether in accordance with the Directions for Use or not. Adama Australia accepts no responsibility for any consequence whatsoever resulting from the use of this product.

® Registered trademark of an Adama Group Company.

07/2020 22013