

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# Qualipro I-pro 365

Fungicide

ACTIVE CONSTITUENT: **365 g/L IPRODIONE**

GROUP **2** FUNGICIDE

A suspo-emulsion formulation with improved rainfastness (over other water based formulations) for the control of fungal diseases in recreational turf as specified in the directions for use



ADAMA

Formulation type  
Suspo-emulsion **SE**

adama.com

CONTENTS: 5 - 20 L

## DIRECTIONS FOR USE

SITUATION	PEST	RATE	CRITICAL COMMENTS
Recreational Turf	Brown Patch ( <i>Rhizoctonia solani</i> )	80 - 125 mL/4 - 150 L of water/100 m <sup>2</sup>	<b>Curative Program:</b> Repeat treatment after 14 days. Use the higher rate under conditions of severe disease pressure. <b>For preventative treatment:</b> Apply as a monthly spray program. Repeat application at monthly intervals from February to May. This use is subject to an AFRAC anti-resistance strategy. DO NOT apply more than two consecutive sprays of QUALIPRO I-PRO 365 Fungicide or related dicarboximide fungicides unless tank mixed with a fungicide from a different activity group.
	Dollar Spot ( <i>Sclerotinia homoeocarpa</i> )	80 mL/4 - 110 L of water/100 m <sup>2</sup>	<b>For curative treatment:</b> Repeat treatment in 14 days. <b>For preventative treatment:</b> Apply monthly. Repeat application at monthly intervals from February to May. This use is subject to an AFRAC anti-resistance strategy. DO NOT apply more than two consecutive sprays of QUALIPRO I-PRO 365 Fungicide or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.
	Winter Fusarium or Fusarium patch ( <i>Fusarium rivale</i> ) (NOT Qld)	125 mL/4 - 150 L of water/100 m <sup>2</sup>	
	Helminthosporium disease affecting leaf, crown and roots: (Black and white Helminthosporium spot <i>Bipolaris</i> spp. <i>Drechslera</i> spp. <i>Exserohilum</i> spp.)	65 mL/4 - 150 L of water/100 m <sup>2</sup>	<b>For curative treatment:</b> Apply as a foliar spray and repeat application after 7 - 14 days. For white spot symptoms on couch grass ( <i>Cynodon dactylon</i> ) only use higher water volumes of 100 - 150 L of water per 100m <sup>2</sup> . <b>For preventative treatment:</b> Apply as a high volume foliar spray. Repeat application at monthly intervals from February to May. This use is subject to an AFRAC anti-resistance strategy. DO NOT apply more than two consecutive sprays of QUALIPRO I-PRO 365 Fungicide or related dicarboximide fungicides, unless tank mixed with a fungicide from a different activity group.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

## GENERAL INSTRUCTIONS

NOTE QUALIPRO I-PRO 365 Fungicide may be very unstable in conditions where the pH is 7 or higher. It is therefore essential to check the pH of the spray solution before adding QUALIPRO I-PRO 365 Fungicide.

A suitable registered buffering agent may have to be added to bring pH down below 7.

## APPLICATION

Good control requires good coverage. Application should be made using sufficient water to ensure thorough coverage.

## TURF NUTRITION

This product acts very rapidly and where the turf is of a low nutritional status, a slight yellowing may be seen. This will be corrected by a nitrogenous fertiliser application. Where possible, it is recommended that the fertiliser program should precede the fungicide application.

## COMPATIBILITY

The use of a surfactant or spray oil is not recommended with QUALIPRO I-PRO 365 Fungicide as it may result in damage to sensitive plants.

DO NOT mix with fertilisers. Mixtures with some fertilisers, e.g. urea, may cause foliar damage. As formulations of other manufacturers' products are beyond the control of Adama Australia, all mixtures should be tested prior to mixing commercial quantities.

## FUNGICIDE RESISTANCE WARNING

For fungicide resistance management

QUALIPRO I-PRO 365 Fungicide is a Group

GROUP	<b>2</b>	FUNGICIDE
-------	----------	-----------

2 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 2 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by the product and other Group 2 fungicides, thus resulting in a reduction in efficacy. Since the occurrence of resistant fungi is difficult to detect prior to use, Adama Australia accepts no liability for any losses that may result from the failure of the product to control resistant fungi.

## PRECAUTION

DO NOT GRAZE TREATED AREAS OR FEED TURF CLIPPINGS FROM TREATED AREAS TO ANIMALS INCLUDING POULTRY.

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

## STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal at an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product must not be burnt.

## SAFETY DIRECTIONS:

May irritate the eyes. Avoid contact with the eyes. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and water-proof gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

## FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

## SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for QUALIPRO I-PRO 365 Fungicide is available from adama.com or call Customer Service on 1800 423 262.

**CONDITIONS OF SALE:** The use of QUALIPRO I-PRO 365 Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Adama Australia regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Adama Australia accepts no responsibility for any consequence whatsoever resulting from the use of this product.

®Registered trademark of an Adama Group Company

06/2017 9089