



ADAMA

Custodia® Forte

Fungicide

Product overview

Custodia® Forte fungicide provides protectant and systemic activity against the key diseases in almonds, grapes and macadamias. Custodia Forte is a concentrated suspension concentrate formulation that combines two different modes of action (Group 3 + 11) for robust disease control and resistance management.

Key features

- Preventative control or suppression of key diseases protecting growers yield potential and quality
- Two modes of action for improved disease control and resistance management
- Excellent protectant and systemic activity, suitable to use in IPM programs
- Suspension concentrate formulation with low odour
- Highly compatible formulation for tank mixing

Harvest Withholding Period

Almonds: Not required when used as directed

Grapevines: Do not harvest for 4 weeks after application

Macadamias: Do not harvest for 15 days after application

Product information

Active ingredient:

Azoxystrobin 222 g/L +
Tebuconazole 370 g/L

Mode of action:

Group 3 + 11

Formulation:

Suspension Concentrate

Registered crops:

Almonds, macadamias, grapes

Target diseases:

Brown rot/Blossom blight,
Shot-hole, Rust, Hull rot,
Powdery mildew, Downy mildew,
Botrytis bunch rot, Husk spot

Re-entry period:

Once the spray has dried

Custodia® Forte

Crop	Target disease	Application rate	Timing	Spray frequency
Almonds	Brown Rot/Blossom blight and Shot-hole	32 mL/100 L or 540 mL/ha	From early (10%) bloom before early hull split	Apply in rotation with fungicides from alternative modes of action on a 7 to 14-day interval targeting key growth stages including full bloom, petal fall and shuck fall After shuck fall, apply Cavalry® Weatherguard for Shot-hole control on a 14-day spray interval Do not apply consecutive applications and no more than three sprays per season
	Rust		Commence spray program from late flowering and/or when first new leaves emerge	Apply up to two applications per season in rotation with fungicides from a different mode of action group on a 14-day spray interval
	Hull rot (suppression only)		Early hull split	Apply a single application. If a second fungicide application is required, rotate with an alternative registered fungicide after a 10 to 14-day interval
Grapes	Powdery Mildew	35 mL/100 L	Before E-L 19 (before flowering) and under low disease pressure up to E-L 29	Maximum two applications per season. Always apply as a preventative spray and never as a curative spray
			From after E-L 19 (start of flowering) and under high disease pressure up to E-L 29	
	Downy Mildew	54 mL/100 L or 540 mL/ha	Seasonal preventative spray program commencing when shoots are 20 cm long, up to E-L 29	Apply in rotation with fungicides from other modes of action on a 7 to 21-day spray interval. Under high disease pressure and during flowering, apply in rotation with Axiom® Plus
Botrytis Bunch Rot (low disease pressure only)	Use no later than E-L 29		If three or fewer Botrytis sprays are applied in a season, use no more than one Custodia Forte spray during the season If four or more Botrytis sprays are to be applied in a season, up to two Custodia Forte sprays can be applied	
Macadamia	Husk spot	32 mL/100 L*	From match head timing	Two applications per season on a 14 to 28-day spray interval dependent on disease pressure, with fungicides from other modes of action groups.

**Applications of Custodia Forte with Hasten* as per label guidelines can improve husk spot control under conditions of high disease pressure. However, we advise growers to evaluate this mix in a small area before commercial use*



Scan here for more information

Refer to label for specific application recommendations.

™Trademarks or ®Registered trademarks of an ADAMA Agricultural Solutions Ltd Company.

*Registered trademarks. Please note: This information is not intended to replace the product label.

Always read the complete product label appearing on the container before opening or using products. ADA24448



FUNGICIDE

ADAMA.COM

1800 4 ADAMA

