



ADAMA

# Custodia®

## A new solution for managing husk spot.

### About Custodia

Custodia provides reliable efficacy on husk spot, reducing premature nut abscission and protecting yield potential and quality. It contains 120 g/L azoxystrobin and 200 g/L tebuconazole in a suspension concentrate formulation for improved compatibility and reduced odour.

### Two modes of action

Custodia contains two different modes of action for improved disease control and resistance management. Azoxystrobin (Group 11) is a strobilurin fungicide, which inhibits mitochondrial respiration and disrupts energy production in fungi. Tebuconazole (Group 3) is a DMI fungicide, which inhibits ergosterol biosynthesis and cell wall production and mycelial/hyphal development in fungi. Custodia has excellent protectant activity and targets fungi at different stages of the infection process.


### At a glance

<b>Robust protection</b>	Custodia provides reliable efficacy on husk spot, reducing premature nut abscission and protecting yield potential and quality.
<b>Two modes of action</b>	Custodia contains two different modes of action (Group 3 and 11) for improved disease control and resistance management.
<b>Protectant and systemic activity</b>	Custodia has protectant and systemic activity, protecting crops at multiple stages of the fungal infection process.
<b>Unique formulation</b>	Custodia contains 120 g/L azoxystrobin and 200 g/L tebuconazole in a suspension concentrate for improved compatibility and reduced odour.

### Tebuconazole

Inhibits ergosterol biosynthesis, disrupting cell wall production and mycelial growth and development in target fungi.

Rapidly absorbed into leaves and green tissue and transported acropetally.



### Azoxystrobin

Disrupts respiration and inhibits energy production.

Inhibits spore germination and host penetration during the early stages of fungal vegetative growth.

Extended residual activity on spore germination.

# Using Custodia®

## Application rate

Apply Custodia at 60 mL/100 L as a dilute application to the point of runoff. For advice on concentrate spraying, please contact your local Adama representative.

## Application timing





Apply up to two applications of Custodia as part of a preventative spray program for husk spot commencing at the match head timing. Use Custodia in rotation with other mode of action fungicides on a 14–28 day spray interval. Use the shorter interval under conditions where higher disease pressure is expected.

## Withholding period

DO NOT harvest for 15 days after application.

## Re-entry interval

Workers may re-enter areas sprayed with Custodia once the spray has dried.



Scan here for more information

Simply. Grow. Together.



ADAMA

Visit [adama.com](https://adama.com)

Phone 1800 4 ADAMA

Follow us  