

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Victory®

Herbicide

ACTIVE CONSTITUENT: **300 g/L CLOPYRALID**

present as the TRIISOPROPANOLAMINE SALT

GROUP

HERBICIDE

For the control of a wide range of broadleaf weeds in wheat, barley, oats, triticale, canola, pastures, fallow land, forests and industrial situations as specified in the Directions for Use table

Formulation type
Soluble
Concentrate

SL



ADAMA

adama.com

CONTENTS: 5 L, 10 L, 20 L, 110 L, 1000 L

DIRECTIONS FOR USE

1. CROPS, PASTURES AND FALLOW LAND

CROP	CROP STAGE	WEED	WEED STAGE	RATE mL/ha	STATE	CRITICAL COMMENTS
Barley, Oats, Triticale, Wheat	Pre-sowing	Capeweed	Up to 8 leaf and maximum 10 cm diameter	150 + knockdown herbicide	WA only	Pre-sowing: This rate should only be used in tank mixture with formulations of paraquat/diquat or glyphosate.
	Post-sowing Pre-emergence through to 3 leaf			150 + diuron at 300 mL/ha		Post-sowing pre-emergent to 3 leaf: This rate should only be used in tank mixture with diuron for control of transplants.
				Early post-emergence (2 leaf to jointing)		150
	4 - 5 leaf onwards	Capeweed, Soldier Thistle	Up to 10 cm diameter (4 - 8 leaf)	300	NSW, ACT, Vic, Tas, SA, WA only	Weeds should be young and actively growing. Weeds will become stunted and will not be competitive soon after application although final results may not show for some weeks. Faba Beans and Lupins will only be suppressed. Use 75 mL/ha rate only in combination with MCPA LVE.
		Volunteer Chickpeas, Lentils, Safflower	Up to 6 leaf	250		
		Volunteer Faba Beans, Lupins	Up to 4 leaf			
		Volunteer Field Peas	Maximum 10 cm high or 6 nodes	150 75 + 700 mL/ha MCPA LVE		
		Volunteer Medics, Seedling Lucerne	Up to 8 leaf	150		
		Volunteer Sub-clover	Up to 6 leaf			
Volunteer Vetch	Runners up to 10 cm maximum 16 leaf	100 75 + 700 mL/ha MCPA LVE				
Prickly Lettuce	4 - 6 leaf and maximum 8 cm diameter		150 + 700 mL/ha MCPA LVE			

CROP	CROP STAGE	WEED	WEED STAGE	RATE mL/ha	STATE	CRITICAL COMMENTS
Barley, Oats, Triticale, Wheat <i>continued</i>	4 - 5 leaf onwards	Thistles including: Nodding, Saffron, Scotch, Slender, Spear, Stemless, Variegated	Rosettes up to 10 cm maximum diameter	50 + 1.0 L/ha MCPA amine (500 g/L) or 50 + 700 mL/ha MCPA LVE	Qld, NSW, ACT, Vic, Tas, SA, WA only	Weeds should be young and actively growing. Weeds will become stunted and will not be competitive soon after application although final results may not show for some weeks.
	5 leaf to late tillering	Skeleton Weed	5 - 15 cm rosettes	500 + 1.0 L/ha MCPA amine (500 g/L)	NSW, ACT, Vic, SA, WA only	Weeds should be a minimum 5 cm in diameter, and growing actively. This rate will give control until harvest and substantially reduce weed numbers the following season.
Canola	2 - 8 leaf	Capeweed, Saffron Thistle, Skeleton Weed, Soldier Thistle	Up to 10 cm diameter (4 - 8 leaf)	300	Qld, NSW, ACT, Vic, Tas, SA, WA only	Weeds should be young and actively growing. Weeds will become stunted and will not be competitive soon after application although final results may not show for some weeks. Faba Beans and Lupins will only be suppressed. For the control of annual grasses, VICTORY® may be tank mixed with Verdict® Herbicide.
		Volunteer Chickpeas, Lentils, Safflower	Up to 6 leaf	250		
		Volunteer Faba Beans, Lupins	Up to 4 leaf	150		
		Volunteer Field Peas	Maximum 10 cm high or 6 nodes			
		Volunteer Medics, Seedling Lucerne	Up to 8 leaf			
		Volunteer Sub-clover	Up to 6 leaf			
		Volunteer Vetch	Runners up to 10 cm maximum 16 leaf	100		
Pastures and Fallow Land	Post-emergence	Hardhead Thistle (creeping Knapweed, Russian Knapweed)	Actively growing plants	Hand gun: 500 mL/100 L of water Boom spray: 2 or 4 L/ha	Vic only	NOTE: DO NOT USE ON LUCERNE. CLOVERS AND MEDICS WILL BE ELIMINATED FOR AT LEAST ONE YEAR. Victoria only: Use the lower rate only on light soils (sand and sandy loam) where a slightly lower degree of control is acceptable. Use the higher rate on all soil types where complete control is required. Addition of a wetting agent at label rates is recommended of Hardhead Thistle. Spray between September and April on actively growing plants for effective control. Thorough coverage is essential. Apply in 200 to 250 L of water/ha. Boom Spraying: Use the higher rates of VICTORY® plus MCPA on multi-crowned plants or rosettes larger than 30 cm in diameter. Spraying may be done at any time during active growth, usually in early winter or spring. Avoid spraying during the dormant winter period or at any time when thistles are not actively growing. DO NOT spray flowering thistles. Pre-spray Management: The pasture should be slightly grazed prior to spraying to reduce clover and grass cover and expose the smaller thistles to the spray. The grazed pasture should be left seven days to allow thistles to freshen prior to treatment. Post-treatment Management: Response of thistles to treatment with the VICTORY® plus MCPA mixture will be slow compared to the standard treatments with 2,4-D or MCPA. If possible delay grazing of sprayed thistles for 14 days after treatment. Clover Damage: The VICTORY® plus MCPA mixture can be very damaging to Subterranean Clover. The lower rate is no more damaging than label rates of 2,4-D or MCPA. The higher rate of the VICTORY® plus MCPA mixture will reduce the clover component of the pasture for about two months. Clover recovery will be quicker during periods of active growth. Hand gun (Spot spray): Treat from rosette stage to early flowering. Thorough spraying is necessary. Drench gun: Apply 10 mL to rosette crown. To multi-crown plants, apply 10 mL to each crown.
				Hand gun: 500 mL/100 L of water Boom spray: 4 L/ha	Qld only	
		Thistles including: Nodding, Scotch, Saffron, Slender, Spear, Variegated	Rosette stage prior to stem elongation. Treat rosette stage prior to stem elongation.	50 or 70 mL/ha + 1 - 1.5 L MCPA amine (500 g/L)/ha Drench gun: 50 mL/1 L of water Hand gun: 250 mL/100 L of water	Qld, NSW, ACT, Vic, Tas, SA only	

CROP	CROP STAGE	WEED	WEED STAGE	RATE	STATE	CRITICAL COMMENTS
Pastures and Fallow Land <i>continued</i>	Post-emergence	Variegated Thistle	Rosette stage prior to stem elongation. Treat rosette stage prior to stem elongation.	100 mL/ha 50 ¹ or 70 mL/ha + 400 mL/ha MCPA amine (500 g/L)	WA only	Apply the 100 mL rate in areas where Restricted Spraying Regulations prohibit the use of phenoxy herbicides. ¹ Use the lower rate on rosettes smaller than 15 cm. Subterranean Clovers will be damaged by these treatments.
		Nodding Thistle	Rosettes up to 20 cm diameter	100 mL/ha	NSW, ACT only	Apply the spray from September to October. Apply by boom spray only. DO NOT apply to thistles over 20 cm in diameter. When thistles are over 20 cm in diameter use VICTORY [®] plus MCPA (referred to above). Clover Damage: Damage to white clover will be no greater than damage with MCPA alone and less than damage from VICTORY [®] plus MCPA mixtures. Damage to Sub-clover may be greater than with MCPA or 2,4-D alone. DO NOT use for spot treatment.
		Californian Thistle	From early buds to flowering (December to February)	Hand gun: 250 mL/100 L of water Boom spray: 2 L/ha	Vic, Tas only	Addition of a wetting agent at label rates is recommended. Retreatment of regrowth in the year following treatment will usually be necessary to achieve a high level of control. Note: Clovers and Medics will be eliminated for at least one year.
Pasture, Forests, Rights-of-Way and Industrial Situations	Post-emergence	Groundsel Bush	Young seedlings to mature plants	Hand gun: 330 or 500 mL/100 L of water	Qld, NSW, ACT only	Spray foliage when growth is active. Use the lower rate on young seedlings and the higher rate on plants more than 2 m tall or when growth is slow.
Pasture	Post-emergence	Silver Wattle	Active growth Spring to Summer	Hand gun: 500 mL/100 L of water Boom and aerial spray: 5 L/ha (weeds 30 cm - 2 m) 7.0 L/ha (weeds 2 - 4 m) 8.5 L/ha (weeds 4 - 8 m)	NSW, ACT, Vic, Tas only	For effective control apply when bushes are growing actively. Large trees will not show complete necrosis. Hand gun: Means high volume NOT low volume knapsack. Spray to give full coverage of leaves and stems. Add organosilicone surfactant at 200 mL/100 L for optimum results. See General Instructions for hand gun spraying. Boom and Aerial spraying: For boom spraying in 150 to 200 L of water/ha. For aerial treatment apply in a minimum of 50 L/ha of water containing 25 - 50 % by volume of anti-evaporant oil such as SACOA ANTIEVAP Spray Adjuvant [†] . Mix VICTORY [®] and water first and then add SACOA ANTIEVAP Spray Adjuvant [†] . Maintain continuous agitation. Clovers and other legumes will be eliminated for at least one year.

2. FORESTS

DIRECTIONS FOR USE: It is essential to select a rate appropriate to weed size. Best results will be obtained when weeds are actively growing at treatment.

RESTRAINTS: DO NOT apply to weeds which may be stressed (inactive growth) due to prolonged periods of extreme heat or cold, moisture stress (water logging or drought) or previous herbicide treatment as reduced levels of control may result. DO NOT spray if rain is likely within 3 hours.

Pre-Planting: Boom and Aerial Application

FORESTS AND PLANTATION TREES INCLUDING <i>EUCALYPTUS</i> SPP., <i>CORYMBIA MACULATA</i> AND <i>PINUS RADIATA</i>				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ha	CRITICAL COMMENTS
Capeweed, Thistles, Volunteer, Legumes, Flatweed, Fleabanes	Pre-emergent	All States	2 - 6 L	Use the higher rate for extended pre-emergence control (> 3 months).
<i>PINUS RADIATA</i> ONLY				
Silver Wattle (suppression only)	Pre-emergence from seeds	NSW, ACT, Vic, Tas, SA only	6 L	For best results apply VICTORY [®] to bare soil just prior to Spring rain or when wattles are expected to germinate. Avoid application to heavy trash situations. A high level of suppression may not be achieved where rain does not fall for an extended period after application (> 1 month), or where very high rainfall occurs after application (>1200 mm/yr).

Post-Planting: High Volume Spraying by Hand Gun

FORESTS AND PLANTATION TREES INCLUDING <i>EUCALYPTUS</i> SPP., <i>CORYMBIA MACULATA</i> AND <i>PINUS RADIATA</i>				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE	CRITICAL COMMENTS
Groundsel Bush	Young seedlings to mature plants	Qld, NSW, ACT only	330 - 500 mL/100 L of water	Spray foliage when growth is active. Use the lower rate on young seedlings and the higher rate on plants more than 2 m tall or when growth is slow.
Ragwort	Actively growing rosettes up to stem elongation and before flowering	All States	200 - 300 mL/100 L of water	Spray from the rosette to the shooting stage of growth. Use the higher rate on large multi-crown plants. Addition of a 100 % non-ionic surfactant such as Wetspray® 1000 at 0.1 % v/v is recommended. Add diquat (200 g/L) at 1 L/100 L water plus a surfactant after opening of the first flowers, to prevent the formation of viable seed. Where diquat is added use a directed spray to avoid tree injury.
Silver Wattle	Active growth Spring to Summer	NSW, ACT, Vic, Tas, SA only	500 mL/100 L of water	For effective control apply when bushes are growing actively. Large trees will not show complete necrosis. Hand gun: Means high volume NOT low volume knapsack. (See General Instructions, Application). Spray to the point of run-off to give full coverage of leaves and stems. Add organosilicone surfactant (e.g. Pulse ¹) at 200 mL/100 L for optimum results.
Cape Ivy	Any growth stage	Vic, Tas only	3.3 L/ha	Application may be made at any time of the year providing foliage is dry at the time. Avoid spraying non-target plants. Low volume application: For application by hand held weed wiper or C.D.A. use at dilutions with water of 100 g/L.

Post-Planting: Boom and Aerial Application

FORESTS AND PLANTATION TREES INCLUDING <i>EUCALYPTUS</i> SPP., <i>CORYMBIA MACULATA</i> AND <i>PINUS RADIATA</i>					
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ha	CRITICAL COMMENTS	
Capeweed, Flatweed, Thistles (except Hardhead Thistle), Volunteer Legumes, Skeleton Weed	Actively growing rosettes, seedlings up to 15 cm diameter or height	All States	500 mL - 1 L	Cupping of the tip leaves and 'weepy leader' symptoms may occur on certain <i>Eucalyptus</i> spp. and <i>Corymbia maculata</i> and are generally transient and do not result in long-term injury. These symptoms may be more obvious at rates of 160 mL/ha or higher or where mixtures are used on blue gum, shining gum and spotted gum. Where 'weepy leader' effect is a concern, use a directed spray. Use the 80 mL rate until 3 months post-planting and the 160 mL rate for trees 3 months and older. Use the low rate only under ideal conditions with excellent weed growth and where knockdown control of small weeds is desired. Use the high rate where longer control is required of larger weeds. For the control of annual and certain perennial grasses VICTORY® can be tankmixed with Verdict ¹ 520 Herbicide. See also comments in Mixing section. Uptake Spraying Oil should not be used in tankmixes with Verdict 520 and VICTORY® on sensitive species such as blue gum, shining gum or spotted gum where rates of VICTORY® are more than 320 mL/ha. Use a 100 % non-ionic surfactant such as Wetspray 1000 at 0.1 % v/v instead.	
Capeweed, Flatweed, Fleabane, Thistles including Hardhead Thistle, Volunteer Legumes, Skeleton Weed	Actively growing rosettes and seedlings greater than 15 cm diameter or height up to stem elongation and before flowering		2 L		
Californian Thistle	From early bud to flowering (December to February)				For best control of California Thistle use a wetter such as Wetspray 1000 at 0.1 % v/v. A second annual application may also be required for best control.
Ragwort	Small rosettes to larger rosettes up to stem elongation and before flowering		1 L or 2 L		Spray from the rosette to the shooting stage of growth. For small rosette seedling plants use the lower rate. For large rosette multi-crown and/or perennial plants use the higher rate. Addition of a 100 % non-ionic surfactant such as Wetspray 1000 at 0.1 % v/v is recommended. Add diquat (200 g/L) at 1 L/100 L water plus a surfactant after opening of the first flowers, to prevent the formation of viable seed. Where diquat is added use a directed spray to avoid tree injury.
Sorrel (suppression only)	Actively growing rosettes, seedlings up to 15 cm diameter or height		6 L - 8.5 L		Higher rates give better suppression. At rates greater than 960 mL use a directed spray to avoid tree injury.

***PINUS RADIATA* AND *EUCALYPTUS* SPP. PLANTATION ONLY**

WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ha	CRITICAL COMMENTS
Silver Wattle	Active growth Spring to Summer (0.5 - 2 m tall)	NSW, ACT, Tas, Vic, SA only	5 L	For effective control apply when bushes are growing actively. Large trees will not show complete necrosis. For boom spraying apply in 150 to 200 water/ha. For aerial treatment apply in a minimum of 50 L/ha of water containing 25 to 50 % by volume of anti-evaporant such as Ulvapon ¹ . Mix VICTORY® and water first and then add Ulvapon. Maintain continuous agitation. At rates of 1120 mL and 1360 mL for <i>Eucalyptus</i> spp. use a directed spray to avoid tree injury.
	Active growth Spring to Summer (2 - 4 m tall)		7 L	
	Active growth Spring to Summer (4 - 8 m tall)		8.5 L	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

PASTURES and FALLOW LAND:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

CEREALS and CANOLA:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

CEREALS:

DO NOT APPLY LATER THAN 10 WEEKS BEFORE HARVEST.

CANOLA:

DO NOT HARVEST FOR 12 WEEKS AFTER APPLICATION.

FOREST, except *Pinus radiata* PLANTATIONS:

DO NOT GRAZE FOR 7 DAYS AFTER APPLICATION.

***Pinus radiata* PLANTATIONS:**

DO NOT GRAZE FOR 14 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

MIXING

Half fill the spray tank with water and add the required quantity of VICTORY® and complete filling. Agitate continuously to ensure thorough mixing before and during application. Only mix sufficient chemical for each day's work.

Tank mixtures: Wettable powder or dry flowable formulations should be added to the spray tank first, followed by suspension concentrates (flowables), aqueous concentrates (VICTORY®) and the emulsifiable concentrate formulations (e.g. Verdict[†] or LVE MCPA).

COMPATIBILITY

VICTORY® is compatible with the following:

Broadleaf Herbicides: Fluroxypyr, Lynx® WG, bromoxynil, chlorsulfuron, diuron, Wipe-Out®, MCPA amine, MCPA LVE, paraquat, diquat + paraquat, terbutryn, 2,4-D amine.

Grass Herbicides in Cereal Crops: Diclofop methyl for Wild Oats control only. Mandate® for Wild Oats and Phalaris control.

Grass Herbicides on Broadleaf Crops: Verdict[†] (haloxyfop).

APPLICATIONS

DO NOT apply this product by air or mister within a Chemical Control Area in Victoria unless a permit has been issued by The Department of Economic Development, Jobs, Transport and Resources (DEDJTR).

Boom Spraying Plantation Trees: Apply VICTORY® in sufficient water to obtain good coverage. It should be applied by an accurately calibrated ground rig or aircraft, delivering 200 - 300 micron droplets and not less than 50 L/ha water volume for boom sprayers or not less than 20 L/ha for aerial application.

Hardhead Thistle – Use a spray volume of 200 - 250 L/ha of water.

Silver Wattle – Use a spray volume of 150 - 200 L/ha of water by ground boom spray and a minimum spray volume of 50 L/ha by aircraft.

Boom Spraying Crop and Pastures: Apply VICTORY® in sufficient water to obtain good coverage. It should be applied by an accurately calibrated ground rig or aircraft, delivering 200 - 300 micron droplets and not less than 50 L/ha water volume for boom sprayers or not less than 20 L/ha for aerial applications.

Hardhead Thistle – Use a spray volume of 200 - 250 L/ha of water.

Silver Wattle – Use a spray volume of 150 - 200 L/ha of water by ground boom spray and a minimum spray volume of 50 L/ha by aircraft.

High Volume Hand Gun: Apply the recommended mix to give full coverage of leaves and stems through a No. 6 - 8 tip at 700 - 1500 kPa. Spray volume for effective coverage of dense two metre high Silver Wattle should be 30 to 40 litres of spray per 100 m² (10 m x 10 m) of infestation. For larger areas an equivalent would be 3000 to 4000 litres per infested hectare.

CLEANING SPRAY EQUIPMENT

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plant and water courses.

PARTIAL CLEANING

(Before spraying other labelled or tolerant crops): After using VICTORY®, empty the tank completely and drain the whole system.

Thoroughly wash inside the tank using a pressure hose. Quarter fill the tank with clean water and circulate through the pump, line, hoses and nozzles. Drain and repeat procedure twice.

COMPLETE CLEANING

(Before spraying susceptible crops): After using VICTORY®, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose. Quarter fill the tank with clean water and circulate as above, then drain. Quarter fill the tank again and add a liquid detergent (e.g. Surf[†], Omo[†], Drive[†]) at 500 mL/100 L water and circulate throughout the system for at least fifteen minutes. Drain, remove filters and nozzles and clean separately. Rinse inside the tank thoroughly using a pressure hose and flush system with clean water.

RESISTANT WEEDS WARNING

VICTORY® Herbicide is a member of the Pyridines group of herbicides. VICTORY® has the disruptors of plant cell growth mode of action. For weed resistance management VICTORY® is a Group I Herbicide. Some naturally occurring weed biotypes resistant to VICTORY® and other Group I herbicides may exist through normal genetic variability in any weed population.

GROUP I HERBICIDE

The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by VICTORY® or other Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Adama Australia accepts no liability for any losses that may result from the failure of VICTORY® to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

Composts and mulches: DO NOT apply VICTORY® to crops or pastures that will be used for the production of compost or mulches or mushroom substrate. Such compost or mulch made from plant material treated with VICTORY® may cause damage to susceptible crops and plants.

Susceptible Crops and plants include, but are not limited to chickpeas, clover, cotton, faba beans, field peas, fruit trees, lentils, lupins, lucerne, medics, ornamentals, potatoes, safflower, tomatoes, vegetables, grape and kiwifruit vines, vetches and wattle. Field peas, faba beans, lentils and vetches are particularly susceptible and should be sown the season following an application of VICTORY® at 500 mL/ha.

Where VICTORY® residue carry over from use rates of less than 500 mL/ha is suspected and susceptible crops are to be planted, test the treated area as follows:

- **Field bioassay** – where rain allows, plant a small area of the susceptible crop 4 to 6 weeks before desired planting date and take note of any symptoms of injury. If any herbicide symptoms are observed, only plant either canola or a cereal (see recommendation for northern and southern Australia below).
- **Pot bioassay** – where not practical to do field bioassay, plant a small number of seeds of the susceptible crop into pots containing soil from the treated field. Do this 4 to 6 weeks before desired planting date. If any herbicide symptoms are observed, only plant either canola or a cereal (see recommendation for northern and southern Australia below).

Stubble from treated crops: Ensure that harvesters effectively spread crop straw and do not leave a heavy 'header' trail after harvest. Burn (if legal in the area), bale and remove, slash or incorporate stubble as soon as practical after harvest and as long as possible before planting next year to allow microbial breakdown of any residues in straw. Heavy stubble loads may carry more residue into the following season. Where heavy stubble burdens and/or non-wetting soils exist and less than recommended rain amount have occurred from application to planting the susceptible crop (see below), only plant a Winter or Summer cereal or canola.

Planting crops following use of VICTORY® in previous crop: Planting crops 'dry' without significant rain (see below) in the 'Autumn break' increases the risk of injury to susceptible crops. This practice should be avoided, or only plant a Winter or irrigated Summer cereal crop or canola. In severely dry conditions, where less than 30 % of average rainfall and/or less than the minimum rain (see below) has fallen between application and planting the next year, only plant a Winter or irrigated Summer cereal or canola.

PLANT BACK PERIODS FOR SOUTHERN AUSTRALIAN WINTER DOMINANT RAINFALL AREAS (Sth NSW, VIC, SA, WA)

Required rainfall: A minimum 25 mm rain event in the post harvest Summer to Autumn period, with a subsequent extended period of at least 1 week where the top 10 cm of the soil stays moist is required to enable breakdown of soil residues. Fastest residue breakdown will occur under good soil moisture and warm conditions, which promote microbial activity. Where significant rain (> 25 mm) has fallen in Summer to Autumn, with soil wetting for at least one week, the following plantback periods apply:

Following Crops	Rate (mL/ha)	Plantback Interval
Clover, Chickpea, Faba Bean, Field Pea, Lentils, Lupins, Medics and Vetch	Up to 300	9 months
	300-500	12 months
	>500	24 months
Barley, Canola, Wheat, Oats	All label rates	1 week

PLANT BACK PERIODS FOR NORTHERN AUSTRALIAN SUMMER DOMINANT RAINFALL AREAS (Nth NSW, QLD)

Required rainfall before plantback:

If planting susceptible Summer crops – at least 100 mm rain.

If planting susceptible Winter crops – at least 150 mm rain.

This rain or irrigation should wet the soil for extended periods (at least one week) this is essential for breakdown of soil residues prior to planting susceptible crops.

If planting a cereal or canola crop – at least 50 mm rain or irrigation is required to enable soil wetting for at least one week. Where these requirements have been met the following plantback periods apply:

Following Crops	Rate (mL/ha)	Plantback Interval
	Up to 75 mL/ha	>75-300 mL/ha
Chickpea, Cotton, Soybean, Sunflower	3 months	6 months
Lucerne	9 months	9 months
Maize, Sorghum	1 week	2 weeks
Wheat, Barley, Oats, Canola	1 week	1 week

Note: Susceptible crops should not be sown for at least 2 years where VICTORY® at more than 300 mL/ha has been used in northern Australia.

PROTECTION OF LIVESTOCK

DO NOT graze or cut treated crops for stock food except as specified under WITHHOLDING PERIODS.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

VICTORY® has low toxicity to fish, birds, honey bees, livestock, earthworms and aquatic organisms. DO NOT contaminate streams, rivers or waterways with chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank.

DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or a designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

Envirodrum Micro Matic Valve (110 L): Store the original sealed Envirodrum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the Envirodrum to the point of purchase. The Envirodrum remains the property of Adama Australia.

1000 L: Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs Adama Australia should be advised immediately. This minibulk container is reusable and remains the property of Adama Australia. DO NOT rinse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return to Adama Australia for cleaning, relabelling and refilling.

SMALL SPILL MANAGEMENT

Wear protective equipment (See Safety Directions). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal. (See Storage and Disposal section). If necessary, wash the spill area with an alkali detergent and water and absorb as above, the wash liquid for disposal.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing the spray, wear elbow-length PVC gloves and a face shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for VICTORY® is available from adama.com or call Customer Service on 1800 423 262.

CONDITIONS OF SALE: The use of VICTORY® Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Adama Australia, regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Adama Australia accepts no responsibility for any consequence whatsoever resulting from the use of this product.

® Trademarks of an Adama Group Company

† Other trademarks

05/2017 9041

UN No. 3082
ENVIRONMENTALLY HAZARDOUS
SUBSTANCE, LIQUID, N.O.S.
PACKAGING GROUP: III
HAZCHEM CODE: 3Z

