

Product overview

Venturi® MAX is a concentrated SC formulation containing 480 g/L of methoxyfenozide. It is an effective lepidoptera insecticide registered for use in various fruit, nut and vine crops. It is ideal for use with IPM programs.

Key features

- Double strength formulation for ease of handling
- Broadest label claims for methoxyfenozide
- Highly effective against eggs and larvae
- Residual activity of up to 21 days
- IPM compatible with low impact on beneficial insects, including bees and common lepidopteran predators and parasitoids
- Alternative mode of action to reduce the risk of insecticide resistance
- Compatible with a wide range of products

Crop	Pest	Venturi® MAX rate and adjuvant recommendation
Almonds	Carob moth (1st and 2nd generation)	40 mL/100 L + wetter
Apples and pears, blueberry, grapevines, kiwifruit, persimmons	Light brown apple moth	
Apples and pears	Loopers	
Avocado, coffee	Avocado leafroller	12.5 mL/100 L
Custard apple, persimmons	Yellow peach moth	
Macadamia	Macadamia flower caterpillar	
Citrus	Light brown apple moth	12.5 mL/100 L + wetting agent at 10 mL/100 L
Macadamia, longan, lychee	Macadamia nut borer	20 mL/100 L
Tomatoes, peppers (capsicum and chilli), eggplant and okra	Native budworm, Tomato grub, Cluster caterpillar	63–85 mL/100 L or 625–850 mL/ha

Product information

Active ingredient:

480g/L methoxyfenozide

Mode of action:

Group 18 insect growth regulator

Formulation:

Suspension Concentrate (SC)

Registered crops:

Almond, apples and pears, avocado, blueberries, capsicum, chilli, citrus, coffee, custard apple, eggplant, grapevines, kiwifruit, longan, lychee, macadamia, okra, persimmon and tomato.

Application information

Adjuvant:

Where recommended apply with Agral® Spray Adjuvant, Wetter 1000 or equivalent

Incompatible with mineral oils

Rainfast:

Once spray is dry (observing the 6 hour run-off restraint)

Ground:

Concentrate or dilute spraying with sufficient water to ensure adequate coverage

Air:

DO NOT apply by air



Venturi® MAX

Withholding periods

Harvest:

Almond, blueberry and persimmons:

DO NOT harvest for 7 days after application

Avocado, coffee, custard apple, kiwifruit, lychee and pome fruit:

DO NOT harvest for 14 days after application

Citrus fruits:

DO NOT harvest for 1 day after application

Grapes (for domestic consumption):

DO NOT harvest for 21 days after application

Grapes (for export or wine production):

Consult your winery, industry spray diary[^] or peak industry body for the recommended withholding period (export harvest interval) to meet export residue requirements.

^AWRI Dog Book 2021/22 guideline for methoxyfenozide is "Use no later than 80% capfall". Refer to current/online version.

Longan:

DO NOT harvest for 3 days after application

Macadamia:

DO NOT harvest for 28 days after application

Tomato, peppers (capsicum and chilli), eggplant and okra:

Not required when used as directed

Grazing:

All Crops:

DO NOT graze treated crops

DO NOT allow livestock to graze within treated orchards, plantations or vineyards.

Compatibility

Venturi MAX is physically compatible with a range of fungicides and insecticides. The following products have been tested to be compatible at common label rates: Mirador® 625, Custodia® Forte, Cavalry® Weatherquard, Kocide* Blue, Mancozeb, Tri Base Blue*, Filan*, Fontelis*, Cabrio*, Trivor®, Sorcerer® 36, Transform*, Movento*, Paramite* and Success* Neo. As formulations of other manufacturers' products are beyond the control of ADAMA Australia, all mixtures should be tested prior to mixing commercial quantities. Venturi MAX is not compatible with mineral oils.







®Registered trademarks of an ADAMA Agricultural Solutions Ltd Company. Please note: This information is not intended to replace the product label. Always read the complete product label appearing on the container before opening or using products. ADA22131.



