

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Ardeo®

Herbicide

ACTIVE CONSTITUENT: **224 g/L ACIFLUORFEN**

present as the SODIUM SALT

GROUP

14

HERBICIDE

Uses: For the control of certain broadleaf weeds and grasses as per the Directions for Use

Crops: Green Beans, Mung Beans, Peanuts, Soybeans, and Seed Crops of Siratro and Stylo

Formulation type

Soluble
Concentrate

SL



adama.com

CONTENTS: 1 L to 1000 L

RESTRAINTS

DO NOT mix with oils, fertilisers or other pesticides, except where indicated.

DO NOT apply when crops or weeds are under stress conditions.

DO NOT apply as a post-emergence treatment to control hard to kill weeds when rainfall is expected within 6 hours.

DIRECTIONS FOR USE

Table 1. POST-EMERGENCE TREATMENTS

CROP	WEEDS CONTROLLED	WEED SIZE	RATE / ha	CRITICAL COMMENTS
Soybeans (Qld, NSW, WA only), Peanuts (Qld, WA only), Mung Beans (Qld only), Adzuki Beans (<i>Vigna angularis</i>) (All States)	Apple of Peru, Turnip Weed, Wild Radish	Cotyledon to 2 true leaves	1.0 L	Apply with thorough coverage to young actively growing weeds.
		2 to 6 true leaves	1.5 L	
	Prince of Wales Feather, Redshank, Thornapple	Cotyledon to 2 true leaves		
		2 to 6 true leaves	2.0 L	
	Pigweed	Cotyledon to 50 mm diameter	1.0 L	
		50 to 100 mm diameter	1.5 L	
	Annual Ground Cherry, Wild Gooseberry	Cotyledon to 4 true leaves	1.0 L	
		4 to 8 true leaves	1.5 L	
		8 leaves to 200 mm high	2.0 L	

Table 1. POST-EMERGENCE TREATMENTS - continued

CROP	WEEDS CONTROLLED	WEED SIZE	RATE / ha	CRITICAL COMMENTS
Soybeans (Qld, NSW, WA only), Peanuts (Qld, WA only), Mung Beans (Qld only), Adzuki Beans (<i>Vigna angularis</i>) (All States) - continued	Blackberry Nightshade, Bladder Ketmia, Capeweed, Fumitory, Noogoora Burr	Cotyledon to 4 true leaves	2.0 L	Apply with thorough coverage to young actively growing weeds.
	Bellvine	2 to 4 true leaves	1.5 L	
		4 to 6 true leaves	2.0 L	
	Giant Pigweed	Cotyledon to 2 true leaves	2.0 L	
	Jute	Cotyledon to 4 true leaves	1.5 L	
		4 to 6 true leaves	2.0 L	
	Sesbania	Cotyledon to 200 mm high		
Mung Beans (NSW only)	Bladder Ketmia, Noogoora Burr	Cotyledon to 4 true leaves		
Mung Beans (Qld, NSW only)	See weeds as above	Apply at the weed stage as indicated above	1.0 L plus Hasten* at 1 L/100 L water (1% v/v)	DO NOT exceed 1 L/ha ARDEO® with this mixture.
Green Beans (All States)	Prince of Wales Feather (<i>Amaranthus powellii</i>)	2 to 4 leaf stage	250 to 500 mL	DO NOT exceed 2 L of product per hectare per season. Apply up to 3 times by boom spray: • after the fully expanded unfoliate leaf stage of the crop. • when crop is at 6 to 7 leaf stage and/or when crop is at early flowering stage. See APPLICATION section below.
			250 to 750 mL	
Peanuts (Qld only)	Bellvine	2 to 6 true leaves	1.0 L ARDEO® PLUS 1.0 to 1.5 L of 400 g/L formulation of 2,4-DB/ha	Apply with thorough coverage as a tank mix to young actively growing weeds. Use the higher rate of 2,4-DB where larger weeds are present. The mixture will give additional control of larger weeds than given by ARDEO® alone.
	Noogoora Burr	Cotyledon to 6 true leaves		
Seed crops of Siratro and common and fine-stem Stylo (<i>Stylosanthes guianensis</i> only) (Qld only)	Billygoat Weed, Gambia Pea, Glycine, Greenleaved Desmodium, Five-leaved Cassia	Cotyledon to 4 true leaves	2.0 L	Apply with thorough coverage to young actively growing weeds. DO NOT use on other Stylos.

Treatment of weeds larger than those listed is not recommended because control may be incomplete due to increased weed tolerance and poor spray coverage. However, in a crop salvage situation, ARDEO® may be used at rates up to 4 L/ha, providing the application is not made within 50 days of harvest.

Table 2. PRE-EMERGENCE TREATMENTS: SOYBEANS AND PEANUTS ONLY. NOT RECOMMENDED FOR MUNG BEANS.

CROP	WEEDS CONTROLLED	RATE / ha	CRITICAL COMMENTS
Peanuts, Soybeans (Qld, WA only)	Apple of Peru, Black Bindweed, Blackberry Nightshade, Fat Hen, Fumitory, Jute, Narrow Leaf Plantain, Noogoora Burr, Pigweed, Potato Weed, Redshank, Scarlet Pimpernel, Turnip Weed, Wild Gooseberry, Wild Radish	3.0 L	Apply as a broadcast spray after sowing but before the crops and weeds emerge. Applications should be made to a well prepared, clod free soil of relatively fine tilth. Mechanical soil incorporation is not recommended as this reduces the activity of ARDEO®.
	Annual Ground Cherry, Barnyard Grass, Bladder Ketmia, Liverseed Grass, Mossman River Grass	4.0 L	

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

Harvest

GREEN BEANS, ADZUKI BEANS: DO NOT HARVEST FOR 28 DAYS AFTER APPLICATION.

CROP SALVAGE SITUATION WHEN ARDEO® IS USED AT UP TO 4 L/ha: DO NOT HARVEST FOR 50 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

MIXING

ARDEO® mixes readily with water. Fill the spray tank about two-thirds full with water and add the required amount of ARDEO®. Fill the spray tank and mix. Maintain sufficient agitation during mixing and spraying to ensure a uniform spray mixture.

Use of Surfactant: The use of low foam, non-ionic surfactant at the labelled rate should be used for maximum weed control. This surfactant should be added to the spray mixture after adding the required amount of ARDEO®.

Use of Crop Oil Adjuvants: ARDEO® may be applied to mung beans at the reduced rate of 1 L/ha in combination with Hasten* at 1 L/100 L (1 %v/v). The adjuvant should be added to the spray mixture after adding the required amount of ARDEO®. Use of this mixture may increase crop phytotoxicity, however the plants will soon recover and yield will be unaffected.

APPLICATION

Spray Equipment: ARDEO® can be applied by ground equipment using standard low pressure herbicide sprayers, equipped with hollow cone or flat fan nozzles. The spray equipment should be calibrated to deliver between 100 and 300 litres of spray mixture per hectare at a pressure of 250 to 300 kPa at the nozzle tips.

ARDEO® may be applied as a banded treatment directly over the crop row. When this method of application is used, it is important that the spray equipment be calibrated to give the correct dosage per sprayed hectare, and that the spray pattern gives maximum coverage of the weed foliage present.

Before using ARDEO®, it is important to check that the equipment is clean and that all materials from previous uses are completely flushed from the system.

Aerial Application: Queensland only. ARDEO® can be applied from a fixed-wing aircraft or helicopter, using a minimum of 48 litres of spray per hectare. Nozzles on spray booms should not be placed any closer to wing tips than three-quarters of the wing span; this will minimise the formation of spray or wing tip vortice roll. A spraying height of 2 to 3 metres over the crop is recommended. Swirl jet or hollow cone nozzles are suggested. Keep pressures below 275 kPa. Nozzles should point to the rear of the aircraft and not be pointed downward more than 20 degrees. DO NOT USE MICRONAIRE EQUIPMENT FOR ARDEO® APPLICATION.

GREEN BEAN CROPS

Early application is important. Treatment later than the recommended weed growth stage may result in inadequate control of Prince of Wales Feather (*Amaranthus powellii*).

The application of ARDEO® may cause some crop phytotoxicity: e.g. some tip burning and stunting.

The beans usually grow out of this, though they may remain about 10 cm shorter than unsprayed beans.

This does not appear to reduce yield. It is recommended that users consult an experienced field agronomist prior to using this product. Additionally it is recommended to apply in the evening if possible. Increased phytotoxicity may occur under warm humid conditions.

A non-ionic wetting agent may be added at 0.1% v/v. Thorough coverage is essential.

DO NOT apply earlier than fully-expanded unifoliate leaf stage.

Use higher rate after the beans have reached the first fully-expanded trifoliate leaf stage, to target *Amaranthus* which is greater than the 2-leaf stage. Multiple applications should only be used if necessary to treat the weed problem.

POST-EMERGENCE TREATMENTS

Best results are obtained when the crop and weeds are free from moisture and temperature stress, and when the relative humidity is around 65% or higher.

DO NOT apply ARDEO® if soil temperature is below 15°C or air temperature is below 20°C.

Crop damage will also be increased when ARDEO® is applied when air temperatures are above 30°C.

Since it is important that the maximum possible weed coverage is obtained, application should be made before the crop canopy shields the weeds from spray contact but after the maximum number of weed seedlings has emerged. This will usually occur between the 2nd and 5th trifoliate stages of the crop, or about one month after sowing.

Thorough coverage is essential for optimum weed control.

After treatment, soybean leaves may exhibit leaf cupping, crinkling and speckling, particularly on the youngest leaves present at the time of application. Soybeans recover from this injury and continue growing at a normal rate with no adverse effect on crop yield.

COMPATIBILITY

ARDEO® is compatible with 2,4-DB, Broadstar®, Firepower®, fluzifop-P, Platinum® and Sertin*.

RESISTANT WEEDS WARNING

ARDEO® Herbicide is a member of the Diphenyl ether group of herbicides.

GROUP	14	HERBICIDE
-------	-----------	-----------

The product has the inhibitor of protoporphyrinogen oxidase mode of action. For weed resistance management the product is a Group 14 herbicide. Some naturally occurring weed biotypes resistant to the product and other Group 14 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group 14 herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Adama Australia accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Wash sprayer thoroughly after use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT apply when weather conditions favour run off or drift from areas treated.

STORAGE AND DISPOSAL

1 to 100 L

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

This container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank.

DO NOT dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection or similar container management program site. The cap should not be replaced, but may be taken separately.

Returnable Containers (110 L, 1000 L)

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply or other designated collection point for refill or storage. This container remains the property of Adama Australia.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin.

If product on skin, immediately wash area with soap and water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

SDS

Additional information is listed in the safety data sheet (SDS). A SDS for ARDEO® Herbicide is available at adama.com or call Customer Service on 1800 423 262.

CONDITIONS OF SALE: The use of ARDEO® Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Adama Australia regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Adama Australia accepts no responsibility for any consequence whatsoever resulting from the use of this product.

® Registered trademark of an Adama Group Company.

* Other registered trademark.

NOT A DANGEROUS GOOD ACCORDING TO THE AUSTRALIAN DANGEROUS GOODS (ADG) CODE

11/2024 10086