

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# Folpan® 800 WG

## Fungicide

ACTIVE CONSTITUENT: **800 g/kg FOLPET**

GROUP

**M4**

FUNGICIDE

A preventative treatment for control of Downy mildew, Powdery mildew, Grey mould, Phomopsis cane and leaf spot, and Black rot in Grapes as shown under the directions for use.

Formulation type

Water Dispersible Granule

**WG**



# ADAMA

adama.com

NET CONTENTS: 1 kg - 1000 kg

### RESTRAINTS

**DO NOT** apply by aircraft.

**DO NOT** apply if heavy rains or storms that are likely to cause runoff from treated areas are forecast within 3 days.

**DO NOT** irrigate to the point of water runoff from the treatment area for at least 3 days after application.

**DO NOT** apply more than five applications per season in grapes.

### SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at [apvma.gov.au/spraydrift](http://apvma.gov.au/spraydrift)

**DO NOT** allow bystanders to come into contact with the spray cloud.

**DO NOT** apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The advisory buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

**DO NOT** apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

**DO NOT** apply if there are surface temperature inversion conditions present at the application site during the time of application. These conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

**DO NOT** apply by a vertical sprayer unless the following requirements are met:

- spray is not directed above the target canopy
- the outside of the sprayer is turned off when turning at the end of rows and when spraying the outer row on each side of the application site
- for dilute water rates up to the maximum listed for each type of canopy specified, minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for vertical sprayers') are observed.

| Buffer zones for vertical sprayers |                 |                       |                  |                  |                 |
|------------------------------------|-----------------|-----------------------|------------------|------------------|-----------------|
| Type of target canopy              | Bystander Areas | Natural Aquatic Areas | Pollinator Areas | Vegetation Areas | Livestock Areas |
| Grapevines                         | 0 metres        | 10 metres             | 0 metres         | 0 metres         | 0 metres        |

## DIRECTIONS FOR USE

Apply FOLPAN® 800 WG as a part of preventative fungicide spray program.

| CROP   | DISEASE   | RATE  | CRITICAL COMMENTS  |
|--------|---|---|--|
| Grapes | Downy mildew<br>( <i>Plasmopara viticola</i> )                | <b>Dilute spraying</b><br>125 g/100L<br><br><b>Concentrate spraying</b><br>Refer to the Mixing/<br>Applications section | FOLPAN® 800 WG is a multi-site fungicide with a preventative mode of action.<br><br>Apply preventatively from budburst as a part of downy mildew protectant spray program, with a 10-14 day spray interval. When the conditions favour disease development use the shorter spray interval.<br><br>Apply to point of run-off. |
|        | Powdery Mildew<br>( <i>Uncinular necator</i> )                |   | Incorporate FOLPAN® 800 WG as part of a season long spray program in support of sulfur or other single-site Mode-of-Action (MoA) treatments.<br><br>FOLPAN® 800 WG should be used as part of a powdery mildew preventative program   |
|        | Grey mould<br>( <i>Botrytis cinerea</i> )                     |   | Apply in a preventative spray program from early flowering until pre-harvest.<br><br>Apply to point of run-off.<br><br>FOLPAN® 800 WG can be used in conjunction with a systemic Botrytis product to assist with resistance management.  |
|        | Phomopsis cane and leaf spot<br>( <i>Phomopsis viticola</i> ) |   | Apply in a preventive spray program when shoots are 2-5 cm long.   |
|        | Black rot<br>( <i>Guignardia bidwellii</i> )                  |   | Apply just before bloom, just after bloom, and in first cover spray.<br>Maximum of 5 total applications per season with a minimal retreatment interval of 10-14 days.  |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS:**

**HARVEST**

**Grapes: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION**

**GRAZING**

**TREATED CROPS ARE NOT TO BE GRAZED OR FED TO ANIMALS.**

**DO NOT GRAZE VINEYARDS AFTER APPLICATION.**

### EXPORT TRADE ADVICE

TREATED CROP: Treated crop commodities destined for export may require extra time being allowed between application and harvest to be accepted in some export markets. Before you use this product you are advised to contact ADAMA Australia Pty Ltd and/or your industry body about any potential trade issues and their management.

### GENERAL INSTRUCTIONS

#### MIXING

Add the required amount of product to the 70% filled spray tank with the agitator in operation. Fill the tank with water and continue agitation during spraying and after a stoppage. Do not leave mixture in tank overnight. Rinse out spray tank, pumps and nozzles at the end of the day.

#### APPLICATION

##### Ground application

##### **Dilute Spraying - Grapes**

Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

##### **Concentrate Spraying:**

Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (see Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way.

##### **Example only**

1. Dilute spray volume as determined above: For example 1000L/ha
2. Your chosen concentrate spray volume: For example 500 L/ha
3. The concentration factor in this example is: 2X (i.e 1000L ÷ 500 L = 2)
4. If the dilute label rate is 125 g/100L, then the concentrate rate becomes 2 x 125 that is 250 g/100L of concentrate spray.

The chosen spray volume, amount of product per 100 L of water and the sprayer set up and operation may need to be changed as the crop grows. DO NOT use at a concentration rate higher than that specified in the Critical Comments as crop safety has not been assessed. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best practices.

#### COMPATIBILITY

If FOLPAN® 800 WG is to be mixed with other products, the physical compatibility of the mixture should be tested prior to use. For information on tank mix compatibility please contact Adama Australia.

#### FUNGICIDE RESISTANCE WARNING

For fungicide resistance management FOLPAN® 800 WG Fungicide is a Group M4 fungicide. Some naturally occurring fungi resistant to the product and other Group M4 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if FOLPAN® 800 WG or other Group M4 fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group M4 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

|       |           |           |
|-------|-----------|-----------|
| GROUP | <b>M4</b> | FUNGICIDE |
|-------|-----------|-----------|

Since occurrence of resistant fungi is difficult to detect prior to use, ADAMA Australia accepts no liability for any losses that may result from the failure of this product to control resistant fungi. FOLPAN® 800 WG may be subject to specific resistance management strategies. For further information contact your local supplier, ADAMA Australia or local agricultural department agronomist.

#### PRECAUTION

Do not apply using spraying equipment carried on the back of the user.

#### Re-entry Period

Do not allow entry into treated areas until the spray has dried. When prior entry is necessary wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), a washable hat and elbow-length chemical resistant gloves. Clothing must be laundered after each day's use.

#### PROTECTION OF LIVESTOCK

DO NOT graze or feed treated crops to animals.  
DO NOT graze vineyards after application.

#### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants, crops, cropping lands or pastures.

#### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

#### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Single-rinse or shake remainder into spray tank. DO NOT dispose of undiluted chemicals on site. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, dispose of empty container or unused product in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

#### SAFETY DIRECTIONS

Harmful if inhaled. Will damage eyes, will irritate skin and may irritate the nose and throat. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow length chemical resistant gloves, washable hat and face shield or goggles.

If applying by closed cab tractor wear cotton overalls buttoned to the neck and wrist (or equivalent clothing).

If applying by open cab tractor, wear cotton overalls buttoned to the neck and wrist and a washable hat, gauntlet length chemical resistant gloves and a disposable mist facemask covering mouth and nose.

Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

#### SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for FOLPAN® 800 WG Fungicide is available from adama.com or call Customer Service on 1800 423 262.

**CONDITIONS OF SALE:** The use of FOLPAN® 800 WG Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by ADAMA Australia regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and ADAMA Australia accepts no responsibility for any consequence whatsoever resulting from the use of this product.

® Registered trademark of an ADAMA Group Company

05/2026 25072