

STOP! READ THE PAMPHLET BEFORE USING THE PRODUCT
AND CONSULT A PROFESSIONAL IN AGRICULTURAL
SCIENCES.



Mirador 25 SC
FUNGICIDE – STROBILURIN
AZOXYSTROBIN

CAUTION

ANTIDOTE: DO NOT EXIST

**THIS PRODUCT MAY BE FATAL IF SWALLOWED
AND/OR IF INHALED MAY CAUSE DAMAGE TO THE
EYES AND SKIN BY EXPOSURE**

**DO NOT STORE IN DWELLINGS
KEEP OUT OF THE REACH OF CHILDREN,
MENTALLY UNSOUND PERSONS, DOMESTIC
ANIMALS, FOOD AND MEDICATIONS**



AGRONOMIC USE

MODE OF ACTION: MIRADOR 25 SC is a fungicide with protectant, curative, eradicant, translaminar and systemic properties. Inhibits spore germination and mycelial growth, and also shows antisporulant activity.

APPLICATION EQUIPMENT: MIRADOR 25 SC can be used with any conventional spray equipment. Before using make sure all application equipment and nozzles are properly calibrated and are in good working condition and without leaks. In terrestrial applications use a mixture volume of 200 L/ha and on aerial sprays use a volume of 50-80 L/ha. Use hollow cone spray nozzles.



HOW TO PREPARE THE MIXTURE: Fill the tank with water until half full then pour in the required dosage of **MIRADOR 25 SC**, complete the filling of the tank with water and stir the solution. The mixture should be used on the same day of preparation. Persons mixing and washing the equipment should wear the following protective gears; gloves, mask or respirators, goggles and rubber boots.



RECOMMENDATIONS FOR USE:

CROP	CONTROLLED DISEASES	RECOMMENDED DOSE
Tomato <i>Lycopersicum esculentum</i>	Early Blight <i>Alternaria</i> sp.	0.30-0.75 l/ha
Rice <i>Oryza sativa</i>	<i>Rhizoctonia solani</i>	0.60-0.70 l/ha
Banana <i>Musa</i> spp	Black sigatoka <i>Mycosphaerella fijiensis</i> var <i>difformis</i> Postharvest: Crown rot <i>Verticillium</i> sp., <i>Cephalosporium</i> sp.,	0.40 l/ha 300 a 400 ppm

	<i>Botryodiplodia</i> sp., <i>Fusarium</i> sp.	
Melon <i>Cucumis melo</i>	Downy mildew <i>Pseudoperonospora cubensis</i>	0.40 l/ha

APPLICATION INTERVAL: Spaced 8-day applications.

PREHARVEST INTERVAL: Rice and banana 0 days. Tomato 7 days. Melon: 7-10 days.

INTERVAL FOR REENTRY INTO TREATED AREAS: 24 hours or when the product has totally dried out.

PHYTOTOXICITY: Not phytotoxic at recommended doses.

COMPATIBILITY: Compatible with most products.

PRECAUTIONS AND WARNINGS FOR USE:

STORAGE AND TRANSPORT: Do not store or transport along with foodstuff, cooking utensils, and other household items. Store away from animal feed and medications, in dry and well ventilated areas away from heat sources. Keep this product in its original container hermetically sealed and duly labeled. Don't store along with fertilizers, insecticides or fungicides.

**DO NOT STORE IN DWELLINGS
KEEP OUT OF THE REACH OF CHILDREN.**



**DO NOT SMOKE, EAT OR DRINK WHILE
APPLYING OR HANDLING THIS PRODUCT.
SHOWER AFTER WORKING WITH THIS PRODUCT
AND WEAR CLEAN CLOTHES.**



SIGNS OF INTOXICATION: Irritation of the skin, eyes.

FIRST AID:

INHALATION: Take victim to fresh air. In case of breathing problems, give artificial breathing. Take patient to the doctor.

INGESTION: Wash mouth with plenty of clean water. Do not give anything by mouth to an unconscious person.

SKIN CONTACT: Take off contaminated clothes. Wash skin with plenty of water and soap.

EYES CONTACT: Wash with plenty of clean water for 15 minutes keeping eyelids open. Take patient to the doctor.

DO NOT GIVE TO DRINK NOR INDUCE VOMIT IN AN UNCONSCIOUS PERSON.

ANTIDOTE AND MEDICAL TREATMENT: There is no specific antidote. Treat symptomatically and give supportive therapy.

NATIONAL INTOXICATION CENTERS

INSTITUTION

COUNTRY	TELEPHONE
Centro de Información y Asistencia Toxicológica	Guatemala 251-3560
Centro de Intoxicaciones Karl Heusner Memorial Hospital	Belize 231-1548
Hospital San Rafael, Santa Tecla	El Salvador 2288-0417
Secretaría de Agricultura y Ganadería	Honduras 232-6213
Centro Nacional de Toxicología	Nicaragua 289-4514
Centro Nacional de Intoxicaciones	Costa Rica 223-1028
Centro de Investigaciones e Información de Medicamentos y Tóxicos	Panama 269-2741

ENVIRONMENTAL PROTECTION MEASURES**TOXIC TO FISH AND/OR CRUSTACEANS.**

**DO NOT CONTAMINATE RIVERS,
LAKES OR PONDS WITH
THIS PRODUCT OR ITS EMPTY CONTAINERS OR
PACKAGES.**

**MANAGEMENT OF CONTAINERS, PACKAGES, WASTE
AND REMNANTS**

Use the entire contents of the bottle after finishing the product, wash and rinse three times with plenty of water and dispose the resulting solution to the already prepared application mix. Perforate empty bottles. Take empty containers to the nearest recollection center or dispose it according to the instructions of the distributor of the product. Remnants of pesticides should be kept in its properly closed, labeled and original container.

**USING CONTAINERS AND PACKAGES FOR PURPOSES
OTHER THAN THOSE FOR WHICH THEY WERE
DESIGNED, MAY BE HAZARDOUS TO HUMAN HEALTH
AND THE ENVIRONMENT.**

DECLARATION OF WARRANTY: The manufacturer guarantees that this product is in concordance with the chemical description on the label and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal conditions of use. The manufacturer is not responsible for the improper use of this product.

FORMULATOR:

**ADAMA ANDINA B. V. SUCURSAL COLOMBIA
CALLE 1C NÚM. 7-53 INTERIOR ZONA FRANCA
BARRANQUILLA, COLOMBIA
TEL.: (571) 5-3799771, FAX: (571) 5-3799715**

IMPORTED BY:

COUNTRY	REGISTRATION NUMBER	REGISTRATION DATE
GUATEMALA	898-67	17-08-12
BELICE	0268-3	29-05-09
HONDURAS	433, F261, TII	07-02-12
NICARAGUA	MC-274E-2-2009	20-11-09
PANAMÁ	5168	17-12-09
REP. DOMINICANA	3187	16-03-09