

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Bobcat® i-MAXX

Herbicide

ACTIVE CONSTITUENTS:

125 g/L HEXAZINONE
25 g/L IMAZAPIC

present as the AMMONIUM SALT

SOLVENTS: 25 g/L N-METHYL PYRROLIDONE
250 g/L HYDROCARBON LIQUID

GROUP	B C	HERBICIDE
-------	-------	-----------

For the control of a wide range of annual and perennial grasses and broadleaf weeds in Sugarcane as specified in the Directions for Use table

Formulation type

Soluble
Concentrate

SL



ADAMA

adama.com

CONTENTS: 5 L, 10 L, 20 L, 110 L, 200 L, 1000 L

DIRECTIONS FOR USE

RESTRAINTS: DO NOT apply to fields where the slope exceeds 5%.
DO NOT apply BOBCAT® i-MAXX and any other product containing imazapic or hexazinone more than once per calendar year.
DO NOT apply in the planting furrow in plant cane.
DO NOT apply to soils saturated with water.
DO NOT apply by aircraft.

Table 1. Pre-emergence application in ratoons

SITUATION	WEEDS CONTROLLED	RATE PER ha	CRITICAL COMMENTS
Sugarcane ratoon crops (Qld, NSW and WA only)	Awnless Barnyard Grass (<i>Echinochloa colona</i>), Barnyard Grass (<i>E. crus-galli</i>), Blackberry Nightshade (<i>Solanum nigrum</i>), Blue Billygoat Weed (<i>Ageratum houstonianum</i>), Calopo Vine (<i>Calopogonium mucunoides</i>) Common Sida (<i>Sida rhombifolia</i>), Giant/Black Pigweed (<i>Trianthema portulacastrum</i>), Glossy Nightshade (<i>Solanum americanum</i>), Green Amaranth (<i>Amaranthus viridis</i>), Green Summer Grass (<i>Brachiaria subquadripara</i>), Ipomoea spp., Guinea Grass (<i>Panicum maximum</i>), Liverseed Grass (<i>Urochloa panicoides</i>), Milkweed (Mexican Fire Plant) (<i>Euphorbia heterophylla</i>), Pigweed (<i>Portulaca oleracea</i>), Sowthistle (<i>Sonchus oleraceus</i>), Summer Grass (<i>Digitaria ciliaris</i>)	2.9 - 3.8 L	Apply as pre-emergent application after harvest and before new cane growth appears. If applying as a band treatment, avoid throwing excessive untreated soil onto the treated band when inter-row cultivating. Use the higher rate of BOBCAT® i-MAXX for increased residual control of weeds or anticipated high weed pressure. Heavy trash layers may prevent the product from reaching the soil. Apply to weed-free soil prior to weed germination. Sufficient rainfall, or overhead irrigation, is required after application and prior to weed emergence to wet soil to a minimum depth of 5 cm. Emerged weeds must be controlled by prior cultivation, or application of an appropriate herbicide. Weed control can be improved under dry conditions or where incorporating moisture has been insufficient, by cultivating the treated soil above setts. Care needs to be taken to minimise exposure of untreated soil or setts. Prolonged wet soil and/or cool conditions can increase crop damage, especially in sandy soil. Normal growth will return when conditions become warm and drier. Pre-emergence weed control may be limited on Krasnozem or red brown Ferrosol soil types, where moisture is not maintained in the top 5 cm of soil. Refer to Soil Type in the General Instructions. Weed escapes may require follow up application of an appropriate herbicide. Contact Adama for advice on variety susceptibility before using this product (refer to the CROP SAFETY section of the label). DO NOT apply more than 3.8 L/ha per year. DO NOT apply to blocks that are to be replanted soon after harvest.

Table 2. Application in emerged cane as a directed spray

SITUATION	WEEDS CONTROLLED	RATE PER HA	CRITICAL COMMENTS
Sugarcane plant and ratoon crops (Qld, NSW and WA only)	As per weeds listed in Table 1 plus control of those weeds expected with these rates of 250 g/L paraquat products alone	2.9 - 3.8 L plus 1.6 L of Spraytop® 250 SL Herbicide (or other registered 250 g/L paraquat product)	<p>Directed inter-row spray application in emerged cane: When applying BOBCAT® i-MAXX in emerged cane, always apply in a tank mix with paraquat. DO NOT apply in the planting furrow in plant cane. Only apply to plant cane as a directed inter-row spray after the final cultivation and hill formation.</p> <p>Apply as a directed inter-row spray, full width between rows or as a band treatment after final cultivation and hill formation to a weed-free surface. Use the higher rates of BOBCAT® i-MAXX for increased residual control of weeds or anticipated high weed pressure. Use as a directed spray to minimise contact with sugarcane foliage. Contact with sugarcane foliage may cause temporary crop damage observed as interveinal yellowing and reduction in crop biomass or crop height. Refer to Spraytop® 250 SL Herbicide label for further instructions on spray boom types and safety precautions. Reduce the incidence of “fines” by using the correct nozzles and operating pressures. Replace worn or old nozzles and calibrate equipment on a regular basis. DO NOT cultivate within 1 hour of application. DO NOT spray plants covered with heavy dew. Prolonged wet soil and/or cool conditions can increase crop damage, especially in sandy soil. Normal growth will return when conditions become warm and drier. Pre-emergence weed control may be limited on Krasnozem or red brown Ferrosol soil types, where moisture is not maintained in the top 5 cm of soil. Refer to Soil Type in the General Instructions. Contact Adama for advice on variety susceptibility before using this product (refer to the CROP SAFETY section of the label). DO NOT use any more than 3.8 L/ha per year. DO NOT apply to blocks that are to be replanted soon after harvest.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED

GRAZING: DO NOT GRAZE CROP OR SPRAYED VEGETATION, OR CUT FOR STOCKFOOD FOR 6 WEEKS AFTER APPLICATION

GENERAL INSTRUCTIONS

All Applications: Reduce the incidence of “fines” by using the correct nozzles and operating pressures. Replace worn or old nozzles and calibrate equipment on a regular basis.

BOBCAT® i-MAXX is a soil residual, broad spectrum, general herbicide for control of many annual and perennial grasses and broadleaf weeds. Best results occur if BOBCAT® i-MAXX is applied shortly before weed germination and growth occurs.

Well established weeds should first be eliminated by mechanical or other means. For best results soil should be well prepared as free as possible from trash and clods. Results vary with soil type (lower rates are effective on lighter soils, higher rates may be necessary for effective control on heavier soils) and environmental conditions. Effects are slow to appear and may not become apparent until sufficient moisture (12 - 25 mm on moist soils, 25 - 50 mm on dry soils) in the form of rainfall or irrigation occurs after treatment to carry the chemical into the root zone of the germinating weeds. Best results are obtained if rainfall or irrigation occurs within 3 - 4 days after application.

Pre-emergence: Where possible, apply to soil two days prior to incorporation. Incorporate with adequate irrigation or rainfall to a minimum depth of 5 cm. When applying BOBCAT® i-MAXX in moist soil conditions, allow weeds to germinate prior to application and apply mixed with a recommended knockdown herbicide. Whenever emerged weeds are present BOBCAT® i-MAXX should be mixed with an appropriate knockdown herbicide.

Post-emergence: BOBCAT® i-MAXX must always be mixed with registered rates of paraquat (i.e. Spraytop® 250 SL herbicide) and applied as a carefully directed inter-row spray.

INCORPORATION

Incorporate via rainfall or irrigation within 3 - 4 days after application. For incorporation by irrigation, the entire treated area must be thoroughly wetted; however, care should be taken not to over-water and cause runoff.

APPLICATION

DO NOT apply by aircraft.

All dosages of BOBCAT® i-MAXX are expressed as overall rates. For band treatment use proportionately less, i.e. use only half of the overall rate when treating one half of the area. Temporary chlorosis of lower leaves may occur. To minimise this injury use low directed spray application. Tolerance of cane variety should be determined before treating large areas.

BOBCAT® i-MAXX is a water miscible liquid formulation. The product is mixed with water and applied at recommended rates with ground equipment. Apply in a spray volume of 400-600 L/ha with a properly calibrated boom.

When mixing BOBCAT® i-MAXX, partially fill the spray tank with water and add the correct amount of the product for the area to be sprayed with the agitation system engaged. Top up to correct volume with water, and for best results keep the spray solution agitated whilst spraying. Spray booms must be shut off while starting, turning, slowing or stopping, as overdose may result in injury to the crop. Use sufficient water (400 - 600 L/ha of area actually sprayed) to provide thorough and uniform coverage. Use sufficient pressure to allow to penetrate, 250 - 350 kPa is the pressure range generally used. Higher pressures should be avoided to prevent spray drift on to non-target areas.

FOLLOW CROPS

Under conditions which do not favour breakdown of this product, carry-over soil residues can affect susceptible follow crops. As environmental and agronomic factors make it impossible to eliminate all risks associated with this product, rotational crop injury is always possible. The following minimum re-cropping intervals (months after application) should be observed.

Re-cropping intervals following application:

MONTHS AFTER APPLICATION	
24	36
Cotton [†] , Sunflower, Mungbeans, Peanuts, Soybeans, Maize [#] , Sorghum [#] , Chickpeas, Lucerne [*] , Barley [*] , Wheat [*] , Lupins, Pasture Legumes	Any other crops

* The following additional requirements apply if it is intended to plant wheat (except for varieties with CLEARFIELD[^] Technology), barley or lucerne during the next Winter season.

DO NOT apply BOBCAT[®] i-MAXX in areas where rainfall from spraying to sowing of cereals or lucerne is expected to be below 200 mm.

[†] **DO NOT** apply BOBCAT[®] i-MAXX in areas where rainfall from spraying to sowing of cotton is expected to be below 550 mm.

[#] **DO NOT** apply BOBCAT[®] i-MAXX in areas where rainfall from spraying to sowing of sorghum or maize is expected to be below 500 mm.

CROP SAFETY

This product may, in some circumstances, lead to transient crop yellowing and temporary slowing of growth in sugarcane but plants soon recover and yield is unaffected. This effect may be more pronounced when the product is used under conditions where soil moisture is high within 2 days following application and on light/sandy soil. Contact Adama for advice on variety susceptibility before using this product.

RESISTANT WEEDS WARNING

BOBCAT[®] i-MAXX Herbicide contains members of the imidazolinone and triazinone groups of herbicides. BOBCAT[®] i-MAXX has the inhibitors-of-acetolactate-synthase (ALS) and inhibitor of photosynthesis at photosystem II modes of action. For weed resistance management BOBCAT[®] i-MAXX contains Group B and C herbicides. Some naturally occurring weed biotypes resistant to BOBCAT[®] i-MAXX and other Group B and C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by BOBCAT[®] i-MAXX or other Group B or C herbicides. Since the occurrence of resistant individuals is difficult to detect prior to use, Adama Australia accepts no liability for any losses that may result from the failure of BOBCAT[®] i-MAXX to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultants, local Department of Agriculture of Primary Industries, or Adama representative.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS THIS PRODUCT MAY KILL MANY TREES AND OTHER DESIRABLE PLANTS IF NOT CORRECTLY APPLIED.**

DO NOT apply or drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. On storage at elevated temperature, the product may produce carbon dioxide which may lead to pressure build-up in the container. A cap with pressure relief may be used for HDPE containers.

5 L, 10 L, 20 L containers only: Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or a designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

Returnable containers: If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured. Empty container by pumping through dry-break connection system. DO NOT attempt to breach the valve system or the filling point, or contaminate the container with water or other products. Ensure that the coupler, pump, meter and hoses are disconnected, triple rinsed with clean water and drained after each use. When empty, or contents no longer required, return the container to the point of purchase. This container remains the property of Adama Australia.

1000 L: Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs Adama Australia should be advised immediately. This minibulk container is reusable and remains the property of Adama Australia. DO NOT rinse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return to Adama Australia for cleaning, relabelling and refilling.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. When opening the container and preparing product for use, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical-resistant gloves and face shield or goggles. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If sprayed on skin, wash thoroughly. If sprayed in mouth, rinse mouth with water. If swallowed, DO NOT induce vomiting. Give a glass of water. If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for BOBCAT[®] i-MAXX Herbicide is available from adama.com or call Customer Service on 1800 423 262.

CONDITIONS OF SALE: The use of BOBCAT[®] i-MAXX Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Adama Australia regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Adama Australia accepts no responsibility for any consequence whatsoever resulting from the use of this product.

[®] Registered trademark of an Adama Group Company

[^] Other trademarks

NOT A DANGEROUS GOOD ACCORDING TO THE AUSTRALIAN DANGEROUS GOODS (ADG) CODE.

08/2017 9261