



ADAMA

# BADGE®

Get proven control of almost 30 broadleaf weeds in cereals, corn and flax with easy-to-use tank-mix options for one-shot weed control.



## Active Ingredients:

Bromoxynil and MCPA

## Herbicide Groups:

Group 4 (MCPA)

Group 6 (Bromoxynil)

## Formulation:

Bromoxynil (as octanoic ester) 225 g/L and MCPA (as 2-EH Ester) 225 g/L = EC

## Application Rates and Packaging:

- 0.5 L/ac
- 20 ac/jug
- 2 x 10 L jug/case

## REGISTERED CROPS:

### Field crops:

- Barley
- Canary seed
- Corn
- Fall rye
- Flax
- Oats
- Wheat (spring, winter)

### Seedling grasses:

- Bromegrass
- Creeping red fescue
- Crested wheatgrass
- Intermediate wheatgrass
- Meadow fescue
- Meadow foxtail
- Reed canary grass
- Russian wild rye
- Seedling meadow bromegrass
- Seedling tall fescue
- Slender wheatgrass
- Tall wheatgrass
- Timothy

## WEEDS CONTROLLED:

- American nightshade
- Ball mustard
- Bluebur
- Canada thistle<sup>1</sup>
- Cocklebur
- Common buckwheat
- Common groundsel
- Common ragweed
- Cow cockle<sup>2</sup>
- Flixweed
- Green smartweed
- Kochia<sup>3</sup>
- Lady's thumb
- Lamb's quarters
- Night-flowering catchfly
- Pale smartweed
- Perennial sow thistle<sup>1</sup>
- Redroot pigweed
- Russian thistle<sup>3</sup>
- Scentless chamomile<sup>4</sup>
- Shepherd's purse
- Stinkweed
- Tartary buckwheat
- Velvetleaf<sup>5</sup>
- Volunteer canola (all types)
- Volunteer sunflower
- Wild buckwheat
- Wild mustard
- Wild tomato
- Wormseed mustard

<sup>1</sup>Top growth control

<sup>2</sup>Up to 4-leaf stage

<sup>3</sup>When sprayed before plants are 2 inches high

<sup>4</sup>Spring annual only

<sup>5</sup>When sprayed before plants are 3 inches high



# BADGE®

## HOW IT WORKS:

A combination of systemic and contact activity with weeds yellowing within 2–4 days and exhibiting abnormal growth (twisting and cupping of leaves) in 2–10 days.

## CROP STAGING:

Crop	Stage
Barley, Oats, Wheat (spring)	2-leaf to early flag
Canary seed	3- to 5-leaf
Corn	4- to 6-leaf
Fall rye	When growth commences in spring to early flag leaf
Flax (including low-linolenic varieties)	2 inches to early bud stage. Best tolerance occurs when flax is 2–4 inches tall.
Seedling grasses	2- to 4-leaf
Winter wheat	2- to 4-leaf stage in the fall or after growth resumes up to early flag leaf

## WATER VOLUME:

Ground:

- 20–40 L/ac in cereals and flax
- 80–120 L/ac in corn
- 60 L/ac in forages

Aerial (wheat, barley and oats only): 8–20 L/ac

## REGISTERED AND SUPPORTED TANK MIXES:

- Corn: atrazine
- Flax: ARROW® 240 EC, ARROW ALL IN™ or Poast®
- Oats: MCPA ester
- Spring wheat and barley: Ally®, MCPA ester, Refine Extra®, BISON® 400 L or Traxos® (spring wheat only)
- Winter wheat: Refine Extra® or Traxos®

## MIXING INSTRUCTIONS:

1. Fill clean spray tank ½ full with water.
2. Add the required amount of tank-mix partner and agitate thoroughly.
3. Add the required amount of BADGE® and agitate thoroughly.
4. Fill the tank and agitate again before use.
5. When tank mixing, follow instructions on both labels.

## RAINFASTNESS:

1 hour

## CROP ROTATIONS:

No re-cropping restrictions the year after treatment.

## PRE-HARVEST INTERVAL:

Flax: 60 days

## STORAGE:

Avoid freezing.

## GRAZING RESTRICTIONS:

- Do not graze treated grain or established timothy crops or cut for feed within 30 days of application.
- Do not graze meadow foxtail in the year of treatment.
- Do not graze other treated forage grasses within 56 days of treatment.

## QUICK TIPS:

Avoid spraying if temperatures are above 25°C.



ADAMA

Always read and follow label directions.  
Toll-free: 1.855.264.6262  
Website: ADAMA.COM/CANADA

®/™ ARROW, BADGE® and BISON® are registered trademarks and ARROW ALL IN™ is a trademark of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies. © 2021 ADAMA Agricultural Solutions Canada Ltd.