

# FIFRA 24(c) Special Local Need Label (SLN)

For distribution and use only in the state of Montana

For use on Alfalfa Grown for Seed for control of Plant bugs (*Lygus* sp.),  
Alfalfa Weevil and Spotted Alfalfa Aphid

## CORMORAN™

**For Agricultural or Commercial Use Only**

**EPA Reg. No. 66222-264**

**SLN No. MT-180001**

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2021

### Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This state-specific Section 24(c) labeling and federal label for this product must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, and precautions on the EPA registered label for EPA Reg. No. 66222-264.

CROP	Insect	RATE OF APPLICATION
Alfalfa Grown for Seed	Plant Bugs ( <i>Lygus</i> sp.) Aphids (including Blue Alfalfa Aphid, Cowpea Aphid, Pea Aphid, Spotted Alfalfa Aphid) Alfalfa Weevil (suppression)	9.0 – 12.0 fl oz per acre

#### APPLICATION RATE:

Make application at the rate of 9.0 – 12.0 fl oz per acre. Make applications only in accordance with directions and conditions of use described in the EPA registered Cormoran™ label.

#### METHOD OF APPLICATION:

Cormoran™ may be applied in a minimum of 15 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.

#### TIMING/FREQUENCY OF APPLICATION:

Repeat applications if needed to maintain control, but do not make applications less than 7 days apart.

#### SPECIFIC USE RESTRICTIONS:

- Restricted Entry Interval is 12 hours.
- Preharvest interval of 14 days
- Do not apply more than 0.23 lb Novaluron active ingredient containing products per acre per season.
- Do not apply more than 0.3 lb Acetamiprid active ingredient containing products per acre per season.
- Do not apply more than 36.0 fl oz of formulated product per acre per season.

**OBSERVE THE FOLLOWING USE RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.**

**USE RESRICTIONS**

For ground application (all crops): Do not apply by ground equipment within 75 feet of bodies of water such as lakes, reservoirs, rivers, permanent streams, natural ponds, marshes or estuaries. All applications must include a 25-foot vegetative buffer strip within the buffer zone to decrease runoff.

For aerial application (except cotton): Do not apply by air equipment within 150 feet of bodies of water such as lakes, reservoirs, rivers, permanent streams, natural ponds, marshes or estuaries. All applications must include a 25-foot vegetative buffer strip within the buffer zone to decrease runoff.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to birds and aquatic invertebrates. This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treatment area. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Do not contaminate water when disposing of equipment washwater or rinsate. Do not contaminate water used for irrigation or domestic purposes. This product may contaminate water through drift of spray in wind. This product has a potential for runoff for several days to weeks after application. Poorly draining soil with shallow water tables is more prone to produce runoff. A level, well maintained vegetative (grass) buffer strip between areas to which this product is applied and the surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination. In order minimize the possibility of developmental effects on pollinator larvae, including honey bee brood, do not use CORMORAN™ on blooming crops.

24(c) Registrant:  
Makhteshim Agan of North America, Inc. d/b/a ADAMA  
3120 Highwoods Blvd., Suite 100  
Raleigh, NC 27604

Cormoran is a trademark of ADAMA



