

NIMITZ®

Nematicide

EPA REG. NO. 66222-243

SUPPLEMENTAL LABELING

This label expires on June 6, 2021 and must not be distributed or used after that date.

READ THE ENTIRE LABEL FOR NIMITZ® BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

“Label” as used in this supplemental labeling refers to the label booklet for NIMITZ® and this supplement.

ACTIVE INGREDIENT:	% BY WT.
Fluensulfone: 5-chloro-2 (3,4,4-trifluoro-but-3-ene-1-sulfonyl)-thiazole	40.0%
OTHER INGREDIENTS*:	60.0%
TOTAL:	100.0%

Contains 4 pounds fluensulfone per gallon in an Emulsifiable Concentrate formulation.

*Contains petroleum distillates

BRASSICA (COLE) LEAFY VEGETABLES (Crop Group 5), LEAFY VEGETABLES (Crop Group 4), Tobacco

KEEP OUT OF REACH OF CHILDREN

CAUTION/PRECAUCION

*Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)*

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This supplemental labeling must be in the possession of the user at the time of pesticide application.
- Read the “Limitation of Warranty and Liability” in the label booklet for NIMITZ® before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.
- These directions can be found on the currently registered EPA Stamped Label.

BRASSICA (COLE) LEAFY VEGETABLES (Crop Group 5) Broccoli, Cabbage, Cauliflower and other brassicas including: broccoli, Chinese (gai lon); broccoli raab (rapini); Brussels sprouts; cabbage, Chinese (bok choy); cabbage, Chinese (napa); cabbage, Chinese mustard (gai choy); cavalo broccolo; collards; kale; kohlrabi; mizuna; mustard greens; mustard spinach; rape greens, Japanese mustard spinach and hot herb mustard	3.5 to 7 pints/ treated acre (56 to 112 fl. oz./treated acre)	Transplanting: Apply at a minimum of 30 days prior to transplanting. Do not apply more than one application per crop and no more than 3.5 lbs. fluensulfone per acre per calendar year.
--	--	---

<p>LEAFY VEGETABLES (Crop Group 4)</p> <p>Celery, Lettuce (head and leaf), Spinach and other leafy vegetables including:</p> <p>Amaranth (Chinese spinach); arugula (roquette); cardoon; celery, Chinese; celtuce; chervil; chrysanthemum, edible-leaved; chrysanthemum, garland; corn salad; cress, garden; cress, upland; dandelion; dock (sorrel); endive (escarole); fennel, Florence; orach; parsley; purslane, garden; purslane, winter; radicchio (red chicory); rhubarb; spinach, New Zealand; spinach, vine; Swiss chard</p>	<p>3.5 to 7 pints/ treated acre (56 to 112 fl. oz./treated acre)</p>	<p>Transplanting: Apply at a minimum of 7 days prior to transplanting.</p> <p>Do not apply more than one application per crop and no more than 3.5 lbs. fluensulfone per acre per calendar year.</p>
<p>TOBACCO</p>	<p>3.5 to 7 pints/ treated acre (56 to 112 fl. oz./treated acre)</p>	<p>Apply at a minimum of 7 days before transplanting.</p> <p>Do not apply more than one application per crop and no more than 3.5 lbs. fluensulfone per acre per calendar year.</p>

NIMITZ is a trademark of an ADAMA Group Company.

EPA 06-09-2016