



ADAMA

Elegant®

Stronger, more flexible
broad- leaved weeds control








ELEGANT is a unique new post-emergent herbicide providing robust control of a wide range of broadleaf weeds in wheat and barley.

Simply. Grow. Together.

ADAMA.COM

Benefits to the Farmer

-  Very broad spectrum
-  Flexibility - Makes it easier to work around weather events and apply early or late
-  Increased efficacy on difficult to control broadleaf with a wide spectrum of activity weeds
-  Excellent value for money when compared to other premium broadleaf options
-  Stops weeds growing almost immediately improving yields from reduced competition

ELEGANT 2FD 306 SE controls a wide range of sensitive broadleaved weeds, among them:



BlackJack
Bidens pilosa



Datura
stramonium



Macdonalds eye
Galinsoga paviflora




Mallow





Mexican marigold
Tagetes minuta



Sow Thistle
Sonchus oleraceus



Wandering Jew
Commelina bengalensis



Mode of Action of Elegant

ELEGANT contains 2,4-D that produces uncontrollable, unsustainable weed growth, as well as florasulam that inhibits synthesis of amino acids. ELEGANT has a dual mode of action, therefore it is an excellent resistance management tool.

| Parameter | Florasulam | 2,4 D Ester LV |
|---------------------|--|--|
| Concentration (g/l) | 6.25 | 300 |
| HRAC Group | B | O |
| Mode of Action | Inhibitor of acetolactate synthase (ALS inhibition) | Synthetic Auxins |
| Plant Uptake | Foliar, shoots, roots | Foliar |
| Activity | Systemic (xylem and phloem) | Systemic |
| Visual symptoms | Weeds will appear stunted and chlorotic, will often have necrotic leaf margins, and will exhibit gradual death | Twisting/distortion of stems and leaves followed by wilting, necrosis and death. |
| First symptoms | 2-4 Days | |

Direction for use: Mix with water and apply onto broad leaved weeds during early vegetative stages of weed growth (with 4 to 6 leaves).



Application rates:

| Time of application | Rate/ha in 200 L of water | Rate in 20 L of water |
|--|---------------------------|-----------------------|
| Early vegetative stages of weed growth (with 4 to 6 leaves). | 500 mL | 50 |



Find us on Facebook
ADAMA East Africa


Pre-harvest interval: 25 days

Please read the label carefully before use.

Application Window

From 5 Leaf stage to Booting stage

Skala BBCH



Manufactured by:



ADAMA

For further information please contact:

Robert Wachira

Tel: +254 713 463 937

www.adama.com/east-africa

Distributed by:



Grebe Enterprises Limited

Hoska Plaza, Pandit Nehru Road

P.O. Box 14828-20100

Nakuru, Kenya

Mobile: +254(0)722603054

www.grebeenterprises.co.ke

ADAMA