



ADAMA

CAPRI 2,4-D

Classics of protection for maize



HERBICIDE

A selective post-emergence herbicide for control of broadleaved weeds in maize mono crop



Fast, long-lasting effect
the complete death of weeds occurs after 15 days, and the product has a protective action period of up to 30 days.



Flexibility of use
Capri® is perfectly combined with other herbicides



Cost - effective
Saves the high cost of manual weeding



High efficiency
Wide range of sensitive broadleaved weeds



Safety of use
Safe in recommended dosages

*Available in 3 pack sizes:
20 L, 5 L and 1 L

Simply. Grow. Together.

ADAMA.COM/EAST-AFRICA

Capri 2,4-D controls a wide range of sensitive broadleaved weeds, among them:



BlackJack
Bidens pilosa

Datura
stramonium

Macdonalds eye
Galinsoga paviflora

Mallow

Mexican marigold
Tagetes minuta

Sow Thistle
Sonchus oleraceus

Wandering Jew
Commelina bengalensis

MODE OF ACTION:

CAPRI 600 SL contains 2,4-D that has systemic activity which hinders plant cell growth in newly forming stems and leaves of broad leaved weeds.

DIRECTIONS FOR USE:

Mix with water and apply onto broad leaved weeds during early vegetative stages of weed growth (four weeks after maize emergence).

APPLICATION TIMING AND TECHNIQUE:

| Crop | Time of application | Rate/ha in 100 L of water | Rate in 20 L of water |
|-------|-----------------------------------------------------------------|---------------------------|-----------------------|
| Maize | Early stages of weed growth (four weeks after maize emergence). | 1 L | 200 ml |

NOTE: Avoid spray drift to any other desirable vegetation. Do not spray during periods of high winds.

Manufactured by:



For further information please contact:
Robert Wachira
Tel: +254 713 463 937
www.adama.com/east-africa

Distributed by:



Grebe Enterprises Limited
Hoska Plaza, Pandit Nehru Road
P.O. Box 14828-20100
Nakuru, Kenya
Mobile: +254(0)722603054
www.grebeenterprises.co.ke



Find us on Facebook
ADAMA East Africa

