

**General information:** Rimon Supra 10SC is growth regulator insecticide based on benzoylphenyl urea compound for control of caterpillars at all stages on Roses.

**Mode of action:** Rimon 10SC has a strong stomach activity and slight contact activity. It does not have ovicidal activity, but gives a high mortality of first instars hatching from eggs laid on sprayed foliage. Use of Rimon Supra 10SC also reduces the number of offspring arising from susceptible insect species. Rimon Supra 10SC works by inhibiting chitin formation in the cuticle thereby causing abnormal cuticle formation or failure to mature in susceptible insect species.

**RECOMMENDATION FOR USE:**

CROP	TARGET PEST	APPLICATION RATE
Roses	Caterpillars	0.5-0.75L/Ha, 50–75ml/100L water, 10-15ml/20ltr of water

**Preparation of Spray Mixture:** Fill the spray tank with half amount of water. Add the required amount of Rimon Supra 10SC, then fill the water to the required level and mix or agitate thoroughly to uniformity. Ensure that the spray mixture is used the same day

**Application timing:** Rimon Supra 10SC has limited knock down activity and is relatively slow acting. Therefore, for best results With Rimon Supra 10SC is achieved when it's applied after the eggs and larva are cited on the crop. Apply at least two consecutive applications of Rimon Supra 10SC at 14 days interval to ensure full effectiveness. Do not use on gyphosphilla

**Applications Technique:** Rimon Supra 10SC works by stomach and contact action and is not systemic in the plant. As such it is essential for applicator to ensure that good spray coverage of the target crop foliage is achieved. Use a high spray volume of 800-1000L per Ha.

**Resistance Management:** Mode of action of Rimon Supra 10SC is different from that of most other insecticides and therefore it is useful component in Insect Resistance Management Programmes. Rimon Supra 10SC should be applied strictly at the recommended rates in alteration with insecticides that have different mode of action. **RE-ENTRY PERIOD:** Spayed area can be re-entered after 6 hours in the field or after the spray has dried in greenhouse conditions. **HAZARDS/PRECAUTIONS: Handling:** Wear protective clothing namely overalls, footwear, gloves, and face shield when preparing and using the product. Do not eat, drink or smoke when working with the product.

**Storage:** Keep out of reach of children. Store in tightly closed original container in a cool dry well ventilated place, under lock key so that unauthorized persons do not have access. Keep away from food, drinks and animal feeding stuffs. **Disposal:** Triple rinse, puncture or crush empty containers before disposal in a safe and authorized place in accordance with National regulations on chemical disposal. Do not re-use empty containers for any other purpose. Pesticide spray mixture or rinsate that cannot be used must be disposed where it cannot contaminate surface water and ditches

**Pesticide Leakage and Spillage:** In event of heavy spillage, collect the contaminated soil layer and bury in an authorized place away from water supplies. Soaking in sand, soil or sawdust may collect small spillage. Contaminated absorbent material must be stored in marked waste bins prior to disposal in authorized landfill site. The pesticide store must be decontaminated by washing with a lot of water.

**Wildlife and Environmental Protection:** Rimon Supra 10SC is not toxic to fish and other aquatic organism. It is harmful to birds, animals and relatively non-toxic to bees. Keep livestock and wildlife out of treated areas.

**FIRST AID INSTRUCTIONS:** If poisoning is suspected, contact the doctor, or poisons information center.

**Skin contact:** Remove affected clothing and wash all exposed skin area with soap and water, followed by warm water rinse. Consult a doctor in the event of any complaints.

**Eye contact:** Wash off with plenty of water (15 min.) If pain persists, consult an eye specialist.

**If inhaled:** Remove victim to fresh air. Consult a doctor in the event of any complaints.

**If swallowed:** Do not induce vomiting. Wash out mouth with plenty of water. Never give anything by mouth to an unconscious person. Consult a doctor in the event of any complaints.

**TOXICOLOGICAL INFORMATION:** This product is harmful if swallowed, inhaled or absorbed through the skin. May cause irritation of the skin, eyes, and mucous membranes and may cause skin sensitization.

**Symptoms of Poisoning:** Signs of poisoning in mammals include Eye irritation and may affect nervous system

**Antidote:** There is no specific antidote. **NOTE TO PHYSICIAN:** Treat symptomatically.

**In case of emergency call toll free number: 0800720021/0800730030.**


**NOTICE TO USERS:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions.

**WARRANTY:** Seller's guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.



# Rimon Supra<sup>®</sup>

## 10SC



Insecticide for control of Catepillars on Roses  
(Ni dawa inayoangamiza viwavi kwa mmea wa waridi)

**COMMERCIAL & AGRICULTURAL CLASS.**  
(KUNDI LA KILIMO NA BIASHARA.)  
**GUARANTEE (DHAMANA);** *Novaluron 100g/liter*  
**REGISTRATION NO. (NAMBARI YA USAJILI);**  
PCPB (CR) 1139

**READ THE LABEL BEFORE USING**  
(SOMA KIBANDIKO KABLA YA KUTUMIA)

**KEEP OUT OF REACH OF CHILDREN**  
(WEKA MBALI NA WATOTO)

**Shelf life:** 2 years from date of manufacture when stored in original unopened container in a cool dry place.  
**(Maisha Rafuni):** Miaka 2 baada ya kutengenezwa kama imewekwa kwenye chombo chake halisi hakijafunguliwa mahali pakavu pasipo na joto jingi).

**Mft date/Tarehe ya kutengenezwa:**  
**Batch No./Nambari ya Kifurushi:**  
**Expiry date/Tumia kabla ya:**

**In case of Emergency call toll free No.**  
(Kwa hali ya Dharura piga simu bila malipo-masaa 24):  
0800720021/ 0800730030(24hrs)



**Distributed By / Wasambazaji:**  
Amiran Kenya Ltd.  
Tel: 0719 095000  
Fert@amirankenya.com  
www.amirankenya.com

**Manufacturer / Watengenezaji:**  
Adama Makhteshim Chemical Ltd, Israel.

WHO CLASS II  
**Net Contents / Ujazo 1Lt**



**Maelezo:** Rimon Supra 10SC ni kiua dudu kilicho kundi la benzoylphenyl urea ina uwezo wa kuangamiza na kudhibiti viwavi katika kila hatua kwenye waridi.

**Jinsi Inavyofanya kazi:** Rimon Supra 10SC huangamiza wadudu iwapo imeliwa. Ingawa ina uwezo mdogo wa kuua kwa mguso. Pia inauwezo wa kuangamiza mayai ya wadudu lakini na huangamiza viwavi wachanga. Rimon Supra 10SC huangamiza wadudu kwa kuzuia ukuwaji wa ngozi.

**MAPENDEKEZO YA MATUMIZI:**

MIMEA	WADUDU	VIPIMO VYA MATUMIZI
Waridi	Viwavi	Lita 0.5-0.75/Hekta, mililita 50-75 kwa lita 100 za maji, mililita 10-15/lita 20 za maji

**Kutayarisha Dawa:**

Changanya kiasi kinachohitajika cha Rimon Supra 10SC pamoja na kiwango kidogo cha maji. Ongeza kiasi cha maji kilichosalia na uendelea kutingiza bomba au kukoroga mchanganyiko wa dawa na maji. Tumia mchanganyiko wa dawa siku hiyo uliotayarishwa.

**Wakati na Jinsi ya Kutumia:** Kwa vile Rimon Supra 10SC haina uwezo wa kuua mara moja. Matokeo ya hali ya juu itapatikana iwapo dawa itanyunyuziwa wakati mayai ya wadudu au viwavi vimeonekana kwa mimea. Nyunyiza mara mbili ukifuatana baada ya siku 14. Usitumie kwa 'Gyphosphilla'

**MUDA SALAMA WA KUNINGIA SHAMBA LILIONYUNYUZIVA DAWA:** Masaa 6 kwa shamba la kawaida Jumba la mimea "greenhouse". Baada ya dawa kukauka.

**HATARI NA TAHADHARI:**

**Matumizi:** Vaa ngao la kujikinga, viatu na vizuizi vya mikononi na osho (face shield), unapotayarisha mchanganyiko wa dawa na maji na unaponyunyiza dawa. Jihadhari usipumue mvuke wa dawa, mchanganyiko wa dawa usingie kwenye macho au kumwagikia ngozi na nguo ulizovalia. Usinyunyize dawa hii ukielekea upande upepo unatoka. Usivute sigara na usile wala kunyua wakati unapotumia dawa hii.

**Baada ya kazi:** Vua nguo za kazi na vizuizi vya kujikinga, oga mikononi na uso kwa kutumia sabuni na maji. Safisha bomba la kunyunyiza dawa na mavasi yaliyomwagikiwa na dawa.

**Kuhifadhi:** Hifadhi dawa hii katika chombo chake asili ambacho kimefunikwa sawa sawa na kuwekwa kwenye ghala la madawa lililodhinishwa. Nilazima watoto na watu wasionuhusiwa wakatazwe kuingia kwenye ghala la madawa.

**Utupaji:** Safisha mkoba wa dawa iliyoisha kwa kusunza mara tatu kisha utoboe au ubonyeze na utupilie au uzike mbali na maji ambapo ni salama na panapodhinishwa kulingana na sheria za utupaji. Mchanganyiko wa dawa ambao haujatumiwa au maji yaliyoosha vyombo ni lazima yatupiliwe mahali salama na ambapo pameidhinishwa. Dawa ambayo haiwasi kutumiwa ni lazima itupiliwe mbali mahali ambapo ni salama, panapokubalivaa, na mbali na maji. Usitumie mkoba wa dawa kwa matumizi mengine yoyote.

**Dawa Ikiwagikia au Kuvuja:** Chukua mchanga uliomwagikiwa na dawa, uzike mahali ambapo ni salama na mbali na maji. Dawa ikivunja, iloweshe kwenye sehemu ndogo ya mchanga au 'sawdust' kisha ufungie mahali ambapo ni salama kabla ya kutupilia mbali mahali ambapo pameidhinishwa. Ghala la madawa ni lazima lioshwe mara kwa mara ili kutoa takataka ya dawa iliomwagikia au kuvunja.

**MADHARA KWA MAZINGIRA:** Rimon Supra 10SC si sumu kwa samaki, viumbe vingine vinavyo ishi kwenye maji na nyuki. Inaweza kudhuru ndege na wanyama. Usichafue mazingira kwa kunyunyiza, kumwaga mabaki ya dawa, kutupa, au kusunza mkoba wa dawa. Usiruhusu mifugo na wanyama wa pori kuingia mahali dawa hii imenyunyuziwa

**MAELEZO YA HUDUMA YA KWANZA: Ikipumuliwa:** Toa mgonjwa mahali penye hatari na umpumzisha mahali ambapo ni salama. Kama mjeruhi anashida ya kupumua, msaidie apumue, kisha umpeleke kwa daktari.

**Ikiwagikia ngozi:** Vua mavazi ya kujikinga na uoge kwa kutumia maji safi na sabuni. Iwapo maumivu itaendelea, muone daktari. **Ikingia kwa macho:** Fungua macho na uyasafishe kwa maji safi yanayotiririka kwa muda wa dakika 15. Iwapo maumivu itaendelea, muone daktari. **Kimezwa:** Usimfanye mgonjwa aliye katika hali maauti kutapika. Muoshe mdomo, kisha umuone daktari. **PELEKA MUADHIRIWA KWA DAKTARI**

**MAELEZO KWA DAKTARI:** Hakuna tiba maalum. Tibu mgonjwa kulingana na dalili za maumivu ili asaidiwe kupumua na kufanya kazi kwa pigo la moyo. Matibabu mengine ya nyongeza ni yale ya kutoa sumu mwilini au ya kuharisha. **Wakati wa dharura piga simu bila malipo (masaa 24): 0800720021/0800730030**

**ILANI KWA MTUMIAJI:** Dawa hii sharti itumie kulingana na maagizo yaliyo kwenye kibandiko. Ni hatia chini ya sheria za madawa ya kuangamiza wadudu vabaribifu kutumia au kuhifadhi dawa hii katika hali isiyokuwa salama. **THIBITISHO:** Dhamaana ya muuzaji ni maelezo yaliyo kwenye kibandiko ambayo ni lazima yafuatwe kikamilifu. Mlinuzi lazima akubali hatari zozote zinazoweza kutokana na matumizi ya dawa hii kwa binadamu au mali na lazima aikubali dawa hii katika hali hiyo.