



# BIGO® 400SC

Fungicide for the control of Late Blight in Potatoes and Tomatoes, Downy mildew and Botrytis in Roses  
 (Kiukuvu cha kudhibiti 'late blight' kwenye viazi na nyanya, 'downy mildew' na 'botrytis' kwenye waridi)

**Manufactured by / Imetengenezwa na:**  
 Adama Makhteshim Ltd  
 P.O.Box 60, BEER-SHEVA, 84100 - ISRAEL

**Registered by/Imezajiliwa na:**  
 Adama Agriculture East Africa Ltd  
 P.O. BOX 40034 Nairobi 00100 Kenya  
 Web: www.adama.com/east-africa

**Distributed by/Wazambazaji:**  
 Amiran Kenya Ltd, Old Airport North Road  
 P.O.Box 30327-00100 Nairobi, Kenya Tel. 0800 720 720  
 Email: fert@amirankenya.com Web: www.amirankenya.com

Batch Number (Nambari ya kifurushii):  
 Date of Manufacture (Tarehe ya kutengenezwa):  
 Expiry date (Tumia kaba ya):


Shelf life: Two years from date of manufacture if kept in unopened original container and in a cool dry place.  
 Maisha Rafuni: Miaka miwili baada ya kutengenezwa kama imehifadhiwa kwenye chombo chake halisi ambacho hakijafunguliwa mahali pakavu na pasipo na joto.

**GUARANTEE (DHAMANA):**  
 Dimethomorph 200g/L and Fluazinam 200g/L

Commercial and Agricultural class • Kundi la bishara na kilimo  
 Registration number /Nomba ya Usajili: PCPB (CR)1490

READ THE LABEL BEFORE USING • KEEP OUT OF REACH OF CHILDREN  
 SOMA KIBANDIKO KABA YA KUTUMIA • WEKA MBALI NA WATOTO

1 Liter



OPEN HERE



# ADAMA



X  
HARMFUL  
(Inadharu)



# ADAMA



**MAELEZO:** BIGO® 400SC ni sumu inayoangamiza ‘late blight’ katika viazi na nyanya, pamoja na ‘downy mildew’ na ‘botrytis’ kwenye waridi.

**Jinsi inavyofanya kazi:** BIGO® 400SC inajumuisha dimethomorph na fluazinam ambazo zinazuia kuenea kwa ugonjwa za ‘fungi’ kwenye viazi na waridi.

**JINSI YA KUTUMIA:** Changanya na maji kulingana na kiwango kilichoandikwa kisha unyonyize kwenye majani ya viazi, nyanya na waridi, kabla ya ugonjwa kuonekana.

**Kipimo:**

Mmea	Ugonjwa	Wakati wa kунуnyizia katika msimu	Kipimo kwa lita 1000 za maji	Kipimo katika lita 20 za maji
Viazi	Late blight	Rudia mara tatu kila siku 7	Mililita 750	Mililita 15
Nyanya	Late blight	Rudia kila siku saba na usirudie zaidi ya mara nne	Mililita 750	Mililita 15
Waridi	Downy mildew na botrytis	Rudia mara tatu kila siku 10	Mililita 750	Mililita 15

**Muda kabla ya kuvuna:** Unatakiwa kungoja siku saba kabla ya kuvuna viazi na nyanya.

**Tahadhari ya utumiaji:** BIGO® 400SC ni dawa ya kuangamiza ‘fungi’ mashambani. Haistahili kutumiwa kwa njia nyingine.

**Hadhari inapotumia:** Weka mbali na watoto, wanyama na chakula cha watu na wanyama. Yaweza kuleta muwasho kwenye mapua, koo, macho na ngozi. Tumia ‘overall’, mipira ya mikono na vazi la kufunika uso. Usile, kunywa wala kuvuta sigara utumiapo dawa.

**Kuchanganya kwa usalama:** Vaa mavazi ya kujikinga na dawa unaponyunyizia. Usijichafue na maji ya dawa. Hakikisha upopo hauisukumi dawa upande wako.

**Usalama baada ya kutumia:** Nawa mikono na mwili vizuri. Usitumie mkebe mtupu kwa kazi ingine yoyote. Tupa vyema mkebe mtupu.

**Uhifadhi:** Weka na kufungia BIGO® 400SC mbali na watoto na wanyama. Weka mahali pakavu ambapo kipimo cha joto hakizidi 30°C. Weka mbali na moto na ndimi zake. Hakikisha dawa hazisongomani kwenye stoo.

**Kutupa:** Usitumie kifaa kilichokuwa na dawa hii kwa namna nyingine yoyote.



Osha mikebe mara tatu kwa kutumia maji kisha uitupe kulingana na sheria za utupaji.

**Ikivuja au Ikimwagika:** Funika au zika dawa iliyomwagika kwa mchanga au vumbi ya mbao. Kusanya mchanganyiko huo na uutupe kulingana na sheria za utupaji. Osha kabisa sakafu iliyochafuka kwa maji na sabuni.

**MADHARA KWA MAZINGIRA:** BIGO® 400SC yawenza kudhuru samaki. Usiruhusu dawa hii illoweke karibu na chemichemi za maji.

**MASHARTI YA HUDUMA YA KWANZA:**

Simamisha kazi mara moja. Chunguza kama kuna dalili za uchovu. Toka pahali palipo chafuliwa na dawa. Vua nguo ikiwa zimechafuliwa. **Ikimezwa**, osha mdomo kwa kutumia maji mengi. Usilazimishe muathiriwa kutapika. Usimpe mtu aliyezimia kitu chochote kupitia mdomo. Peleka muathiriwa hospitalini. **Ikiingia machoni**, osha mara moja kwa maji mengi kwa dakika 15. **Ikigusa ngozi**, osha kwa maji mengi na sabuni. Kisha osha tena na maji yasiyo na moto kabisa. **Ikivutwa wakati wa kupumua**, mwondoe mwadhiriwa na umweke mahali penye hewa safi. Hakikisha muathiriwa hahisi baridi. Tafuta matibabu zaidi.

**PELEKA MJERUHI KWA DAKTARI**

**Ikiwa kuna tukio la madhara piga simu bila malipo 0800 730030/720021 (masaa 24)**

**ILANI KWA MUTUMIAJI**

Dawa hii lazima itumike kulingana na maelezo yaliyoko katika kibandiko hiki. Ni hatia chini ya sheria za viuadudu kuitumia au huhifadhi dawa hii kwa njia nyingine kinyume na ile iliyoelezewa hapa.

**THIBITISHO:** Dhamana ya muuzaji imeshauriwa na masharti yaliyowekwa kwenye kijikaratasi na label. Muuzaji hahusiki kamwe na madhara yoyote yanayotokana na utumiaji wa dawa hii kwa njia zinazohatarisha maisha.



**DESCRIPTION:** BIGO® 400SC is a systemic fungicide for the control of late blight (*Phytophthora infestans*) in potatoes and tomatoes, as well as downy mildew and botrytis in roses.

**Mode of action:** BIGO® 400SC contains dimethomorph and fluazinam which prevent growth and development of fungal cells and infective structures.

**Directions for use:** Mix with water at the recommended rates and foliar apply on potatoes, tomatoes and roses before disease appears.

#### Application rates:

Crop	Target Disease	Frequency of application	Rate in 1000 L of water	Rate/20 L of water
Potato	Late blight	Repeat 3 times every 7 days	750 mL	15 mL
Tomato	Late blight	Repeat 4 times every 7 days	750 mL	15 mL
Roses	Downy mildew and botrytis	Repeat 3 times every 10 days	750 mL	15 mL

**Pre-harvest interval:** 7 days on potatoes and tomatoes.

**Re-entry interval:** On drying of spray for roses.

**Use limitations:** BIGO® 400SC is an agricultural fungicide. It should be used only as recommended.

**Handling precautions:** Keep out of reach of children. Do not ingest. It may cause irritation of nose, throat, eyes and skin. Do not breathe spray mist. Use overall, gloves and mask when handling. Do not eat, drink or smoke while handling the product.

**Safety when mixing:** Wear protective clothing when mixing. Avoid contamination from spray liquid by spillage and drift.

**Safety when spraying:** Wear protective clothing when spraying. Avoid contamination from spray liquid by spillage and drift.

**Safety after spraying:** Wash hands and body thoroughly after spraying. Do not reuse the empty container. Dispose the container safely.

**Storage:** Store BIGO® 400SC under lock and key and out of reach in a cool dry place away from food, feed, children and animals and from direct sunlight. Store below 30°C.



**Disposal of waste and empty containers:** Triple rinse the containers and puncture holes in them. Dispose in accordance with national legislation and regulations. Do not allow material to enter sewers or bodies of water. Do not re-use the empty containers for any other purpose.

**Leakage and spillage:** If leakage of the liquid or spray mix occurs, soak up material with sand, soil or sawdust. Collect this contaminated material and bury in a safe place away from ware sources.

**ENVIRONMENTAL HAZARDS:** BIGO® 400SC is toxic to fish. Do not contaminate of water sources.

**FIRST AID MEASURES:** If poisoning is suspected, stop work immediately. Remove victim from the area of contamination and if clothing is contaminated, remove it. **In case of contact with eyes**, wash with plenty of water for at least 15 minutes keeping the eyelids open. **In case of contact with skin**, wash thoroughly with soap and water, followed by a warm water rinse. **If inhaled**, move patient to fresh air. Keep victim warm and at rest. If not breathing, give artificial respiration. If breathing is difficult, give oxygen. Get medical attention. **If swallowed**, rinse mouth with plenty of water. Do not induce vomiting. Never give anything by mouth to an unconscious person. Get medical attention.

**TOXICOLOGICAL INFORMATION:**

**Symptoms of poisoning:** Redness in the eyes, irritation, tears. Allergic reactions may be observed.

**Antidote:** No specific antidote

**Note to the physician:** Treat symptomatically and give supportive therapy.  
**In case of poisoning call toll free 0800 730030/720021 (24 hours)**

**NOTICE TO USER:** This pest control product is to be used in accordance with the direction on this label. It is an offence under the pest control Products Act to use or store a pest control product under unsafe conditions.

**WARRANTY:** Seller's guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.



**Shelf life:** 2 years from date of manufacture if stored in a cool, dry, well-ventilated place in original unopened container.

**Maisha Rafuni:** Miaka miwili kutoka tarehe ya kutengenezwa ikiwa imehifadhiwa katika mkoba wake asili ambao haujafunguliwa pahali pakavu pasipo na joto jingi.

**Manufactured by / Imetengenezwa na:**

Adama Makhteshim Ltd  
P.O.Box 60, BEER-SHEVA, 84100 - ISRAEL

**Registered by/Imezajiliwa na:**

Adama Agriculture East Africa Ltd  
P.O. BOX 40034 Nairobi 00100 Kenya  
Web: [www.adama.com/east-africa](http://www.adama.com/east-africa)

**Distributed by/Wazambazaji:**

Amiran Kenya Ltd, Old Airport North Road  
P.O.Box 30327-00100 Nairobi, Kenya  
Tel. 0800 720 720  
Email: [fert@amirankenya.com](mailto:fert@amirankenya.com)  
Web: [www.amirankenya.com](http://www.amirankenya.com)

ADAMA