

General Information

Folpan 50 SC is a protectant fungicide that belongs to phthalimides group for the control of Downey mildew on Rose.

Mode of Action

Folpan 50SC is a contact fungicide that controls susceptible crop disease by binding to the sulphur-hydrogen bond of the fungi, and thereby interferes with respiration process in fungi.

Direction For Use

Crop	Disease	Rates	R.E.I
Roses	Downey mildew	2.0 litres/Ha, 200ml/100 L of water 40ml/20L of water	6 hours/ Restricted until the spray has dried

Preparation of spray mixture: Fill the tank with half the required amount of water, shake Folpan 50SC gently and add the required amount, and adjuvant if required, then mix thoroughly. Fill the tank with water to the required level.

Timing of application: Start applying Folpan 50 SC when the weather condition favours disease development in order to protect the crop. Make protective sprays of Folpan 50SC regularly at a 7-days interval; or at a short interval during periods of intense-shoot growth and when serious infection is likely.

Application Technique: Use a spray volume of 1000 to 1500L per Ha for ornamental crops depending on the size of crop canopy to ensure thorough coverage.

Resistance Management: To prevent risk of developing pest resistance, it is recommended that Folpan 50SC be alternated with products of different modes of action.

HAZARDS/PRECAUTIONS

Handling Folpan 50 SC: Wear protective clothing: overalls, footwear, gloves, and face shield when preparing and using the product. Do not eat, drink or smoke while using it. Avoid inhaling spray mist, contact with skin, eyes and clothing. Do not apply upwind and stay out of the spray mist.

Ater Wwork: Wash hands before breaks and at end of work. Change clothes and wash hands and face with soap and water thoroughly. Wash the spraying equipment and any contaminated clothing carefully.

Storage: Keep out of reach of children. Store in tightly closed original container in a cool dry and well ventilated place, under lock and key so that unauthorized persons do not have access. Keep away from food, drinks and animal feeds.

Disposal: Triple rinse the bottle and puncture or perforate it and dispose off in a landfill site approved for pesticides away from water resources and in accordance with National Regulators on disposal or return to stockist/Distributor. The rinsate should be disposed off away from surface water or avoid contamination. Do not re-use the empty container for any other purpose.

Leakage and spillage: Wear protective clothing. Use sand or sawdust to absorb the chemical and collect it using a shovel. Put in authorized disposing containers. Do not contaminate waterways and rivers. In case of large spills, consult the distributor.

Wildlife And Environmental Protection: Folpan 50 SC is toxic to fish and other aquatic organisms. Do not contaminate streams, rivers or waterways with the chemical or used packs. Livestock should not be allowed to feed on freshly sprayed herbage.

FIRST AID INSTRUCTIONS: If poisoning is suspected, contact the doctor, or Poisons Information Centre.

Skin contact: Take off clothing; wash the skin immediately with plenty of water and soap. **Eye contact:** Hold the eye open and rinse thoroughly with running water for 15 minutes. **If inhaled:** Move the patient to fresh air. If the subject is not breathing, give artificial respiration, preferably mouth-to-mouth and call the doctor immediately. **If swallowed:** Induce vomiting. Wash out mouth with plenty of water. Never give anything by mouth to an unconscious person. Seek medical advice immediately.

TOXICOLOGICAL INFORMATION

Symptoms of poisoning: Irritant to eyes; May cause sensitization by skin contact.

Antidote: There is no specific antidote known. **Note to physician/Treatment:** Treat symptomatic and supportively. If swallowed, perform gastric lavage and administer activated charcoal.

In case of emergency call toll free number (24hrs): 0800720021/0800730030.

NOTICE TO USERS: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions.

WARRANTY: Seller's guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.



Amiran
Folpan®
50 SC

A fungicide for the control of Downey mildew on Rose (Kiua kuuu kinacho kinga kuuu yenye unga kwenye waridi)

COMMERCIAL & AGRICULTURAL CLASS.

(KUNDI LA BIASHARA NA KILIMO.)

GUARANTEE (DHAMANA): Folpet.....500g/l

REGISTRATION NO. (NAMBARI YA USAJILI):

PCPB (CR) ..0537

**READ THE LABEL BEFORE USING
(SOMA KIBANDIKO KABLA YA KUTUMIA)**

**KEEP OUT OF REACH OF CHILDREN
(FUNGIA MBALI NA WATOTO)**

**In case of Emergency call toll free No.
(Wakati wa Dharura piga simu bila malipo-masaa 24):
0800720021 / 0800730030(24hrs)**

Distributor / Wasambazaji:
Amiran Kenya Ltd.
Tel: 0719 095000
Fert@amirankenya.com
www.amirankenya.com

Member Of
Agro Chemical
Association Of Kenya
Registration Number.
AAK 219



Warning (Itani)
Toxic to aquatic life,
may cause long term adverse effects
in the aquatic environment.



Manufacturer / Watengenezaji:
Adama Makhteshim Ltd, Israel

WHO
CLASS II

Net Contents / Ujazo 1Lt



**CAUTION
(TAHADHARI)**



Maelezo

Folpan 50 SC ni kiua kuuu kinacho kinga waridi kutokana na kuuu. Kiko katika kikundi cha futhalimidi.

Jinsi inavyofanya kazi:

Folpan 50 SC hufanya kazi kwa mgusu na kuinga kuuu kwenye waridi kwa kuharibu mbinu za upumuaji wa kuuu.

Kipimo

Mimea	Ugonjwa	Vipimo	Subiri
Waridi	Kuuu yenye unga	Lita 2/Hektari, Mililita 200/Lita 100, Mililita 40/Lita 20 za maji	Kutoidhinishwa hadi manyonyu ya kemikali ikauke

Utayarishaji wa kemikali: Jaza tangi nusu maji, tikisa Folpan 50 SC, ongeza kwenye maji kiasi kilicho pendekezwa na ukoroge vizuri. Jaza maji kwenye tangi kama inavyopendekezwa.

Wakati wa kunyunyizia: Folpan 50 SC hunyunyiziwa msimu wenye ugonjwa hujitokeza. Nyunyizia mara kwa mara baada ya siku 7 au kabla ya siku 7 kuisha; wakati wa uvamizi wa hali ya juu.

Jinsi ya kunyunyizia: Tumia maji kiasi cha lita 1000-1500 kwa hektari kulingana na kiwango cha majani kwenye mimea

Upinzani: Kuzuia upinzani wa kemikali kutokea, inapendekezwa kuwa Folpan 50SC itumiwe kwa kubadilisha na kemikali zenye mbinu tofauti za utenda kazi.

HATARI NA TAHADHARI

Matumizi: Vaa ngao la kujikinga, viatu na viziwi vya mikono na uso, unapotarisha mchanganyiko wa dawa na maji na unaponyunyiza dawa. Jihadhari usipumue mvuke wa dawa, mchanganyiko wa dawa usingie kwenye macho au kumwagikia ngozi na nguo ulizovalia. Usinyunyizie dawa hii ukielekea upande upepo unatoka. Usivute sigara na usile wala kunyua wakati unapotumia dawa hii.

Baada ya kazi: Vaa nguo za kazi na viziwi vya kujikinga, Nawa mikono kati ya mapumziko na hata baada ya kazi. Osha bomba uliyotumia kisha Oga na ufue mavazi uliyotumia kwa maji na sabuni.

Kuhifadhi: Hifadhi Folpan 50 SC kwenye chombo chake asili kichofungwa vyema na kukiweka pahali pakavu, pasipo na joto na panapo hewa vizuri-kwenye ghala lake lililofungwa ambalo watu wasioidhinishwa kuingia. Weka mbali na watoto, chakula, vinywaji na vyakula vya mifugo

Utupaji: Suiza mkoba mara tatu, kitobee kisha ukitupe sehemu ambayo imetengwa na kuidhinishwa mbali na sehemu zilizo na chemi chemi ya maji kulingana na sheria za utupaji au ukirejisha kwenye msambazaji. Msuzo unafaa umwagwe mbali na sehemu za maji kuzuia machafuko ya mazingira.

Ikiujia/mwagika: Vaa mavazi ya kinga. Zuia machafuko wa maji au mchanga. Fagia kwa kutumia unga wa mbao au mchanga. Kusanya kwa kijiko cha mchanga kasha weka ndani ya mikoba inayoidhinishwa kuondoa kemikali. Kwa wingi wa kemikali, fuata mashauri ya msambazaji.

TAHADHARI KWA MAZINGIRA: Folpan 50 SC ni sumu kwa samaki na viembe vya majini. Usichafue mito, mabawa au njia za maji. Usiwaruhusu wanyama kwenye sehemu zilizo nyunyiziwa kemikali hii.

MAELEZO YA HUDUMA YA KWANZA: Kwa ngonzi: Vaa nguo na kuoga mara moja kwa maji mengi na sabuni. **Kwa macho:** Fumbua jicho au macho na kushikilia gigubiko cha jicho kisha tirisha maji mengi ndani ya macho au jicho kwa muda wa dakika 15. Muone daktari. **Ikiwitwa:** Mpeleke mjeruhi palipo na hewa safi, ikiwa anaugumu akipumua, mpe hewa kwa kutumia mdomo wako na wake. Muone daktari.

Ikimezwa: Vtapishe. Tumia maji mengi kusukutua mdomo. Usimpe kitu chochote mjeruhiwa aliyezirai. Muone daktari. PELEKA MJERUHIWA KWA DAKTARI.

TIBA: Tibu kulingana na dalili za ugonjwa

Kwa hali ya dharura, piga simu bila malipo 0800720021/0800730030, masaa 24

ILANI KWA MTUMIAJI: Dawa hii sharti itumiwe kulingana na maagizo yaliyo kwenye kibandiko. Ni hatia chini ya sheria za madawa ya kuangamiza wadudu waharibifu kutumia au kuhifadhi dawa hii katika hali isiyu salama.

THIBITISHO: Dhamana ya muuzaji ni maelezo yaliyo kwenye kibandiko ambayo ni lazima yafuatwe kikamilifu. Mhunuzi lazima akubali hatari zozote zinazoweza kutokana na matumizi ya dawa hii kwa binadamu au mali na lazima akubali dawa hii katika hali hiyo.