



MAXIDOR 250 SC

GUARANTEE (DHAMANA):

Azoxystrobin 250g/L

Fungicide for the control of rust and ring spot in carnations, powdery mildew on roses and rust on French beans
 (Dawa ya kuthibiti powdery mildew kwenye waridi, rust na ring spot kwenye carnations pamoja na rust kwenye French beans)

Manufactured by/Imetengenezwa na:

ADAMA Makhteshim Ltd P.O.BOX 60 Beer Sheva
 84100 Israel. Web: www.adama.com/east-africa

Registered by/Imezajiliwa na:

Adama Agriculture East Africa Ltd P.O. BOX 40034 Nairobi
 00100 Kenya. Web: www.adama.com/east-africa

Distributed by/Wazambazaji:

CropTech (K) LTD, Phenom Estate 238 Off Langata Road
 P.O. Box 137-00618 Ruaraka, Nairobi.

Tel. +254 722 703 911

Shelf Life: 2 years from date of manufacture if stored in a cool, dry, well-ventilated place in original unopened container.

Maisha Rafuni: Miaka miwili kutoka tarehe ya kutengenezwa ikiwa imewekwa katika mkoba wake halisi ambao haujafunguliwa.



SCAN FOR
WEBSITE



IRRITANT
HUWASHA



FLAMMABLE
LIQUID



ADAMA
ESSENTIALS

1 Liter

97101649D X 201510



CAUTION / ILANI



ADAMA



Maelezo

Maxidor SC ni dawa ya kuthibiti powdery mildew kwenye waridi, rust na ring spot kwenye carnations pamoja na rust kwenye French beans.

Jinsi inavyofanya kazi

Azoxystrobin ni dawa za kuangamiza 'fungi' kwa kupitia majani na mizizi.

Jinsi ya kutumia

Changanya na maji kulingana na kiwango kilichoandikwa kisha unyunyize kwenye majani ya maua na French beans.

Kuchanganya na wakati wa kutumia

Weka kipimo kinachofaa cha dawa ya Maxidor SC kwenye maji na unyunyizie kwenye mimea ili kukinga dhidi ya fungi. Usitumie kwenye mimea ambayo tayari imeambukizwa ugonjwa. Tumia mara mbili peke yake katika msimu. Usitumie pamoja na dawa zingine za aina ya 'strobilurin'.

Mmea	Ugonjwa	Kipimo kwenye lita 1000 za maji	Kipimo kwa lita 20 za maji	Wakati wa kunyunyiza
Carnations	Fungi aina ya 'rust' na 'leaf spot'			Nyunyiza kabla ya dalili za ugonjwa kuonekana.
Waridi	powdery mildew	mililita 750	mililita 15	Nyunyizia mara mbili au tatu kwa msimu ukingoja siku 10 kabla ya kunyunyiza tena.
French beans	Rust			

Muda wa kungoja kabla ya kuingia shamba ya maua: Masaa sita.

Muda wa kungoja kabla ya kuvuna French beans: Siku 3

Uchanganyikaji na madawa mengine

Maxidor SC inaweza kuchanganywa na kemikali zingine lakini inastahili kufanya utafiti na majaribio kwanza kabla ya kutumia michanganyiko yoyote.

Hadhari inapotumia

Weka mbali na watoto, wanyama na chakula cha watu na wanyama.

Yaweza kuleta muwasho kwenye mapua, koo, macho na ngozi. Tumia 'overall', mipira ya mikono na vazi la kufunika uso. Usile, kunywa wala kuvuta sigara utumiapo dawa.



Kuchanganya kwa usalama

Epuka kugusana na dawa unapoichanganya. Vaa 'overall', mipira ya mikono na vazi la kufunika uso unapoichanganya.

Kunyunyizia kwa usalama

Vaa mavazi ya kujikinga na dawa unaponyunyizia. Usijichafue na maji ya dawa. Hakikisha upepo hauisukumi dawa upande wako.

Usalama baada ya kutumia

Nawa mikono na mwili vizuri. Usitumie mkebe mtupu kwa kazi ingine yoyote. Tupa vyema mkebe mtupu.

Uhifadhi

Weka na kufungia Maxidor SC mbali na watoto na wanyama. Weka mahali pakavu ambapo kipimo cha joto hakizidi 30°C. Weka mbali na moto na ndimi zake. Hakikisha dawa hazisongomani kwenye stoo.

Kutupa

Usitumie kifaa kilichokuwa na dawa hii kwa namna nyingine yoyote. Osha mikebe mara tatu kwa kutumia maji kisha utupe kulingana na sheria za utupaji.

Ikivuja au Ikimwagika

Funika au zika dawa iliyomwagika kwa mchanga au vumbi ya mbao. Kusanya mchanganyiko huo na utupe kulingana na sheria za utupaji. Osha kabisa sakafu iliyochafuka kwa maji na sabuni.



MASHARTI YA HUDUMA YA KWANZA

Simamisha kazi mara moja. Chunguza kama kuna dalili za uchovu. Toka pahali palipo chafuliwa na dawa. Vua nguo ikiwa zimechafuliwa. **Ikimezwa**, usilazimishe muathiriwa kutapika. Peleka muathiriwa hospitalini na uonyeshe daktari kibandiko hiki.

Ikiingia machoni, osha mara moja kwa maji mengi. Maumivu yakizidi, mpeleke mjeruhi kwa daktari.

Ikigusa ngozi, vua mavazi iliyoguzwa na uoshe kwa maji mengi na sabuni. Kisha osha tena na maji yasiyo na moto kabisa. Maumivu yakizidi, mpeleke mjeruhi kwa daktari.

Ikivutwa wakati wa kupumua, mwondoe mwadhiriwa na umweke mahali penye hewa safi. Hakikisha muathiriwa hahisi baridi. Tafuta matibabu zaidi.

PELEKA MJERUHI KWA DAKTARI

Ikiwa kuna tukio la madhara pigi simu bila malipo **0800 730030/720021** (masaa 24).

ILANI KWA MUTUMIAJI

Dawa hii lazima itumike kulingana na maelezo yaliyoko katika kibandiko hiki. Ni hatia chini ya sheria za viuadudu kuitumia au huhifadhi dawa hii kwa njia nyingine kinyume na ile iliyoelezewa hapa.

THIBITISHO

Dhamana ya muuzaji imeshauriwa na masharti yaliyowekwa kwenye kijikaratasi na label. Muuzaji hahusiki kamwe na madhara yoyote yanayotokana na utumiaji wa dawa hii kwa njia zinazohatarisha maisha.



Description

Maxidor is an agricultural systemic and translaminar fungicide for the control of rust and ring spot in carnations, powdery mildew on roses and rust on French beans.

Mode of action

Maxidor SC contains azoxystrobin, a fungicide with protective, curative, eradicative, translaminar and systemic properties. It inhibits spore germination and mycelia growth; it also shows anti-sporulant activity.

Directions for use

Mix with water at the recommended rates and foliar apply on carnations, roses and French beans.

Mixing and timing of application

Mix the required amount of Maxidor SC with sufficient water and do not apply more than twice each season. Do not mix with any other strobilurin fungicide during application.

Crop	Disease	Rate/ha in 1000 L of water	Rate/20 L of water	Timing of application
Carnations	Rust and ring spot	750 ml	15	Before onset of disease. Apply 2 – 3 times per season with 10 days between applications
Roses	Powdery mildew			
French beans	Rust			

Re-entry interval for roses and carnations: 6 hours

Pre-harvest internal on French beans: 3 days

Handling precautions

Keep out of reach of children. Do not ingest. It may cause irritation of nose, throat, eyes and skin. Do not breathe spray mist. Use overall, gloves and mask when handling. Do not eat, drink or smoke while handling the product.



Safety when mixing

Wear protective clothing when mixing. Avoid contamination from spray liquid by spillage and drift.

Safety when spraying

Wear protective clothing when spraying. Avoid contamination from spray liquid by spillage and drift.

Safety after spraying

Wash hands and body thoroughly after spraying. Do not reuse the empty container. Dispose the container safely and according to national regulations.

Storage

Store Maxidor SC under lock and key and out of reach in a cool dry place away from food, feed, children and animals and from direct sunlight. Store below 30°C.

Disposal of waste and empty containers

Dispose in accordance with national regulations. Do not allow material to enter sewers or bodies of water. Triple rinse the containers. Do not re-use the empty containers for any other purpose.

Leakage and spillage

If leakage of the liquid or spray mix occurs, soak up material with sand, soil or sawdust. Collect this contaminated material and dispose according to national regulations.

ENVIRONMENTAL HAZARDS

Azoxystrobin is toxic to aquatic organisms but has low toxicity to birds, bees and terrestrial organisms. Do not contaminate water bodies.



FIRST AID MEASURES

If poisoning is suspected, stop work immediately. Remove victim from the area of contamination and if clothing is contaminated, remove it. **In case of contact with eyes**, wash off with plenty of water for at least 15 minutes, including the area under the eyelids. If irritation persists, consult an eye specialist.

In case of contact with skin, remove affected clothing and wash all exposed skin area with soap and water, followed by warm water rinse. Consult a doctor if irritation persists. Wash contaminated clothing before re-use.

If inhaled, remove victim to fresh air. Keep victim warm and at rest. If breathing is irregular or stopped, give oxygen. Seek medical attention.

If swallowed, do not induce vomiting. Seek medical attention immediately and show this label.

TOXICOLOGICAL INFORMATION

Symptoms of poisoning: Vomiting, nausea, respiratory tract irritation, reddening of eyes, burning and itching sensations.

Antidote: No specific antidote

Note to the Physician: Treat symptomatically and give supportive therapy. If ingested perform gastric lavage and administer activated charcoal.

In case of poisoning call toll free 0800 730030/720021 (24 hours).

NOTICE TO USERS

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions.

WARRANTY

Seller's guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.



Manufactured by / Imetengenezwa na:
ADAMA Makhteshim Ltd
P.O.BOX 60 Beer Sheva
84100 Israel
Web: www.adama.com/east-africa

Registered by/Imezajiliwa na:
Adama Agriculture East Africa Ltd
P.O. BOX 40034 Nairobi 00100 Kenya
Web: www.adama.com/east-africa

ADAMA