



Sphinx Extra®

713WDG



Fungicide for control of Late blight on tomatoes and Downy mildew on Roses
(Kiua kuvu kinacho zuia kuvu aina za "late blight" kwa nyanya na kuvu aina ya "downy mildew" kwa waridi.)

**COMMERCIAL & AGRICULTURAL CLASS.
(KUNDI LA KILIMO NA BIASHARA.)**

GUARANTEE (DHAMANA):

Dimethomorph 113g/Kg + Folpet 600g/Kg

REGISTRATION NO. (NAMBAARI YA USAJILI):

PCPB (CR) 1115

**READ THE LABEL BEFORE USING
(SOMA KIBANDIKO KABLA YA KUTUMIA)**

**KEEP OUT OF REACH OF CHILDREN
(WEKA MBALI NA WATOTO)**

Shelf life: 2 years from date of manufacture when stored in original container in a cool dry place.

(Maisha Rafuni: Miaka 2 baada ya kutengenezwa kama imewekwa kwenye chombo chake halisi ambacho hakijafunguliwa mahali pakavu na pasipo na joto jingi).

In case of Emergency call toll free No.
(Kwa hali ya Dharura piga simu bila malipo): 0800720021/ 0800730030
(24hrs / Masaa 24)

Member Of
AAK
Reg. No.
AAK 219

Mfg date/Tarehe ya kutengenezwa:

Batch No./Nambari ya Kifurushi:

Expiry date/Tumia kabla ya:

Distributed By / Wasambazaji:

Amiran Kenya Ltd.

Tel: 0719 095000

Fert@amirankenya.com | www.amirankenya.com



Manufacturer/Watengenezaji:
Adama Makhteshim Ltd,
P.O. Box 60, Israel.
Tel: 97286296671.



Warning (Ilani)

Very toxic to aquatic life,
may cause long term adverse effects
in the aquatic environment

Net Contents / Ujazo **1Kg**

WHO CLASS III



**CAUTION
(TAHDHARI)**



Sphinx Extra[®] | 713WDG

General Information: Sphinx Extra 713WDG is a Contact and Systemic fungicide with both Protective and Curative action. Sphinx Extra 713WDG is combination of two active ingredients; *Dimethomorph* and *Folpet* and it involves two different modes of actions, which are used to control downy mildew (*Plasmopara Viticola*) in Roses and Blight in tomatoes.

Mode of Action: *Dimethomorph* inhibits the formation of the Oomycete fungal cell wall affecting the sterol biosynthesis. It has local systemic (translaminar) properties with good protectant and antisporeulant activity. *Dimethomorph* moves acropetally up the treated plant stem and into growing leaves. It has antisporeulant activity prevents the spread of fungi attack.

Folpet has a protective contact action. It binds to sulphur hydrogen bonds that interferes with the fungi respiration process.

Recommendation for use:

Crop	Disease	Application rate
Roses	Downy mildew	2 - 2.5Kg/Ha, 200 - 250g/100L of water, 40 - 50g/20L of water.
Tomatoes	Late blight	2Kg/Ha, 200g/100L of water, 40g/20L of water.

Pre-harvest interval: Tomatoes: 7 days

Re-entry period: 6 hours.

Preparation of Spray Mixture

Half fill the spray tank (knapsack) with water. Add the required amount of Sphinx Extra 713WDG and mix by stirring or agitating, then fill the water to the required level and continue with stirring or agitating until the mixture is thoroughly uniform.

Application timing and technique

Start application at first signs of disease occurrence, Repeat after every 7-14 days as necessary. Use a spray volume of 1,000 to 1,500 L per Ha for roses and 1000L per Ha on tomatoes depending on the size of crop canopy to ensure thorough coverage. Do not closely apply Sphinx Extra 713WDG after an oil application.

Hazards/ precautions

Handling

Wear suitable protective clothing, gloves, face shield/safety goggles and respirator when preparing the spray mixture and in addition impermeable boots when spraying. Avoid, breathing the spray mist, contact with skin, eyes and clothing. Do not apply upwind and stay out of the spray mist. Do not smoke, drink or eat while handling the product.

After work

Change protective clothing and wash hands thoroughly with soap and plenty of water. Wash the spraying equipment and the contaminated clothing carefully.

Storage

Keep out of reach of children. Store in tightly closed original container in a cool dry well ventilated place, under lock key so that unauthorized persons do not have access. Keep away from food, drinks and animal feeding stuffs.

Disposal

Triple rinse, puncture or crush empty pouches before disposal in a safe and authorized place in accordance with National regulations on chemical disposal. Do not re-use empty containers for any other purpose. Pesticide spray mixture or rinsate that cannot be used must be disposed where it cannot contaminate surface water and ditches.

Pesticide Leakage and Spillage

In event of heavy spillage, collect the contaminated soil layer and dispose in an authorized place away from water supplies. Soaking in sand, soil or sawdust may collect small spillage. Contaminated absorbent material must be stored in marked waste bins prior to disposal in authorized landfill site. The pesticide store must be decontaminated by washing with a lot of water.

Environmental hazards

Sphinx Extra 713WDG is toxic to fish, other aquatic organism. It is moderately harmful to birds, animals and relatively non-toxic bees. Do not contaminate ponds and waterways by direct application, cleaning of equipment, disposal of wastes and empty container. Keep livestock and wildlife out of treated areas.

FIRST AID INSTRUCTIONS

Skin contact: Remove contaminated clothing and wash off with plenty of water and soap. Consult the doctor in case the complaint persists. **Eye contact:** Rinse thoroughly with running water for at least 15 minutes. If pain persists, consult an eye specialist. **If inhaled:**

Remove the patient to fresh air. Keep the patient warm and at rest. If the patient is not breathing, apply artificial respiration and seek medical attention. **If swallowed:** Do not induce vomiting. Wash out mouth with plenty of water. Never give anything by mouth to unconscious person. Consult a doctor.

TOXICOLOGICAL INFORMATION

Sphinx Extra 713WDG is slightly irritating to the skin and harmful when swallowed.

Symptoms of Poisoning: Irritation of the mucous membranes and skin. May cause ataxia and temporary sensitization may occur in certain individuals characterised by eye redness, bronchial irritation and skin rash.

Antidote: There is no specific antidote.

NOTE TO PHYSICIAN: Remove the victim to fresh air. Keep the victim warm and at rest and give symptomatic and supportive therapy.

In case of emergency call toll free number: 0800720021/0800730030.

NOTICE TO USERS

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions.

WARRANTY

Seller's guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

Maelezo

Sphinx Extra 713WDG ni kiuu kuuu kinacho angamiza kuuu kwa njia ya mguso na kusambaa yenyen mchanganyiko wa *Dimethomorph* na *Folpet* ambayo uponya na kuzua kuuu aina ya 'downy mildew' kwenye mimea ya waridi na kuuu aina ya 'late blight' kwa nyanya.

Jinsi Inavyofanya kazi

Dimethomorph, huzua utengenezaji wa kuta za chebechebe za kuuu na kuitafiana na na "sterol Biosynthesis". *Dimethomorph* usambaa kuanzia kwa shina ya mmea mpaka kwenye sehemu zinazokua za matawi na kuzua kusambaa kuuu.

Folpet, Ni dawa ifanyayo kazi ya kuzuua magonjwa kwa njia ya mguso. Huangamiza magonjwa ya mimea kwa kuitafiana na utengenezaji wa nguvu kwenye viumbe nya kuuu vinavyo sababisha ugonjwa.

Mapendekezo ya matumizi

Mimea	Ugonjwa	Vipimo vya matumizi
Waridi	Downy mildew	Kilo 2 - 2.5/Hekta, gramu 200 - 250/Lita 100 za maji, gramu 40 - 50 /lita 20 za maji
Nyanya	Late blight	Kilo 2 /Hekta, gramu 200/Lita 100 za maji, gramu 40 /lita 20 za maji

Muda salama wa kuvuna: Nyanya: siku 7

Muda salama wa kuingia shamba lilionyunyiziwa dawa: Masaa 6

Kutayarisha Dawa

Jaza bomba la kunyuyizia dawa maji kiasi nusu, kisha ongeza kiwango kinachohitajika cha Sphinx Extra 713WDG na huku ukikoroga na kujaza kiwango cha maji iliyosalia. Endelea kukoroga mchanganyiko wa dawa hadi uwe sawasawa.

Wakati na Jinsi ya Kutumia

Anza kunyuyizia dawa wakati dalili za ugonjwa zinapodhihirisha. Rudia kila siku 7-15 kulingana na dalili na ali ya anga. Tumia maji kiasi cha lita 1000 – 1,500 kwa hektaa unaponyuyizia mimea ya waridi na lita 1000 kwa hektaa unaponyuyizia mimea ya nyanya kulingana na kanopi ya matawi. Usitumie Sphinx Extra 713WDG baada ya kunyuyiza madawa yaliyo na aina za mafuta.

Hatari na tahadhari

Matumizi: Vaa ngao la kujikinga, viatu na vizuizi nya mikono na osho (face shield), unapotayarisha mchanganyiko wa dawa na maji na unaponyuyiza dawa. Jihadhari usipumue mvuke wa dawa, mchanganyiko wa dawa usingie kwenye macho au kumwagikia ngozi na nguo ulizovalia. Usinyunyizie dawa hii ukielekeupande upepo unatoka. Usivute sigara na usile wala kunya wa kati unapotumia dawa hii.

Baada ya kazi: Vua nguo za kazi na vizuizi nya kujikinga, oga mikono na uso kwa kutumia sabuni na maji. Safisha bomba la kunyuyizia dawa na mavasi yaliyomwagikiwa na dawa.

Kuhifadhi: Hifadhi dawa hii katika pouchi yake asili ambacho kimefunkika sawa sawa na kuwekwa kwenye ghala la madawa lilioidhinishwa. Ni lazima watoto na watu wasioruhusiwa wakatazwe kuingia kwenye ghala la madawa.

Utupaji: Safisha pouchi ya dawa iliyoisha kwa kusuza mara tatu kisha utoboe au ubonyeze na utupilie mbali na maji ambapo ni salama na panapoidhinishwa kulingana na sheria za utupaji. Mchanganyiko wa dawa amba haujatumika au maji yaliyoasha vyombo ni lazima yatupiliwe mahali salama na ambapo pameidhinishwa. Dawa ambayo haiwesi kutumiwa ni lazima itupiliwe mbali mahali ambapo ni salama, panapokubaliwa, na mbali na maji. Usitumie pouchi ya dawa kwa matumizi mengine yoyote.

Dawa Ikimwagika au Kuvuja: Chukua mchanga uliomwagikiwa na dawa, utupe mahali ambapo ni salama na mbali na maji. Dawa ikivunja, iloweshe kwenye sehemu ndogo ya mchanga au 'sawdust' kisha ufungie mahali ambapo ni salama kabila ya kutupilia mbali mahali ambapo pameidhinishwa. Ghala la madawa ni lazima lioshwe mara kwa mara ili kutoa takataka ya dawa iliomwagika au kuvunja.

Madhara kwa mazingira

Sphinx Extra 713WDG ni sumu kwa samaki na viumbe vingine nya maji. inaweza kudhuru ndege na wanyama. Haina madhara kwa nyuki. Usichafue mazingira kwa kunyuyiza, kumwaga mabaki ya dawa, kutupa au kusunza mkoba wa dawa kwa mito, visima au vidimbwi na chemichemi.

MAELEZO YA HUDUMA YA KWANZA

Ikipumuliwa: Toa mgonjwa mahali penye hatari na umpumishe mahali ambapo ni salama. Kama mjeruhi anashida ya kupumua, msaidie apumue, kisha umpeleke kwa daktari. **Ikimwagikia ngozi:** Vua mavazi ya kujikinga na uoge kwa kutumia maji safi na sabuni. Iwapo maumivu itaendelea, muone daktari. **Ikingia kwa macho:** Fungua macho na uyasafishe kwa maji safi yanayotiririka kwa muda wa dakika 15. Iwapo maumivu itaendelea, muone daktari. **Ikimezwia:** Usimfanye mgonjwa aliye katika hali maututu kutapike. Muoshe mdomo, kisha umuone daktari. PELEKA MJERUHIWA KWA DAKTARI

Maelezo kuhusu sumu

Sphinx Extra 713WDG inawasha ikimwangikia ngozi na ni sumu ikimezwia.

Dalili za kusumika: Inaweza kusababisha unyonge wa mwili, kufa ganzi, kupoteza hamu ya chakula, kuwashwa kwa umeo la pumzi na vipelepele katika sehemu zilizo guswa na dawa.

Maelezo kwa daktari: Hakuna tiba maalum. Tibu mgonjwa kulingana na dalili za maumivu.

Wakati wa dharura piga simu bila malipo (masaa 24): 0800720021/0800730030

ILANI KWA MTUMIAJI: Dawa hii sharti itumiwe kulingana na maagizo yaliyo kwenye kibandiko. Ni hatia chini ya sheria za madawa ya kuangamiza wadudu waharibifu kutumia au kuhifadhi dawa hii katika hali isyo salama.

THIBITISHO: Dhamana ya muuza ni maelezo yaliyo kwenye kibandiko ambayo ni lazima yafuatwe kikamilifu. Mnunuzi lazima akubali hatari zozote zinazoweza kutokekana na matumizi ya dawa hii kwa binadamu au mali na lazima aikubali dawa hii katika hali hiyo.