

# PYRINEX

A broad spectrum  
Insecticide/Acaricide  
for wood treatment,  
agricultural use,  
public health &  
termite control



SCAN FOR  
WEBSITE



IRRITANT  
HUWASHA



FLAMMABLE  
LIQUID

**GUARANTEE:** Chloropyrifos 480 g/Lt, EC  
(o,o-Diethyl 0-3, 5-6-trichloro-2-pyridyl phosphorothioate)

Commercial and Agricultural class F.P. 32, UN NOS:1993  
Registration number: UgC/2011/000824/In/RRRR

READ THE LABEL BEFORE USING • KEEP OUT OF REACH OF CHILDREN

## 1 Liter

97070565A/201504

### GENERAL INSTRUCTIONS

PYRINEX 48 EC is an organophosphate insecticide used for the control of a wide range of insect pests in the field, Public health and termite control.

PYRINEX 48 EC is compatible with most commonly used pesticides, but should not be mixed with Eradicane.

If in doubt about compatibility always consult the supplier.

### PRECAUTIONS

- Handle with care: avoid splashing/contact: poisonous if swallowed, inhaled or contacts with the skin.
- Wear full protective clothing, i.e. rubber gloves, rubber boots, overall, hood and respirator.
- Do not eat, drink or smoke while applying this insecticide.
- Remove protective clothing on completion of spraying and wash hands and face thoroughly with soap and water.
- Dangerous to animals, birds, fish and bees.
- Do not contaminate drinking pools, dams, rivers and waterways
- Dangerous to livestock and domestic pets.
- Keep out of reach of children.
- Keep away from food and foodstuffs.
- Store in original container and keep under lock and key.
- Destroy empty containers - do not use for any other purpose.

### WARNINGS

Avoid spray drift onto crops. Minimum number of days from last application to harvesting -14 days.

### SYMPTOMS OF POISONING

These may include excessive sweating, headache, weakness, faintness and giddiness, nausea, stomach pains, small pupils, blurred vision, muscle twitching.

### FIRST AID

If symptoms of poisoning occur, particularly if there is skin contamination or gross exposure, remove contaminated clothing without delay and wash affected skin.

Do not induce vomiting but obtain medical attention immediately. Never give anything by mouth to an unconscious person. Keep patient at rest lying in a face down position and ensuring a clear airway. If breathing has stopped apply artificial respiration. If eyes are contaminated, flush well with clean water.

### NOTICE USERS

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe corruption.

### WARRANTY

Seller's guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or properly arising from the use or handing of this product and accepts the product on that conation.

### DIRECTIONS FOR USE

CROP	PEST	RATE	REMARKS
Wood treatment	Complex wood pests	0,5%-1%	
Maize	Cutworms False wirewoms Black maize beetle	100 ml/100 lt water (1 lL/ha) 10 ml/100 m plant furrow	Use 500 lt. of water/ha Spray soil surrounding plants.
Potatoes	Cutworms, False wirewoms Tuber moth, Army worm- Aphids	10 ml/100 m plant furrow 1.5-2 lt/ha	Spray soil surrounding plants Use 200 - 300 lt. water/ha
Tomatoes	Tomatoes grub. Green vegetable grub Green peach aphid Cutworms, False wirewoms	1.5 - 2 lUhaor 150 - 200 ml/100lt. water	Spray interval: 7-10 days Use high rate under heavy pest pressure.
Tobacco	Cutworms, False wite worms, white grubs	100-150 ml/100 lt. water	Apply 400 -500 lt/ha
Cole crops	Aphids, Diamond back moth, Cabbage white butterfly	150 - 200 ml/100 lt. water or 1.5 - 2 lt/ha	Spray interval: 10-14 days Use high rate under heavy pest pressure
Banana	Thrips Banana Scab moth	75 -100 m/l 100 lt. water 200 m/100 lt. water	
Beans	Black maize beetle	10 ml/100 m plant furrow	Use 3 lt. warter/100 m furrow
Citrus	American bollworm, aphids' Orange dog, Mealybugs	40 - 50 ml/100 lt. water 100 ml/100 lt. water	
Lettuce	Cutworms	1lt/ha, 0.1 lt/100 lt. water	
Ornamentals Rowers Lawns	Crickets Mealybugs Caterpillars, Aphids	50 ml/100 lt. waiter 100 ml/100 lt. water 50 ml/100 lt. water or 250 ml/ha	Full cover spray Full cover spray Repeat when necessary
Pineapples	Pineapple Mealybugs, Ants	100 ml/100 lt. water	Use 3000 lt. water/ha
Sugar Cane	Common Armyworm, locust	700-900 ml/ha	Use higher rate when larvae over 3 cm.
Cereals	Cutworms, wireworms	700 ml/ha	Use min. 100 lt. water/ha
Public health	Cockroaches Spiders, Ants Mosquitoes Mosquito Polluted water Termites	50 ml/10 lt. water 100 ml/10 lt. water 50 -100 ml/ha 20 ml 100 m³water 4 lt/100 lt. water	

### Manufactured by:

ADAMA Makhteshim Ltd

P.O.BOX 60 Beer Sheva

84100 Israel

Web: [www.adama.com](http://www.adama.com)



HARMFUL

