



FARMURON

Herbicide for the control of numerous annual broad-leaved and grass weeds in potatoes, carrots, onions, beans and maize.

Dawa inayochagua na kuua kwekwe zenye majani mapana na pia zile za nyasi kwenye mashamba ya mimea kama vile viazi, karoti, vitunguu, maharagwe na mahindi.



HERBICIDE



GUARANTEE (DHAMANA): Linuron 500 gr/kg, WP

Commercial and Agricultural class • Kwa biashara na kilimo
Registration number/ Namba ya Usajili: TPRI HE/0251

READ THE LABEL BEFORE USING • KEEP OUT OF REACH OF CHILDREN
SOMA KIBANDIKO KABLA YA KUTUMIA • WEKA MBALI NA WATOTO

15 Kg

ADAMA



CAUTION



ADAMA



MAELEZO KWA JUMLA

FARMURON®50WP ni dawa inayochagua na kuuu magugu yenye majani mapana na pia nyasi kwenye mashamba ya mimea kama vile viazi, karoti, vitunguu, maharagwe na mahindi.

JINSI YA KUTUMIA

Viwango vya kutumia

Mimea	Kiwango (Kilo/Hektari)	Kiwango (Gramu kwa Lita 20 za maji)	Maelezo
Viazi	Kilo 2-3	80-120g.(Gramu 60 - 90/ lita 15 za maji)	Nyunyuzla baada ya kutengenaza misingi kabla ya viazi kuota
karoti	Kabla ya kuota Kilo 1.5-2.0	60-80g.(Gramu 45 - 60/ lita 15 za maji)	Nyunyiza muda tu baada ya kupanda. Kabla ya mmea kuota
	Baada ya kuota Kilo 1.5-2.0	60-80g.(Gramu 45 - 60/ lita 15 za maji)	Nyunyiza ikiwa mmea umetoa majani matatu
Vitunguu (baada ya kupanda miche)	Kilo 1.0-1.5 (Kiwango cha chini kwa mchanga mwepesi na kiwango cha juu kwa mchanga mzito)	40-60g.(Gramu 30 - 45/ lita 15 za maji)	Nyunyiza siku 10-14 baada ya kupanda miche
Maharagwe na Njegere	Kabla ya kuota Kilo 1.5-2.0	60-80g.(Gramu 45 - 60/ lita 15 za maji)	Nyunyiza mare tu baada ya kupanda. Isipite siku moja baada ya kupanda
Mahindi	Kabla ya kuota Kilo 1.5-2.0	60-80g.(Gramu 45 - 60/ lita 15 za maji)	Nyunyiza muda tu baada ya kupanda

Tumia kifaa ulichopewa kupimia dawa. Kiwango cha lita 500 za maji kwa hektari kimetumiwa ili kujua kiwango cha FARMURON®50WP kinachohitajika kwa lita 20 za maji.

Matumizi: Nyunyiza FARMURON®50WP kwenye shamba lililopandwa, lenya unyevu na lililotengenezwa vizuri, kabla ya mimea kuota. Panda mbegu chini inavyowezekana. Hakikisha mbegu zimefunikwa vizuri na udongo. Kwa matumizi baada ya mimea kuota, nyunyiza dawa kwenye magugu change (zimetoea majani manne). Usinyunyize dawa baada ya mimea kuota kama mvua inatarajiwa au kama bado magugu yana maji ya mvua au unyevu. Changanya kiwango kinachohitajika cha FARMURON®50WP na lita 500 hadi 1000 za maji kwa hektari moja.

Jinsi ya kunyunyiza: Baada ya utayarishaji wa FARMURON®50WP inyunyize kama ifuatavyo:

Unyunyizaji kabla ya mimea kuota kwenye shamba la viazi, mahindi, maharagwe na njegere, nyunyiza dawa vizuri kwenye shamba lililotengenezwa vizuri. Hakikisha kuwa mbegu zimefunikwa kabisa na udongo.

ADAMA



Unyunyiziaji kabla na baada ya karati kuota: Kwa unyunyiziaji kabla ya karati kuota, nyunyiza FARMURON®50WP muda tu baada ya kupanda. Udongo lazima uwe na unyevu.

Kwa unyunyiziaji baada ya karati kuota, nyunyizia dawa kama mmea umetoa majani mawili hadi manne.

Kiasi Cha Maji Kinachohitajika

Lita 50-100 kwa kila hekta moja kwa unyunyiziaji wa angani (ndege).

Lita 200-500 kwa kila hekta moja kwa mabomba yanayotumia tingatinga.

Lita 500-1000 kwa kila hekta moja kwa kutumia bomba la kubebea mgongoni k.m solo

JINSI YA KUTAYARISHA DAWA

Weka maji hadi nusu ya tenki. Anza kukoroga. Weka kiasi kinachotakiwa cha FARMURON® 50WP ndani ya tenki huku ukiendelea kukoroga na kujaza maji hadi tenki lijae.

Kwa matumizi. Bora, tayarisha dawa na utumie siku hiyohiyo huku ukihakikisha kuwa umenyunyiza sawasawa kote shambani.

UPATANISHI

FARMURON®50WP Inaweza kuchanganywa na dawa zingine za kuua magugu kama Atachlor, Atrazine Bromoxynil na Diuron.

HADHARI

Tumia dawa hii kwa uangalifu. Soma na unzingatie maelezo kamili ya matumizi wa dawa hii. Inadhuru ikimezwa. Inaweza kuwasha macho. Usiwe karibu na dawa kwa muda mrefu. Epuka kuupumua ukungu unapo nyunyiza. Valia mavazi ya kinga pamoja na miwani na glavu. Usile, kunywa au kuvuta sigara wakati unapotumia dawa hii.

Baada ya kutumia dawa:

Osha glavu vizuri na unawe mikono kwa sabuni na maji. Badilisha nguo.

KUHIFADHI

Hifadhi ndani ya chombo chake halisi pahali pasipo na joto na pakavu, weka mbali na watoto, vyakula vya binadamu na wanyama.

KUTUPA VYOMBO VILIVYOTUMIKA

Usitumie tena vyombo vilivyoisha dawa, vichome au uzike. Usihifadhi mchanganyiko wa dawa hii ya kunyunyiza.

MAELEZO YA HUDUMA YA KWANZA

Ikiingia machoni: Suza macho kwa maji mengi kwa muda wa dakika 15. Mwasho ukizidi, pata ushauri wa daktari na umuonyeshe kibandiko.

Ikimwagika kwenye ngozi: vua nguo zilizo chafuliwa na dawa. Osha sehemu hiyo ya mwili kwa kutumia maji mengi na sabuni. Mwasho ukizidi pata ushauri wa daktari na umuonyeshe kibandiko



Ikimezwa: Mpe mgonjwa makaa yaliyosagwa na kuchanganywa na maji. Usimpe mgonjwa maziwa, vyakula vya mafuta au pombe. Usimpe mgonjwa chochote kwa mdomo ikiwa amezlrai. Mwasho ukizidi pata ushauri wa daktari na umuonyeshe kibandiko.

MAELEZO YA SUMU/ TAARIFA KWA DAKTARI

Dalili za sumu: Hakuna visa vyovyote vimeripotwa kuhusu sumu ya dawa hii kwa wanadamu.
Makata: Hakuna makata inayojulikana.

TIBA

Tibu kulingana na dalili mgonjwa anazoonyesha.

Wakati wa dharura piga simu bila malipo (Masaa 24): 00800720021/0800730030

ILANI KWA MTUMIAJI.

Dawa hii lazima itumiwe kulingana na maagizo yaliyo kwenye kibandiko.

Ni hatia chini ya sheria za madawa ya kuangamiza wadudu waharibifu, kutumia au kuhifadhi dawa hii katika hali isiyu salama.

THIBITISHO

Dhamana ya muuzaji ni maelezo yaliyo kwenye kibandiko ambayo ni lazima yafuatwe kikamilifu. Mnunuzi lazima akubali hatari zozote zinazoweza kutokana na matumizi ya dawa hii, kwenye binadamu au mali na lazima aikubali dawa hii katika hali hiyo.



GENERAL INFORMATION

FARMURON® 50WP is a selective pre - and post - emergence herbicide for control of many annual broad-leaved weeds and grass weeds in several crops such as potatoes, carrots, onions, beans and maize.

DIRECTIONS FOR USE

Crops and Rates of Application

Crops	Dosage (Kg/Ha)	Dosage (20 Lt)	Remarks
Potatoes	Pre-emergence 2-3 Kg	80-120 g (60-90 g/15 Lt)	Apply after final ridging before emergence of potatoes
Carrots	Pre-emergence 1.5-2.0 Kg	60-80 g (45-60 g/15 Lt)	Apply soon after sowing
	Post-emergence 1.5-2.0 Kg	60-80 g (45-60 g/15 Lt)	Treat at 3 true leaf stage
Onions (Only after transplanting)	1.0-1.5 Kg (low rates for light soils and high rates for heavy soils)	40-60 g (30-45 g/15 Lt)	Spray 10-14 days after transplanting
Phaseolus beans and peas	Pre-emergence 1.5-2.0 Kg	60-80 g (45-60 g/15 Lt)	Apply immediately and NOT later than 1 day after sowing
Maize	Pre-emergence 1.5-2.0 Kg	60-80 g (45-60 g/15 Lt)	Apply soon after sowing

Use the provided measuring cap. 500 Lt of water per hectare has been considered when calculating quantities of FARMURON 50WP required per 20 Lt Water.

APPLICATION

Apply FARMURON® 50WP preferably as a pre-emergence, spray evenly over a moist and well-prepared seedbed. Plant crop as deep as possible making sure that the seed is covered by soil. With post emergence treatment apply the product to germinating actively growing young weeds, (up to 4-leaf stage). Do not apply post emergence if rainfall is expected, or if weeds are still wet from rain or dew. Mix the recommended quantity of FARMURON® 50WP in 509-1009 liters of water per hectare. As a pre-emergence treatment apply the herbicide EVENLY over MOIST soil and well-prepared seedbed.



HOW TO APPLY FARMURON 50WP

After preparation of FARMURON 50WP spray suspension, apply the herbicide as follows:

Pre-emergence spray on potatoes, maize, beans and peas. Apply the herbicide **EVENLY** over a **MOIST** and well-prepared seedbed making sure that the seed is well covered with soil.

Pre and post emergence treatment on carrots: As a pre-emergence treatment apply FARMURON® 50WP on carrots soon after planting. The soil **MUST** be **MOIST**.

As post emergence treatment, apply the herbicide when the carrots are at 2-4 true leaf stage.

Recommended water volumes

50 -100 Lt/ha for aerial applications

200-500 Lt/ha for tractor mounted spraying machines

500-1000 Lt/ha for knapsack sprayers with flatten nozzles e.g. Teejet

PREPARATION OF SPRAY

Fill half of the tank with water and start agitation. Pour the required amount of FARMURON 50WP on the water surface keeping the agitator running and continue filling the tank with water until the tank is full.

Use the spray mixture the same day of its preparation stirring well before use. Apply as a spray making sure that the mixture is evenly applied over the whole surface.

COMPATIBILITY

FARMURON® 50WP is compatible with other herbicides such as Alachbr, Atrazine, Bromoxynil, Diuron, Trifluralin and Hormone weed killers.

It is however advisable to carry out small trials first, before large-scale applications.

PRECAUTIONS

Handling

Carefully study and strictly observe directions for use. Harmful if swallowed. May cause irritation to eyes. Avoid prolonged exposure to the product. Do not breathe dust or spray mist. Wear protective clothing including goggles and rubber gloves. Do not eat, drink, or smoke when handling.

Afterwork

Wash gloves and hands with soap and plenty of water and change clothing.

STORAGE

Store product in closed original container in a cool dry place and keep away from children and food stuffs.

DISPOSAL

Do not re-use empty containers, burn or bury. Do not store spray mixture.

WILDLIFE AND ENVIRONMENTAL PROTECTION

Do not dispose of spray mixture near waterways. Toxic to fish.



FIRST AID INSTRUCTIONS

In case of eye contact: Flush eyes and skin with plenty of water for 15 minutes. In case of serious irritation, seek medical advice and show physician the label.

Skin contact: remove contaminated clothing. Wash affected areas with plenty of soap and water.

If swallowed: give slurry of activated charcoal in a glass of water. Avoid milk, fat, oil based laxatives and alcohol. Never give anything by mouth to an unconscious person. Seek medical attention.

TOXICOLOGICAL INFORMATION/NOTE TO THE PHYSICIAN

Symptoms of poisoning: No cases of poisoning in man have been reported.

Antidote: no specific antidote known.

In case of Emergency, call toll free number: 00800720021 / 0800730030 (24hrs).

TREATMENT

Symptomatic treatment prescribed. Give supportive therapy.
If ingested, lavage stomach and undertake alkaline catharsis.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control

Products act to use or store a pest control product under unsafe conditions.

WARRANTY

Seller's guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.



Shelf Life: 2 years from the date of manufacture if stored in unopened original container in a dry cool place.

Maisha rafuni: Miaka miwili baada ya tarehe ya kutengenezwa iwapo dawa hii itahifadhiwa kwa chombo chake asili bila kufunguliwa na iwe mahali pakavu na pasipo na joto.

Manufactured by / Imetengenezwa na:

Adama-Agan Ltd. P. O. Box 262 Ashdod 77102
Northern Industrial Zone, Israel
Web: www.adama.com

Approved by / Imethibitishwa na:

Tropical Pesticides Research Institute (TPRI) P.O. Box 3024,
ARUSHA – TANZANIA
TEL: 255 27 2508813/15, 255 27 2505871 Fax: 255 27 2505871

Distributed By / Inasambazwa Na:

TriaChem (T) Ltd.
P.O.Box 2750, Arusha-Tanzania
PLOT NO 49AA, Esso Road,
Tel/Fax: +255272548774
Web: www.triachem.com



Registrant: Farmchem Limited, Nairobi

Expiry date
Date of Manufacture SEE PACKING
Batch Number

T4 945A/1

ADAMA