



ADAMA

FOLPAN® 80 WDG

A multi-site, mode-of-action fungicide that delivers exceptional protection against many of the most damaging diseases in apples, grapes and other fruit and vegetable crops.



GROUP M

Active Ingredient

80% Folpet
(water-dispersible granule)

Packaging

Case: 2 X 5 kg bag

Application Rates and Area Treated

Rate: 1.25–5 kg/ha
Area Treated: 1–4 ha/bag

Water Volume

Ground: 1000–3000 L/ha
(600–1900 US gal/ha)

Aerial: Do not apply by air.

Rainfastness:

Do not apply when heavy rain is forecast.

MAKE A FOLPAN PLAN!

REGISTERED CROPS

- Apples
- Azalea
- Carnation
- Chrysanthemum
- Crabapples
- Field cucumbers
- Field tomatoes
- Grapes
- Iris
- Marigold
- Melons
- Poinsettia (greenhouse)
- Pumpkins
- Snapdragon
- Squash
- Strawberries
- Zinnia

KEY DISEASES CONTROLLED

- Alternaria leaf spot
- Anthracnose
- Black rot
- Brooks spot
- Dead arm
- Didymellina leaf spot
- Downy mildew
- Fly speck
- Fruit rot
- Grey mould
- Leaf spot
- Phythium root rot
- Powdery mildew
- Scab
- Septoria leaf spot
- Sooty blotch
- Stem rot

HOW IT WORKS

FOLPAN® 80 WDG is a multi-site-contact mode-of-action fungicide with protective ability to be used in a regularly scheduled maintenance program.



FOLPAN® 80 WDG

APPLICATION TIMING AND CROP STAGING

Crop	Diseases	Timing
Apples, Crabapples	Alternaria leaf spot, Black rot, Brooks spot, Fly speck, Scab, Sooty blotch	Maximum applications per season: 6 Apply every 7–14 days in a regular protective schedule from green tip up until harvest. See label for additional precautions.
Cranberries	Fruit rot	Apply when 5% of blossoms are open. Repeat 10–14 days later when 50–75% of blossoms are open.
Grapes	Phomopsis	Apply when new shoots are 1–2 inches long and repeat when growth is 4–6 inches long.
	Black rot, Downy mildew	Apply just before bloom, just after bloom and in first cover spray. For downy mildew, an additional application 2–3 weeks later may be needed.
	Powdery mildew	Apply in a tank mix with Kumulus® DF at first sign of mildew and repeat after 10 days.
Field cucumbers, Melons, Pumpkins, Squash	Anthrachnose, Downy mildew	Apply when first true leaves appear. Repeat at 7-day intervals until crop is harvested.
Field tomatoes	Anthrachnose	Apply during first bloom and repeat at 7-day intervals until harvest.
Strawberries	Grey mould, Fruit rot, Leaf spot	Apply before first infection and repeat at 7-day intervals to protect crop until harvest.
Carnation, Poinsettia, Azalea, Marigold, Zinnia, Chrysanthemum, Iris, Snapdragon	Blight, Phythium root rot, Stem rot, Alternaria leaf spot, Septoria leaf spot, Didymellina leaf spot, Anthracnose (depending on ornamental)	Generally, apply when ornamental emerges and repeat at regular intervals. Consult label for timing on specific disease and ornamental.

PRE-HARVEST INTERVALS

- Apples: 1 day
 - » Re-entry intervals:
 - » Hand-thinning: 6 days
 - » All other activities: 12 hours
- Field cucumbers, Field tomatoes, Grapes, Melons, Pumpkins, Squash, Strawberries: 1 day
- Cranberries: 30 days

NOTE: There are no restrictions on crop rotations, grazing or storage.

REGISTERED AND SUPPORTED TANK MIXES

- Compatible with most commonly used insecticides and fungicides. Check compatibilities before making new tank mixes.

STORAGE

May be stored at any temperature

MIXING INSTRUCTIONS

1. Fill spray tank nearly full and pour recommended amount of FOLPAN® 80 WDG on surface of water.
2. Fungicide can be premixed in a bucket ½ filled with water. Mix can be poured through screen into nearly filled spray tank.
3. Finish filling tank.
4. Keep agitator running during filling and spraying.



Always read and follow registered product label instructions. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.
1.855.264.6262 ADAMA.COM

®/™ FOLPAN is a registered trademark of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies.
© 2022 ADAMA Agricultural Solutions Canada Ltd.