



# THRASHER®

Provides excellent dual modes of action for control of 26 tough broadleaf weeds (including Group 2 resistant kochia) in wheat and barley.



GROUPS 4 & 6

## Active Ingredient

Bromoxynil Octanoic Ester  
225 g/L, and 2,4-D EH Ester  
225 g/L = EC

## Packaging

Case: 2 x 10 L jugs

## Application Rates & Acres Treated

Rate: 500 ml/ac  
Area Treated: 20 ac/jug

## Water Volume

Ground: 20 – 40 L/ac  
(5 – 10 US gal/ac)  
Aerial: 12 – 16 L/ac  
(3 – 4 US gal/ac)

## Rainfastness

1 hour

## REGISTERED CROPS

- Barley
- Wheat (spring)

## WEEDS CONTROLLED

- American nightshade
- Ball mustard
- Bluebur
- Cocklebur
- Common buckwheat<sup>1</sup>
- Common ground<sup>1</sup>
- Common ragweed
- Cow cockle
- Flixweed
- Green smartweed
- Kochia (2" high or 1- to 12- leaf)
- Lady's thumb
- Lamb's quarters<sup>1</sup>
- Night-flowering catchfly
- Pale smartweed
- Redroot pigweed<sup>2</sup>
- Russian thistle (2- to 12-leaf)
- Shepherd's purse
- Stinkweed<sup>1</sup>
- Tartary buckwheat<sup>1</sup>
- Velvetleaf (3" high)
- Volunteer canola
- Volunteer sunflower
- Wild buckwheat<sup>1</sup>
- Wild mustard<sup>1</sup>

<sup>1</sup> 8-leaf

<sup>2</sup> including triazine-resistant pigweed

## HOW IT WORKS

A combination of systemic and contact activity with weeds yellowing within 2 – 4 days and exhibiting abnormal growth (twisting and cupping of leaves) in 2 – 10 days.



# THRASHER<sup>®</sup>

## CROP STAGING

4-leaf to early flag leaf.

## REGISTERED AND SUPPORTED TANK MIXES

BISON<sup>®</sup> 400 L

## MIXING INSTRUCTIONS

1. Fill the spray tank  $\frac{1}{2}$  full with water.
2. Add the required amount of THRASHER<sup>®</sup> and agitate thoroughly.
3. Fill the tank with remaining water and agitate again before use.

## CROP ROTATIONS

No restrictions the year after application.

## PRE-HARVEST INTERVAL

30 days

## GRAZING RESTRICTIONS

- Do not permit livestock to graze fields within 30 days of application.
- Do not harvest, forage or cut for hay within 30 days of application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.

## STORAGE

- May be stored at any temperature
- Shake well before using.

### QUICK TIPS:

For best results when there is a heavy crop canopy, or when the majority of the weeds are cow cockle, smartweed or pigweed, use higher water volumes. Spray when weeds are in the seedling stage. Apply in good growing conditions. Application must be made before the crop canopy shields the weeds.



Always read and follow label directions.  
1.855.264.6262 ADAMA.COM

THRASHER<sup>®</sup> and BISON<sup>®</sup> are registered trademarks of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies.  
© 2022 ADAMA Agricultural Solutions Canada Ltd.