



BROMOTRIL®

Tough broadleaf weed control with tank-mix flexibility and excellent crop safety.



Active Ingredient

Bromoxynil Octanoate Ester
235 g/L = EC

Packaging

Case: 2 x 9.7 L jugs

Application Rates & Acres Treated

PRE-PLANT:

Rate: 490 ml/ac
Area Treated: 20 ac/jug

IN-CROP BROADLEAF:

Rate: 490 – 570 ml/ac
Area Treated: 17 – 20 ac/jug

Water Volume

Ground: 20 – 80 L/ac
(5 – 20 US gal/ac)

Aerial:

8 – 16 L/ac (3 – 5 US gal/ac)
Wheat and barley only
Do not apply pre-plant

Rainfastness

30 minutes

REGISTERED CROPS

| Crop | Stage |
|---|--|
| Barley, Oats, Wheat | Pre-seed burn-off with glyphosate |
| Crop | Crop Leaf Stage |
| Alfalfa (seedling) | 2 – 6 trifoliate |
| Alfalfa (established) | Spring: before the crop begins to shield the weeds |
| Barley, Oats, Triticale, Wheat (spring, durum) | 2-leaf to early flag |
| Winter wheat | Fall: 2- to 4-leaf Spring: first growth to early flag |
| Corn (field, sweet) | Beyond 4-leaf |
| Canary seed (seed production only) | 3- to 5-leaf |
| Fall rye | Spring: from first growth to early flag |
| Flax | 2 – 4 inches in height |
| Forage millet, Sorghum | 4-leaf to 8 inches |
| Seedling grasses (seed production only): Bromegrass, Fescue (creeping red, meadow), Orchard grass, Reed canarygrass, Russian wildrye, Timothy, Wheatgrass (crested, intermediate, slender, tall) | 2- to 4-leaf (year of establishment only) |

WEEDS CONTROLLED

Seedling up to 4-leaf stage:

- American nightshade
- Bluebur
- Cocklebur
- Common ragweed
- Cow cockle¹
- Green smartweed
- Kochia²
- Lady's thumb
- Pale smartweed
- Pigweed¹
- Russian thistle²
- Stinkweed¹
- Velvetleaf³
- Wild mustard¹

Seedling up to 8-leaf stage:

- Common buckwheat
- Common groundsel
- Lamb's quarters
- Tartary buckwheat
- Wild buckwheat

¹ In normal conditions, it will be controlled up to 4-leaf stage. Plants beyond this stage are unlikely to be controlled; the higher rate generally gives better results.

² Spray before plants are 2 inches high.

³ Spray before plants are 3 inches high.



BROMOTRIL®

HOW IT WORKS

BROMOTRIL® is a contact herbicide which controls Group 2- and Group 9-resistant biotypes. Leaves will yellow in 2–4 days with complete control in 7–14 days.

REGISTERED AND SUPPORTED POST-EMERGENT TANK MIXES

Post-emergent Herbicides:

- Spring wheat: 2,4-D ESTER 700, ~~LADDER ALL-IN®~~, MCPA ESTER 600, BISON® 400 L, ~~Traxos®~~
- Winter wheat: 2,4-D ESTER 700, MCPA ESTER 600, BISON® 400 L, ~~Traxos®~~
- Barley: 2,4-D ESTER 700, MCPA ESTER 600, BISON® 400 L
- Oats: MCPA ESTER 600
- Corn: Accent®, Atrazine, Banvel®(Dicamba), Ultim®
- Fall rye: MCPA ESTER 600
- ~~Flax: MCPA ESTER 600 or MCPA-K~~
- ~~Canary seed: MCPA ESTER 600~~
- ~~Seedling grasses: MCPA ESTER 600~~

Pre-seed:

- Glyphosate

Fungicides:

- BUMPER® 432 EC

MIXING INSTRUCTIONS

1. Fill spray tank ½ full with water.
2. Add required amount of BROMOTRIL®. Begin agitation.
3. If tank mixing, add the recommended amount of the tank-mix partner product to the spray tank first, agitate and then add BROMOTRIL® (unless otherwise directed by the BROMOTRIL® and tank-mix partner label).
4. Add the remaining amount of water while agitation continues.

CROP ROTATIONS

~~If applying pre-plant, only apply prior to wheat, barley and oats.~~

STORAGE

Do not freeze.

PRE-HARVEST INTERVAL

30 days

GRAZING RESTRICTIONS

- Do not use treated crops for grazing of livestock or green feed until 30 days after application.
- Do not cut treated crops for forage until 30 days after application.

QUICK TIPS:

Avoid spraying if temperatures are above 25° C. Leaf scorching may occur in corn and flax if applied during or after adverse growing conditions, such as cool and wet or hot (above 27° C) and humid weather. For best results, spray when weeds are in the seedling stage.



ADAMA

Always read and follow registered product label instructions. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.
1.855.264.6262 ADAMA.COM

BISON®, BROMOTRIL®, BUMPER®, ~~LADDER ALL-IN®~~ are registered trademarks of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies.

© 2022 ADAMA Agricultural Solutions Canada Ltd.