



ADAMA

# BROMOTRIL®

Tough broadleaf weed control with tank-mix flexibility and excellent crop safety.



GROUP 1

## Active Ingredient

Bromoxynil Octanoate Ester  
235 g/L = EC

## Packaging

Case: 2 x 9.7 L jugs

## Application Rates & Acres Treated

### PRE-PLANT:

Rate: 490 ml/ac  
Area Treated: 20 ac/jug

### IN-CROP BROADLEAF:

Rate: 490 – 570 ml/ac  
Area Treated: 17 – 20 ac/jug

## Water Volume

Ground: 20 – 80 L/ac  
(5 – 20 US gal/ac)

### Aerial:

8 – 16 L/ac (3 – 5 US gal/ac)  
Wheat and barley only  
Do not apply pre-plant

## Rainfastness

30 minutes

## REGISTERED CROPS

Crop	Stage
Barley, Oats, Wheat	Pre-seed burn-off with glyphosate
Crop	Crop Leaf Stage
Alfalfa (seedling)	2 – 6 trifoliate
Alfalfa (established)	Spring: before the crop begins to shield the weeds
Barley, Oats, Triticale, Wheat (spring, durum)	2-leaf to early flag
Winter wheat	Fall: 2- to 4-leaf Spring: first growth to early flag
Corn (field, sweet)	Beyond 4-leaf
Canary seed (seed production only)	3- to 5-leaf
Fall rye	Spring: from first growth to early flag
Flax	2 – 4 inches in height
Forage millet, Sorghum	4-leaf to 8 inches
<b>Seedling grasses (seed production only):</b> Bromegrass, Fescue (creeping red, meadow), Orchard grass, Reed canarygrass, Russian wildrye, Timothy, Wheatgrass (crested, intermediate, slender, tall)	2- to 4-leaf (year of establishment only)

## WEEDS CONTROLLED

### Seedling up to 4-leaf stage:

- American nightshade
- Bluebur
- Cocklebur
- Common ragweed
- Cow cockle<sup>1</sup>
- Green smartweed
- Kochia<sup>2</sup>
- Lady's thumb
- Pale smartweed
- Pigweed<sup>1</sup>
- Russian thistle<sup>2</sup>
- Stinkweed<sup>1</sup>
- Velvetleaf<sup>3</sup>
- Wild mustard<sup>1</sup>

### Seedling up to 8-leaf stage:

- Common buckwheat
- Common groundsel
- Lamb's quarters
- Tartary buckwheat
- Wild buckwheat

<sup>1</sup> In normal conditions, it will be controlled up to 4-leaf stage. Plants beyond this stage are unlikely to be controlled; the higher rate generally gives better results.

<sup>2</sup> Spray before plants are 2 inches high.

<sup>3</sup> Spray before plants are 3 inches high.



# BROMOTRIL®

## HOW IT WORKS

BROMOTRIL® is a contact herbicide which controls Group 2- and Group 9-resistant biotypes. Leaves will yellow in 2–4 days with complete control in 7–14 days.

## REGISTERED AND SUPPORTED POST-EMERGENT TANK MIXES

### Post-emergent Herbicides:

- Spring wheat: 2,4-D ESTER 700, LADDER ALL IN®, MCPA ESTER 600, BISON® 400 L, Traxos®
- Winter wheat: 2,4-D ESTER 700, MCPA ESTER 600, BISON® 400 L, Traxos®
- Barley: 2,4-D ESTER 700, MCPA ESTER 600, BISON® 400 L
- Oats: MCPA ESTER 600
- Corn: Accent®, Atrazine, Banvel®(Dicamba), Ultim®
- Fall rye: MCPA ESTER 600
- Flax: MCPA ESTER 600 or MCPA-K
- Canary seed: MCPA ESTER 600
- Seedling grasses: MCPA ESTER 600

### Pre-seed:

- Glyphosate

### Fungicides:

- BUMPER® 432 EC

## MIXING INSTRUCTIONS

1. Fill spray tank ½ full with water.
2. Add required amount of BROMOTRIL®. Begin agitation.
3. If tank mixing, add the recommended amount of the tank-mix partner product to the spray tank first, agitate and then add BROMOTRIL® (unless otherwise directed by the BROMOTRIL® and tank-mix partner label).
4. Add the remaining amount of water while agitation continues.

## CROP ROTATIONS

If applying pre-plant, only apply prior to wheat, barley and oats.

## STORAGE

Do not freeze.

## PRE-HARVEST INTERVAL

30 days

## GRAZING RESTRICTIONS

- Do not use treated crops for grazing of livestock or green feed until 30 days after application.
- Do not cut treated crops for forage until 30 days after application.

### QUICK TIPS:

Avoid spraying if temperatures are above 25° C. Leaf scorching may occur in corn and flax if applied during or after adverse growing conditions, such as cool and wet or hot (above 27° C) and humid weather. For best results, spray when weeds are in the seedling stage.



ADAMA

Always read and follow registered product label instructions. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.  
1.855.264.6262 ADAMA.COM

BISON®, BROMOTRIL®, BUMPER®, LADDER ALL IN® are registered trademarks of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies.

© 2022 ADAMA Agricultural Solutions Canada Ltd.