



ADAMA

ARROW® 240 EC

Get broad-spectrum grassy weed control
in soybeans, potatoes, beans and
other broadleaf and specialty crops.



GROUP 1

Active Ingredient

Clethodim 240 g/L = EC

Packaging

One case includes:

- 1 x 3 L jug of ARROW® 240 EC
- 1 x 9 L jug of X-ACT® adjuvant

Application Rates & Acres Treated

Rate: 50 – 150 ml/ac

Area Treated:

20 – 60 ac/case

Water Volume

Ground: 20 – 90 L/ac
(5 – 24 US gal/ac)

Aerial: Do not apply by air.

Rainfastness

1 hour

REGISTERED CROPS

- Alfalfa, seedling
- Beans
- Canola
- Coriander
- Cranberries
- Fenugreek
- Field peas
- Highbush blueberries
- Onions
- Potatoes
- Prairie carnations
- Soybeans
- Spinach
- Sunflowers

WEEDS CONTROLLED

Grass Species	Leaf Stage	Application Rates
Foxtail (green, yellow), Wild oats, Volunteer cereals (wheat, barley, oats)	2 – 4	50 ml/ac
Barnyard grass, Fall panicum, Proso millet, Volunteer corn, Volunteer canarygrass, Witch grass	2 – 6	50 ml/ac
Barnyard grass, Crabgrass (smooth, large), Fall panicum, Foxtail (green, yellow), Persian darnel, Proso millet, Volunteer canarygrass, Volunteer cereals (wheat, barley, oats), Volunteer corn, Wild oats, Witch grass Quackgrass (suppression)	2 – 6	75 ml/ac
Quackgrass (control) ¹	2 – 6	150 ml/ac

¹ Use a higher rate (1% v/v) of X-ACT® adjuvant for improved quackgrass control.



ARROW[®] 240 EC

HOW IT WORKS

The active ingredient is translocated from the treated foliage to the growing points of the leaves, shoots and roots. Leaf foliage will first change from green to yellowish, then purplish and finally brown. Newest leaf of affected plant pulls out easily in 3–5 days. Time required for complete control is normally 7–21 days following treatment, depending on growing conditions and crop competition.

CROP STAGING

- Most crops are tolerant at all stages, so target applications at the optimal weed stage.
- Always adhere to the pre-harvest interval for each crop.

REGISTERED AND SUPPORTED TANK MIXES

- Canola: Lontrel™ XC or Muster®
- Clearfield® canola only: PHANTOM® 240 SL
- LibertyLink® canola only: glufosinate
- Field peas: PHANTOM® 240 SL
- Glyphosate-tolerant soybeans: glyphosate

MIXING INSTRUCTIONS

1. Fill clean spray tank ½ full with water. Start agitation.
2. Add the correct amount of ARROW® 240 EC. Continue to agitate.
3. Add the correct amount of adjuvant X-ACT® along with the remaining amount of water necessary to fill the spray tank.
4. Continue to agitate or run the bypass system.
5. After any break in the spraying operation, agitate thoroughly before spraying again.
6. Do not allow the mixture to sit overnight.
7. If tank-mixing, follow label directions for each tank-mix partner.

ADJUVANT RATE

- X-ACT®* @ 0.5–1.0% v/v
- Methylated seed oil (MSO) @ 0.5% v/v
- Non-ionic surfactant (NIS) @ 0.25% v/v

*Use a higher rate (1% v/v) of X-ACT® for improved quackgrass control.

GRAZING RESTRICTIONS

Do not cut treated crops for feed or graze until 60 days after application.

STORAGE

- May be stored at any temperature.
- Shake well before use.

CROP ROTATIONS

No restrictions when applied alone

PRE-HARVEST INTERVALS

- Blueberries, Spinach: 14 days
- Alfalfa, Fenugreek: 30 days
- Onions: 45 days
- Canola, Coriander, Beans, Lentils, Potatoes, Chickpeas: 60 days
- Sunflowers: 72 days
- Soybeans, Field peas: 75 days

QUICK TIPS:

ARROW® 240 EC works best when applied to actively growing weeds. Regrowth of tillers may occur if applied to weeds under stress conditions. ARROW® 240 EC must be used with the adjuvant X-ACT®.



ADAMA

Always read and follow registered product label instructions. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.
1.855.264.6262 ADAMA.COM

© ARROW, BADGE, PHANTOM and X-ACT are registered trademarks of ADAMA Agricultural Solutions Canada Ltd. All other products mentioned are trademarks of their respective companies.
© 2023 ADAMA Agricultural Solutions Canada Ltd.