



BADGE®

Get proven control of almost 30 broadleaf weeds in cereals, flax and corn with easy-to-use tank-mix options for one-shot weed control.



GROUPS 4 & 6

Active Ingredient

Bromoxynil 225 g/L and MCPA 2 EH ester 225 g/L = EC

Packaging

Case: 2 x 10 L jugs

Application Rates & Acres Treated

Rate: 500 ml/ac

Area Treated: 20 ac/jug

Water Volume

Ground:

- Cereals: 20–40 L/ac (5–10 US gal/ac)
- Corn: 80–120 L/ac (20–30 US gal/ac)
- Forages: 60 L/ac (15 US gal/ac)

Aerial:

8–20 L/ac (2–5 US gal/ac)

Rainfastness

1 hour

REGISTERED CROPS

Field crops:

- Barley
- Corn
- Fall rye
- Oats
- Wheat (spring, winter)

WEEDS CONTROLLED

- American nightshade
- Ball mustard
- Bluebur
- Canada thistle¹
- Cocklebur
- Common buckwheat
- Common groundsel
- Common ragweed
- Cow cockle²
- Flixweed
- Green smartweed
- Kochia³
- Lady's thumb
- Lamb's quarters
- Night-flowering catchfly
- Pale smartweed
- Perennial sow thistle¹
- Redroot pigweed
- Russian thistle³
- Scentless chamomile⁴
- Shepherd's purse
- Stinkweed
- Tartary buckwheat
- Velvetleaf⁵
- Volunteer canola (all types)
- Volunteer sunflower
- Wild buckwheat
- Wild mustard
- Wild tomato
- Wormseed mustard

¹Top growth control

²Up to 4-leaf stage

³When sprayed before plants are 2 inches high

⁴Spring annual only

⁵When sprayed before plants are 3 inches high



BADGE®

HOW IT WORKS

A combination of systemic and contact activity with weeds yellowing within 2–4 days and exhibiting abnormal growth (twisting and cupping of leaves) in 2–10 days.

CROP STAGING

| Crop | Timing |
|---|--|
| Barley, Oats, Spring wheat | 2-leaf to early flag |
| Corn | 4- to 6-leaf |
| Fall rye | When growth commences in spring to early flag leaf |
| Timothy (established for seed production) | Prior to shot blade in the seed production year |
| Winter wheat (includes underseeded with red clover) | 2- to 4-leaf stage in the fall or after growth resumes up to early flag leaf |

REGISTERED AND SUPPORTED TANK MIXES

- Corn: atrazine
- Oats: MCPA ESTER 600
- Spring wheat and barley: Avenge®, MCPA ESTER 600, Refine Extra®, BISON® 400 L
- Winter wheat: MCPA ESTER 600, Refine Extra®

MIXING INSTRUCTIONS

1. Fill clean spray tank ½ full with water.
2. Add the required amount of BADGE® and agitate thoroughly.
3. Fill the tank and agitate again before use.
4. When tank-mixing, follow instructions on both labels.

CROP ROTATIONS

No re-cropping restrictions the year after treatment.

PRE-HARVEST INTERVAL

Flax: 60 days

STORAGE

Do not freeze.

GRAZING RESTRICTIONS

- Do not graze treated grain or established timothy crops or cut for feed within 30 days of application.
- Do not graze meadow foxtail in the year of treatment.
- Do not graze other treated forage grasses within 56 days of treatment.

QUICK TIPS:

BADGE® herbicide is well known for being gentle on the crop. Avoid spraying if temperatures are above 25°C.



ADAMA

Always read and follow label directions.
1.855.264.6262 ADAMA.COM

ARROW®, ARROW ALL IN™, BADGE® and BISON® are registered trademarks of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies.
© 2022 ADAMA Agricultural Solutions Canada Ltd.