



# BADGE®

Get proven control of almost 30 broadleaf weeds in cereals and corn with easy-to-use tank-mix options for one-shot weed control.



## GROUPS 4 & 6

### Active Ingredient

Bromoxynil 225 g/L and MCPA 2 EH ester 225 g/L = EC

### Packaging

Case: 2 x 10 L jugs

### Application Rates & Acres Treated

Rate: 500 ml/ac

Area Treated: 20 ac/jug

### Water Volume

#### Ground:

- Cereals: 20–40 L/ac (5–10 US gal/ac)
- Corn: 80–120 L/ac (20–30 US gal/ac)
- Forages: 60 L/ac (15 US gal/ac)

#### Aerial:

8–20 L/ac (2–5 US gal/ac)

### Rainfastness

1 hour

### REGISTERED CROPS

#### Field crops:

- Barley
- Corn
- Fall rye
- Oats
- Wheat (spring, winter)

#### WEEDS CONTROLLED

- American nightshade
- Ball mustard
- Bluebur
- Canada thistle<sup>1</sup>
- Cocklebur
- Common buckwheat
- Common groundsel
- Common ragweed
- Cow cockle<sup>2</sup>
- Flixweed
- Green smartweed
- Kochia<sup>3</sup>
- Lady's thumb
- Lamb's quarters
- Night-flowering catchfly
- Pale smartweed
- Perennial sow thistle<sup>1</sup>
- Redroot pigweed
- Russian thistle<sup>3</sup>
- Scentless chamomile<sup>4</sup>
- Shepherd's purse
- Stinkweed
- Tartary buckwheat
- Velvetleaf<sup>5</sup>
- Volunteer canola (all types)
- Volunteer sunflower
- Wild buckwheat
- Wild mustard
- Wild tomato
- Wormseed mustard

<sup>1</sup>Top growth control

<sup>2</sup>Up to 4-leaf stage

<sup>3</sup>When sprayed before plants are 2 inches high

<sup>4</sup>Spring annual only

<sup>5</sup>When sprayed before plants are 3 inches high



# BADGE®

## HOW IT WORKS

A combination of systemic and contact activity with weeds yellowing within 2–4 days and exhibiting abnormal growth (twisting and cupping of leaves) in 2–10 days.

## CROP STAGING

Crop	Timing
Barley, Oats, Spring wheat	2-leaf to early flag
Corn	4- to 6-leaf
Fall rye	When growth commences in spring to early flag leaf
Timothy (established for seed production)	Prior to shot blade in the seed production year
Winter wheat (includes underseeded with red clover)	2- to 4-leaf stage in the fall or after growth resumes up to early flag leaf

## REGISTERED AND SUPPORTED TANK MIXES

- Corn: atrazine
- Oats: MCPA ESTER 600
- Spring wheat and barley: Avenge®, MCPA ESTER 600, Refine Extra®, BISON® 400 L
- Winter wheat: MCPA ESTER 600, Refine Extra®

## MIXING INSTRUCTIONS

1. Fill clean spray tank ½ full with water.
2. Add the required amount of BADGE® and agitate thoroughly.
3. Fill the tank and agitate again before use.
4. If tank-mixing with a grassy weed herbicide, read both labels and follow the more stringent directions for tank-mixing.

## CROP ROTATIONS

No re-cropping restrictions the year after treatment.

## PRE-HARVEST INTERVAL

Flax: 60 days

## STORAGE

Do not freeze.

## GRAZING RESTRICTIONS

- Do not graze treated grain or established timothy crops or cut for feed within 30 days of application.
- Do not graze meadow foxtail in the year of treatment.
- Do not graze other treated forage grasses within 56 days of treatment.

### QUICK TIPS:

BADGE® herbicide is well known for being gentle on the crop. Avoid spraying if temperatures are above 25°C.



ADAMA

Always read and follow label directions.  
1.855.264.6262 ADAMA.COM

®/™BADGE and BISON are registered trademarks of ADAMA Agricultural Solutions Canada Ltd.  
All other products are trademarks of their respective companies.  
© 2023 ADAMA Agricultural Solutions Canada Ltd.