

2023-1399
2024-03-27

Container label

GROUP 2 HERBICIDE

DAVAI® 80 SL

Herbicide
Solution

FOR SALE FOR USE IN FIELD PEAS, DRY BEANS, IMIDAZOLINONE-TOLERANT
LENTILS AND SOYBEANS

COMMERCIAL

AGRICULTURAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: Imazamox (present as ammonium salt) 80 g/L

REGISTRATION NO. 32929 PEST CONTROL PRODUCTS ACT



POISON

CAUTION – EYE IRRITANT

NET CONTENTS: 1-1050 litres

ADAMA Agricultural Solutions Canada Ltd.

300 – 191 Lombard Avenue
Winnipeg, Manitoba
R3B 0X1
1-855-264-6262

For emergency medical help and health/safety inquiries call ProPharma Group at 1.877.250.9291
(24 hours a day)

For spill, leak or fire call INFOTRAC at 1.800.535.5053 (24 hours a day)

2023-1399
2024-03-27

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

Harmful if absorbed through skin or inhaled. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. May irritate eyes.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear goggles or face shield for mixing/loading, cleanup and repair activities. Gloves are not required during application within a closed cab.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

2023-1399
2024-03-27

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This product contains aromatic petroleum distillates which are toxic to aquatic organisms.

STORAGE

1. Store this product away from food or feed.
2. Store above 5° C in original, tightly closed container.
2. Do not ship or store near seed and fertilizers.
3. Store in cool, dry, locked, well-ventilated area without floor drain.
4. Keep from freezing.

DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2023-1399
2024-03-27

Pamphlet label

GROUP 2 HERBICIDE

DAVAI® 80 SL

Herbicide
Solution

FOR SALE FOR USE IN FIELD PEAS, DRY BEANS, IMIDAZOLINONE-TOLERANT
LENTILS AND SOYBEANS

COMMERCIAL

AGRICULTURAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: Imazamox (present as ammonium salt) 80 g/L

REGISTRATION NO.32929 PEST CONTROL PRODUCTS ACT



POISON

CAUTION – EYE IRRITANT

NET CONTENTS: 1-1050 litres

ADAMA Agricultural Solutions Canada Ltd.

300 – 191 Lombard Avenue
Winnipeg, Manitoba
R3B 0X1
1-855-264-6262

For emergency medical help and health/safety inquiries call ProPharma Group at 1.877.250.9291
(24 hours a day)

For spill, leak or fire call INFOTRAC at 1.800.535.5053 (24 hours a day)

| | |
|---|----|
| GENERAL INFORMATION..... | 5 |
| <i>SECTION 1: THE PRODUCT</i> | 5 |
| SAFETY AND HANDLING..... | 5 |
| <i>SECTION 2: PRECAUTIONS, PROTECTIVE CLOTHING AND EQUIPMENT, AND RE-ENTRY RESTRICTIONS</i> | 5 |
| <i>SECTION 3: FIRST AID AND TOXICOLOGICAL INFORMATION</i> | 5 |
| <i>SECTION 4: ENVIRONMENTAL PRECAUTIONS</i> | 6 |
| <i>SECTION 5: STORAGE</i> | 6 |
| <i>SECTION 6: DISPOSAL</i> | 6 |
| <i>SECTION 7: NOTICE TO USER</i> | 6 |
| DIRECTIONS FOR USE | 7 |
| <i>SECTION 8: PRODUCT OVERVIEW</i> | 7 |
| <i>SECTION 9: APPLICATION INSTRUCTIONS AND USE LIMITATIONS</i> | 7 |
| <i>SECTION 10: PRE-HARVEST, GRAZING AND FEEDING INTERVALS</i> | 7 |
| <i>SECTION 11: SPRAY BUFFER ZONES</i> | 8 |
| <i>SECTION 12: REGISTERED CROPS AND APPLICATION TIMING</i> | 8 |
| <i>SECTION 13: WEEDS CONTROLLED</i> | 11 |
| <i>SECTION 14: MIXING INSTRUCTIONS</i> | 14 |
| <i>SECTION 15: ROTATIONAL CROPS</i> | 14 |
| <i>SECTION 16: RESISTANCE MANAGEMENT RECOMMENDATIONS</i> | 15 |

2023-1399
2024-03-27

GENERAL INFORMATION

THE PRODUCT

DAVAI® 80 SL is a selective herbicide that can be applied as an early post-emergent treatment in field peas, soybeans, dry beans and imidazolinone-tolerant lentils for broad-spectrum weed control.

SAFETY AND HANDLING

PRECAUTIONS, PROTECTIVE CLOTHING AND EQUIPMENT, AND RE-ENTRY RESTRICTIONS

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.

Harmful if absorbed through skin or inhaled. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. May irritate eyes.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear goggles or face shield for mixing/loading, cleanup and repair activities. Gloves are not required during application within a closed cab.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

FIRST AID AND TOXICOLOGICAL INFORMATION

FIRST AID: Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

2023-1399
2024-03-27

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants. Observe spray buffer zones specified under **DIRECTIONS FOR USE**.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This product contains aromatic petroleum distillates which are toxic to aquatic organisms.

STORAGE

1. Store this product away from food or feed.
2. Store above 5° C in original, tightly closed container.
3. Do not ship or store near seed and fertilizers.
4. Store in cool, dry, locked, well-ventilated area without floor drain.
5. Keep from freezing.

DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an

2023-1399
2024-03-27

offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

DIRECTIONS FOR USE

PRODUCT OVERVIEW

DAVAI® 80 SL is a selective herbicide that can be applied as an early postemergence treatment in field peas, soybeans, dry beans, and imidazolinone-tolerant lentils for broad-spectrum weed control.

Cool weather conditions or drought will delay herbicidal activity and if prolonged, may result in poor weed control. Use of DAVAI® 80 SL in hot, humid weather may result in temporary leaf yellowing, leaf flecking, bronzing or burning. The crop usually outgrows this condition within 10 days.

When weeds are stressed due to drought, flooding, hot or prolonged cool temperatures (15°C or less), control can be reduced or delayed since weeds are not actively growing. Weeds escapes or regrowth may occur under prolonged stress conditions or low fertility. Do not make applications to weeds stressed longer than 20 days due to lack of moisture, as unsatisfactory control can result.

APPLICATION INSTRUCTIONS AND USE LIMITATIONS

- DO NOT APPLY BY AIR. Apply using ground equipment only.
- DO NOT apply more than once per year.
- DO NOT apply to crops that have been subjected to stress from conditions such as hail damage, flooding, drought, hot, humid weather, widely fluctuating temperature conditions, prolonged cold weather or injury from prior herbicide applications, as crop injury may result.
- DO NOT apply when weather conditions may cause spray drift from treated areas to adjacent crops.
- As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.
- DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.
- Apply only to agricultural crops when the potential for drift to areas of human habitation and areas of human activity (houses, cottages, schools and recreational areas) is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment and sprayer settings.

PRE-HARVEST, GRAZING AND FEEDING INTERVALS

- Grazing/Feeding: Do not graze the treated crops; Field peas may be fed to livestock 30 days after application. Do not graze the treated imidazolinone-tolerant Lentils within 20 days of application.

2023-1399
2024-03-27

- Pre-harvest Interval: Field peas can be harvested 60 days after treatment. Dry beans can be harvested 75 days after treatment. Soybeans can be harvested 85 days after treatment. Lentils can be harvested 60 days after treatment.

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

SPRAY BUFFER ZONES

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label,
- low-clearance hooded or shielded sprayers that prevent spray contact with crop, fruit or foliage.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands).

| Method of Application | Crop | Spray Buffer Zones (metres) Required for the Protection of Terrestrial Habitat |
|-----------------------|-----------|--|
| Field sprayer | All crops | 1 |

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

REGISTERED CROPS AND APPLICATION TIMING

REGISTERED CROPS:

Soybeans Eastern and Western Canada

DAVAI® 80 SL is a selective herbicide that can be applied as an early post-emergent treatment in soybeans for control of grass and broadleaf weeds.

| APPLICATION INSTRUCTIONS – DAVAI® 80 SL + a non-ionic surfactant (such as ADAMA Adjuvant 80), a methylated seed oil adjuvant (such as MSO Spray Adjuvant), Merge® Adjuvant or Hasten® NT Ultra | |
|---|--|
| Timing | Early post-emergence |
| Rate | DAVAI® 80 SL at 250 mL/ha + a non-ionic surfactant (such as |

2023-1399
2024-03-27

| | |
|----------------------|--|
| | ADAMA Adjuvant 80) at 0.25% v/v, a methylated seed oil adjuvant (such as MSO Spray Adjuvant) at 0.5% v/v, Merge® Adjuvant at 0.5% v/v or Hasten® NT Ultra at 0.5% v/v |
| Water Volume | 100 L/ha |
| Weeds Controlled | Provides control of broadleaf and grass weeds as listed in the WEEDS CONTROLLED section of this label. |
| Pre-harvest Interval | 85 days |
| Remark | Soybeans: Apply from Emergence to 3 expanded trifoliolate leaves after weeds have emerged. Weeds: Apply when broadleaf weeds are from the cotyledon to 4- leaf stage and when grassy weeds are at the 1 - 4 true leaf or early tillering. |

APPLICATION INSTRUCTIONS – DAVAI® 80 SL + a non-ionic surfactant (such as ADAMA Adjuvant 80), a methylated seed oil adjuvant (such as MSO Spray Adjuvant), Merge® Adjuvant or Hasten® NT Ultra.

| | |
|----------------------|--|
| Timing | Early post-emergence |
| Rate | DAVAI® 80 SL at 187.5 mL/ha + a non-ionic surfactant (such as ADAMA Adjuvant 80) at 0.25% v/v, a methylated seed oil adjuvant (such as MSO Spray Adjuvant) at 0.5% v/v, Merge® Adjuvant at 0.5% v/v or Hasten® NT Ultra at 0.5% v/v |
| Water Volume | 100 L/ha |
| Weeds Controlled | Volunteer Canola |
| Pre-harvest Interval | 85 days |
| Remark | Soybeans: Apply from Emergence to 3 expanded trifoliolate leaves after weeds have emerged. Weeds: Apply when Volunteer canola is from the cotyledon to 4- leaf stage |

Field Peas Eastern and Western Canada

DAVAI® 80 SL is a selective herbicide that can be applied as an early post-emergent treatment in field peas for control of broadleaf weeds.

APPLICATION INSTRUCTIONS – DAVAI® 80 SL + a non-ionic surfactant (such as ADAMA Adjuvant 80), a methylated seed oil adjuvant (such as MSO Spray Adjuvant), Merge® Adjuvant or Hasten® NT Ultra

| | |
|--------------|--|
| Timing | Early post-emergence |
| Rate | DAVAI® 80 SL at 250 mL/ha + a non-ionic surfactant (such as ADAMA Adjuvant 80) at 0.25% v/v, a methylated seed oil adjuvant (such as MSO Spray Adjuvant) at 0.5% v/v, Merge® Adjuvant at 0.5% v/v or Hasten® NT Ultra at 0.5% v/v |
| Water Volume | 100 L/ha |
| | Provides control of broadleaf as listed in the WEEDS CONTROLLED |

2023-1399
2024-03-27

| | |
|----------------------|--|
| Weeds Controlled | section of this label |
| Pre-harvest Interval | 60 days |
| Remark | Apply from the 1 - 6 true leaf stage of field peas after weeds have emerged. Weeds: Apply when broadleaf weeds are from the cotyledon to 4- leaf stage and when grassy weeds are at the 1 - 4 true leaf or early tillering. For field peas, initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield. |

APPLICATION INSTRUCTIONS – DAVAI® 80 SL + a non-ionic surfactant (such as ADAMA Adjuvant 80), a methylated seed oil adjuvant (such as MSO Spray Adjuvant), Merge® Adjuvant or Hasten® NT Ultra

| | |
|----------------------|--|
| Timing | Early post-emergence |
| Rate | DAVAI® 80 SL Herbicide at 187.5 mL/ha + a non-ionic surfactant (such as ADAMA Adjuvant 80) at 0.25% v/v, a methylated seed oil adjuvant (such as MSO Spray Adjuvant) at 0.5% v/v, Merge® Adjuvant at 0.5% v/v or Hasten® NT Ultra at 0.5% v/v |
| Water Volume | 100 L/ha |
| Weeds Controlled | Volunteer Canola |
| Pre-harvest Interval | 60 days |
| Remark | Field Peas: Apply from the 1 - 6 true leaf stage of field peas after weeds have emerged. Weeds: Apply when Volunteer canola is from the cotyledon to 4- leaf stage |

Dry Beans Eastern and Western Canada

DAVAI® 80 SL is a selective herbicide that can be applied as an early post-emergent treatment in dry beans for control of grass and broadleaf weeds.

APPLICATION INSTRUCTIONS – DAVAI® 80 SL + a non-ionic surfactant (such as ADAMA Adjuvant 80), a methylated seed oil adjuvant (such as MSO Spray Adjuvant), Merge® Adjuvant or Hasten® NT Ultra

| | |
|----------------------|--|
| Timing | Early post-emergence |
| Rate | DAVAI® 80 SL at 250 mL/ha + a non-ionic surfactant (such as ADAMA Adjuvant 80) at 0.25% v/v, a methylated seed oil adjuvant (such as MSO Spray Adjuvant) at 0.5% v/v, Merge® Adjuvant at 0.5% v/v or Hasten® NT Ultra at 0.5% v/v |
| Water Volume | 100 L/ha |
| Weeds Controlled | Provides control of broadleaf and grass weeds as listed in the WEEDS CONTROLLED section of this label. |
| Pre-harvest Interval | 75 days |

2023-1399
2024-03-27

| | |
|--------|---|
| Remark | Apply from Emergence to 3 expanded trifoliolate leaves after weeds have emerged. Weeds: Apply when broadleaf weeds are from the cotyledon to 4- leaf stage and when grassy weeds are at the 1 - 4 true leaf or early tillering. For dry beans tolerance may vary between varieties. Test new varieties on a small area for tolerance before widespread use. |
|--------|---|

| | |
|---|--|
| APPLICATION INSTRUCTIONS – DAVAI® 80 SL + a non-ionic surfactant (such as ADAMA Adjuvant 80), a methylated seed oil adjuvant (such as MSO Spray Adjuvant), Merge® Adjuvant or Hasten® NT Ultra | |
| Timing | Early post-emergence |
| Rate | DAVAI® 80 SL at 187.5 mL/ha + a non-ionic surfactant (such as ADAMA Adjuvant 80) at 0.25% v/v, a methylated seed oil adjuvant (such as MSO Spray Adjuvant) at 0.5% v/v, Merge® Adjuvant at 0.5% v/v or Hasten® NT Ultra at 0.5% v/v |
| Water Volume | 100 L/ha |
| Weeds Controlled | Volunteer Canola |
| Pre-harvest Interval | 75 days |
| Remark | Dry Beans: Apply from Emergence to 3 expanded trifoliolate leaves after weeds have emerged. Weeds: Apply when Volunteer canola is from the cotyledon to 4- leaf stage |

Dry bean varieties may vary in their tolerance to herbicides, including to DAVAI® 80 SL. Since not all dry bean varieties have been tested for tolerance to DAVAI® 80 SL, first use of this Herbicide should be limited to a small area of each variety to confirm tolerance prior to adoption as a general field practice. Additionally, consult your seed supplier for information on the tolerance of specific varieties of dry common beans to DAVAI® 80 SL.

Imidazolinone-Tolerant Lentils – Only for use in the Prairie Provinces and interior of British Columbia (including the Peace River region) only

DAVAI® 80 SL is a selective herbicide that can be applied as an early post-emergent treatment in imidazolinone-tolerant lentils for control of grass and broadleaf weeds.

| | |
|--|---|
| APPLICATION INSTRUCTIONS – DAVAI® 80 SL + a non-ionic surfactant (such as ADAMA Adjuvant 80), a methylated seed oil adjuvant (such as MSO Spray Adjuvant), Merge® Adjuvant or Hasten® NT Ultra: | |
| Timing | 1–9 node stage, post-emergence |
| Rate | DAVAI® 80 SL at 250 mL/ha + a non-ionic surfactant (such as ADAMA Adjuvant 80) at 0.25% v/v, a methylated seed oil adjuvant (such as MSO Spray Adjuvant) at 0.5% v/v, Merge® Adjuvant at |

2023-1399
2024-03-27

| | |
|----------------------|--|
| | 0.5% v/v or Hasten® NT Ultra at 0.5% v/v |
| Water Volume | 100 L/ha |
| Weeds Controlled | Provides control of broadleaf and grass weeds as listed in the WEEDS CONTROLLED section of this label. |
| Pre-harvest Interval | 60 days |
| Remark | Apply from the 1-9 node stage and after weeds have emerged. Weeds: Apply when broadleaf weeds are at the cotyledon to the 4-leaf stage. Grassy weeds can be treated up to tillering provided they are not beyond the 4 true leaf stage. |

Tank Mixes

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

DAVAI® 80 SL + Phantom® 240 SL - Tank Mixture

| | |
|---|---|
| APPLICATION INSTRUCTIONS – DAVAI® 80 SL + Phantom® 240 SL + a non-ionic surfactant (such as ADAMA Adjuvant 80), a methylated seed oil adjuvant (such as MSO Spray Adjuvant), Merge Adjuvant or Hasten NT Ultra | |
| Timing | Early post-emergence |
| Rate | DAVAI® 80 SL at 200 mL/ha + Phantom® 240 SL at 65 mL/ha + a non-ionic surfactant (such as ADAMA Adjuvant 80) at 0.25% v/v, a methylated seed oil adjuvant (such as MSO Spray Adjuvant) at 0.5% v/v, Merge Adjuvant at 0.5% v/v or Hasten® NT Ultra at 0.5% v/v |
| Water Volume | 100 L/ha |
| Weeds Controlled | Provides control of broadleaf and grass weeds as listed in the WEEDS CONTROLLED section of this label. |
| Pre-harvest Interval | 60 days for field peas; 75 days for dry beans; 85 days for soybeans |
| | Field Peas: Apply from the 1 - 6 true leaf stage of field peas after weeds have emerged. Soybeans: Apply from Emergence to 3 expanded trifoliolate leaves after weeds have emerged. Dry Beans: Apply from Emergence to the second trifoliolate leaf after weeds have emerged. |

2023-1399
2024-03-27

| | |
|--|---|
| | <p>Imidazolinone-Tolerant Lentils¹: Apply from the 1 – 6 node stage after weeds have emerged.</p> <p>Weeds: Apply when broadleaf weeds are from the cotyledon to 4- leaf stage and when grassy weeds are at the 1 - 4 true leaf or early tillering.</p> <p>For field peas, initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield.</p> <p>For dry beans tolerance may vary between varieties. Test new varieties on a small area for tolerance before widespread use.</p> |
|--|---|

¹ For use in the Prairie Provinces and interior of British Columbia (including the Peace River region) only

DAVAI® 80 SL + Basagran® Forte + UAN 28%

| APPLICATION INSTRUCTIONS – DAVAI® 80 SL + Basagran® Forte + UAN 28% | |
|--|---|
| Timing | Early post-emergence |
| Rate | DAVAI® 80 SL at 250 mL/ha + Basagran® Forte at 1.25L/ha + UAN 28% at 2 L/ha |
| Water Volume | 100 L/ha |
| Weeds Controlled | Provides control of broadleaf and grass weeds as listed in the WEEDS CONTROLLED section of this label. |
| Pre-harvest Interval | 60 days for field peas; 75 days for dry beans; 85 days for soybeans |
| Remark | <p>Treated crops are not to be grazed or cut for hay.</p> <p>Field Peas: Apply from the 3 - 6 true leaf stage of field peas after weeds have emerged.</p> <p>Soybeans: Apply from unifoliate to 3 expanded trifoliate leaves after weeds have emerged.</p> <p>Dry Beans: Apply from first to the second trifoliate leaf after weeds have emerged.</p> <p>Weeds: Apply when broadleaf weeds are from the cotyledon to 4- leaf stage and when grassy weeds are at the 1 - 4 true leaf or early tillering.</p> <p>For field peas, initial transient crop yellowing may be observed after application, but this is outgrown and should not affect yield.</p> <p>For dry beans tolerance may vary between varieties. Test new varieties on a small area for tolerance before widespread use.</p> |

2023-1399
2024-03-27

WEEDS CONTROLLED

APPLICATION INSTRUCTIONS – DAVAI® 80 SL used as directed will control:

DAVAI® 80 SL Alone

DAVAI® 80 SL + a non-ionic surfactant (such as ADAMA Adjuvant 80), a methylated seed oil adjuvant (such as MSO Spray Adjuvant), Merge® Adjuvant or Hasten® NT Ultra used as directed will control:

| | | |
|--|--|--------------------------|
| DAVAI® 80 SL at 250 mL/ha + a non-ionic surfactant (such as ADAMA Adjuvant 80) at 0.25% v/v, a methylated seed oil adjuvant (such as MSO Spray Adjuvant) at 0.5% v/v, Merge® Adjuvant at 0.5% v/v or Hasten® NT Ultra at 0.5% v/v | | |
| WEEDS CONTROLLED | | |
| GRASS WEEDS | RECOMMENDED STAGE | |
| Barnyard grass | 1-4 true leaf stage or early tillering | |
| Green foxtail (including Group 1 resistant) ¹ | | |
| Persian Darnel | | |
| Tame oat | | |
| Volunteer barley | | |
| Volunteer canaryseed | | |
| Volunteer wheat (non imidazolinone-tolerant varieties) | | |
| Wild oat (including Group 1 resistant) ¹ | | |
| Yellow foxtail | | |
| BROADLEAF WEEDS | | RECOMMENDED STAGE |
| Cow Cockle | Cotyledon to 4 leaf stage | |
| Flixweed | | |
| Green smartweed | | |
| Lamb’s quarters | | |
| Redroot Pigweed | | |
| Shepherd’s Purse | | |
| Stinkweed | | |
| Stork’s bill | | |
| Volunteer Canola (non imidazolinone-tolerant varieties) | | |
| Wild mustard | | |
| WEEDS SUPPRESSED | | |
| Cleavers | | |
| Japanese brome | | |
| Wild buckwheat | | |

¹ DAVAI® 80 SL will not control weed biotypes that are multiple-resistant to both Group 1 and Group 2 Herbicides

DAVAI® 80 SL + Phantom® 240 SL - Tank Mixture

2023-1399
2024-03-27

DAVAI® 80 SL + Phantom® 240 SL + a non-ionic surfactant (such as ADAMA Adjuvant 80), a methylated seed oil adjuvant (such as MSO Spray Adjuvant), Merge® Adjuvant or Hasten® NT Ultra used as directed will control:

| | |
|--|--|
| DAVAI® 80 SL Herbicide at 200 mL/ha + Phantom® 240 SL at 65 mL/ha + a non-ionic surfactant (such as ADAMA Adjuvant 80) at 0.25% v/v, a methylated seed oil adjuvant (such as MSO Spray Adjuvant) at 0.5% v/v, Merge® Adjuvant at 0.5% v/v or Hasten® NT Ultra at 0.5% v/v | |
| WEEDS CONTROLLED | RECOMMENDED STAGE |
| GRASS WEEDS | 1-4 true leaf stage or early tillering |
| Green foxtail (including Group 1 resistant) ¹ | |
| Tame oat | |
| Wild oat (including Group 1 resistant) ¹ | |
| Yellow foxtail | |
| BROADLEAF WEEDS | RECOMMENDED STAGE |
| Chickweed | Cotyledon to 4 leaf stage |
| Green smartweed | |
| Lamb's quarters | |
| Redroot pigweed | |
| Stinkweed | |
| Wild mustard | |
| WEEDS SUPPRESSED | |
| Barnyard grass | |
| Volunteer barley | |
| Volunteer canola (non imidazolinone-tolerant) | |
| Wild buckwheat | |

¹ DAVAI® 80 SL will not control weed biotypes that are multiple-resistant to both Group 1 and Group 2 Herbicides.

DAVAI® 80 SL + Basagran® Forte + UAN 28% - Tank Mixture

DAVAI® 80 SL tank-mixed with Basagran® Forte and UAN 28% as directed will control:

| | |
|---|--|
| DAVAI® 80 SL at 250 mL/ha + Basagran® Forte at 1.25 L/ha + UAN 28% at 2 L/ha | |
| WEEDS CONTROLLED | |
| GRASS WEEDS | RECOMMENDED STAGE |
| Barnyard grass | 1-4 true leaf stage or early tillering |
| Green foxtail (including Group 1 resistant) ¹ | |
| Persian darnel | |
| Tame oat | |
| Volunteer barley | |
| Volunteer canaryseed | |
| Volunteer wheat (non imidazolinone-tolerant varieties) | |

2023-1399
2024-03-27

| | | |
|---|---------------------------|--|
| Wild oat (including Group 1 resistant) ¹ | | |
| Yellow foxtail | | |
| BROADLEAF WEEDS | RECOMMENDED STAGE | |
| Cow cockle | Cotyledon to 4 leaf stage | |
| Flixweed | | |
| Green smartweed | | |
| Lamb's quarters | | |
| Redroot pigweed | | |
| Prostrate pigweed | | |
| Shepherd's purse | | |
| Stinkweed | | |
| Stork's bill | | |
| Volunteer canola (non imidazolinone-tolerant varieties) | | |
| Wild mustard | | |
| WEEDS SUPPRESSED | | |
| Annual sow-thistle | | |
| Cleavers | | |
| Japanese brome | | |
| Wild buckwheat | | |

¹ DAVAI® 80 SL will not control weed biotypes that are multiple-resistant to both Group 1 and Group 2 Herbicides

MIXING INSTRUCTIONS

- Use 50-100 L/ha of water.
- Use a 50-mesh (or coarser) filter screen.
- Fill the spray tank three-quarters full with water.
- Add the required amount of DAVAI® 80 SL directly into the sprayer through the tank opening.
- Agitate until herbicide is thoroughly mixed.
- Continue agitation and add the required amount of the tank-mix partner.
- Continue agitation while adding the required amount of recommended Adjuvant.
- If excess foaming occurs, a silicone anti-foaming agent may be added (e.g. Halt®).
- Complete filling the tank to the desired level with water.
- Upon completion of spraying, thoroughly flush tank, boom, hoses and in-line and nozzle screens with clean water to avoid possible injury to other crops.
- Repeat sprayer cleanout process using an appropriate spray system cleaner.

ROTATIONAL CROPS

There is the possibility of residual soil activity for DAVAI® 80 SL the year following application. Initial crop injury to non imidazolinone-tolerant canola may be observed. Avoid spray overlap as yield reduction may result.

2023-1399
2024-03-27

Research studies have shown that the following crops may be safely planted the year following DAVAI® 80 SL application:

Spring Barley
Canary seed
Imidazolinone-tolerant Canola
Non imidazolinone-tolerant Canola
Chickpea
Field Corn
Field Pea
Flax
Lentil
Tame Oat
Soybean
Imidazolinone-tolerant Sunflower
Spring Wheat (including Durum wheat)

If tank-mixing DAVAI® 80 SL with another herbicide please consult the re-cropping label for the tank mix partner.

WARNING: Certain environmental conditions may delay the breakdown of herbicide residues in soil. These conditions include but are not limited to drought, extremes in soil pH and excessive cold. Under these conditions, the level of phytotoxic herbicide residues present in the field the season following an application may result in an increased potential for injury to succeeding crops to occur. This potential for increased residues under these environmental conditions is not unique to any specific herbicide or herbicide group but is a property of those herbicides which persist in the soil and are dependent on soil microbial activity and other non-microbial processes (e.g. hydrolysis) to breakdown. There are insufficient data for other follow crops. Conduct a field bioassay (a test strip grown to maturity) the year before growing any crop other than those listed above.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, DAVAI® 80 SL is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to DAVAI® 80 SL and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of DAVAI® 80 SL or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.

2023-1399
2024-03-27

- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

For further information or to report suspected resistance, contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 or at www.adama.com/canada.

Davai and Phantom are trademarks of an ADAMA Group Company.
All other products listed are registered trademarks of their respective companies.