



DAVAI A PLUS™

DAVAI® 80 SL, is conveniently packaged with ARROW ALL IN® to offer broad-spectrum control in dry beans, peas and soybeans.



GROUP 1 & 2

REGISTERED CROPS

- Dry beans
- Field peas
- Soybeans

WEEDS CONTROLLED AND APPLICATION TIMING

BROADLEAF WEEDS

Cotyledon – 4 leaf:

- Cleavers¹
- Cow cockle
- Flixweed
- Green smartweed
- Lamb's quarters
- Redroot pigweed
- Shepherd's purse
- Stinkweed
- Stork's bill
- Volunteer canola*
- Wild buckwheat¹
- Wild mustard

GRASSY WEEDS

1–6 true leaf:

- Barnyard grass
- Crabgrass (smooth, large)³
- Fall panicum
- Green foxtail²
- Japanese brome grass¹
- Persian darnel
- Proso millet
- Quackgrass¹
- Volunteer canary grass
- Volunteer canary seed⁴
- Volunteer cereals (barley, oats, wheat)
- Volunteer corn³
- Wild oats²
- Witchgrass
- Yellow foxtail

¹Suppression

²Including Group 1-resistant weeds. DAVAI A PLUS™ will NOT control weed biotypes that are multiple-resistant to both Group-1 and Group-2 herbicides.

³2- to 6-leaf

⁴1- to 4-leaf

*non imidazolinone-tolerant varieties

HOW IT WORKS

DAVAI A PLUS™ combines two (2) actives to tackle broadleaf and grassy weeds. See DAVAI® 80 SL and ARROW ALL IN® for more information.

Active Ingredients

Imazamox 80 g/L = SL
Clethodim 120 g/L = EC

Packaging

Co-pack includes:

- 1 x 4 L jug of DAVAI® 80 SL
- 1 x 6 L jug of ARROW ALL IN®

Application Rates & Acres Treated

Rate:

- DAVAI® 80 SL: 100 ml/ac
- ARROW ALL IN®: 150 ml/ac

Acres Treated: 40 ac/case

Water Volume

Ground: 40 L/ac (10 US gal/ac)

Aerial: Do not apply by air.

Rainfastness

3 hours



DAVAI A PLUS™

CROP STAGING

- Dry beans, Soybeans: Emergence to 3 expanded trifoliolate leaves
- Field peas: 1 – 6 true leaf

MIXING INSTRUCTIONS

1. Fill clean tank ½ to ¾ full of clean water and turn agitation on.
2. Add the required amount of ARROW ALL IN® herbicide and continue to agitate.
3. Add the required amount of DAVAI® 80 SL herbicide and continue to agitate.
4. Continue agitating and fill the remainder of the spray tank with water.

REGISTERED AND SUPPORTED TANK MIXES

None registered

ADJUVANT RATE

No adjuvant required.

CROP ROTATIONS

- | | | |
|---------------|--------------|-------------------------------------|
| • Barley | • Field peas | • Imidazolinone-tolerant sunflowers |
| • Canary seed | • Oats | • Wheat (spring) |
| • Canola | • Soybeans | |
| • Corn | | |

PRE-HARVEST INTERVAL

- Field peas: 60 days
- Dry beans: 75 days
- Soybeans: 85 days

GRAZING RESTRICTIONS

- Field peas: 30 days
- All other crops: Do not graze.

STORAGE

Do not freeze.

RECROPPING FOLLOWING AN IMIDAZOLINONE APPLICATION

There are several factors that affect the re-cropping following an imidazolinone application. These include (in order of importance):

1. **Product:** Imazethapyr, for example, is more persistent than imazamox.
2. **Soil moisture:** Requires more than 125 mm (5") of rain between herbicide application and August 31 in the year of application.
3. **Organic matter:** Brown soil zones (< 3% organic matter) are more susceptible to carryover crop injury the year after application.
4. **Rate:** Depending on the crop and rates, soil residues can be an issue.
5. **Soil pH:** Product persists longer in a pH < 5.5 – 6.

Depending on the following crop the level of sensitivity will vary. Please contact your local ADAMA Area Business Manager for more details.

WANT TO KNOW MORE?

Click on the QR code to access all the product details including the label and SDS documents



Always read and follow label directions.
1.855.264.6262 ADAMA.COM

®/™ARROW ALL IN and DAVAI are registered trademarks and DAVAI A PLUS is a trademark of ADAMA Agricultural Solutions Canada Ltd.
All other products are trademarks of their respective companies.
© 2024 ADAMA Agricultural Solutions Canada Ltd.

1/2024