



ADAMA

BROMOTRIL®

Tough broadleaf weed control with tank-mix flexibility and excellent crop safety



GROUP 6

Active Ingredient

Bromoxynil Octanoate Ester
240 g/L = EC

Packaging

Case: 2 x 9.7 L jugs

Application Rates & Acres Treated

PRE-PLANT:

Rate: 490 ml/ac
Area Treated: 20 ac/jug

IN-CROP BROADLEAF:

Rate: 490 – 570 ml/ac
Area Treated: 17 – 20 ac/jug

Water Volume

Ground: 20 – 80 L/ac
(5 – 20 US gal/ac)

Aerial:

8 – 16 L/ac (3 – 5 US gal/ac)
Wheat and barley only
Do not apply pre-plant

Rainfastness

30 minutes

REGISTERED CROPS

Crop	Stage
Barley, Oats, Wheat	Pre-seed burn-off with glyphosate
Crop	Crop Leaf Stage
Alfalfa (seedling)	2 – 6 trifoliolate
Alfalfa (established for seed production only)	Spring: before the crop begins to shield the weeds
Barley, Oats, Triticale, Wheat (spring)	2-leaf to early flag
Winter wheat	Fall: 2- to 4-leaf Spring: first growth to early flag
Corn (field, sweet)	Beyond 4-leaf
Fall rye	Spring: from first growth to early flag
Forage millet, Sorghum	4-leaf to 8 inches

WEEDS CONTROLLED

Seedling up to 4-leaf stage:

- American nightshade
- Bluebur
- Cocklebur
- Common ragweed
- Cow cockle¹
- Green smartweed
- Kochia²
- Lady's thumb
- Pale smartweed
- Pigweed¹
- Russian thistle²
- Stinkweed¹
- Velvetleaf³
- Wild mustard¹

Seedling up to 8-leaf stage:

- Common buckwheat
- Common groundsel
- Lamb's quarters
- Tartary buckwheat
- Wild buckwheat

¹ In normal conditions, it will be controlled up to 4-leaf stage. Plants beyond this stage are unlikely to be controlled; the higher rate generally gives better results.

² Spray before plants are 2 inches high.

³ Spray before plants are 3 inches high.



BROMOTRIL®

HOW IT WORKS

BROMOTRIL® is a contact herbicide which controls Group 2- and Group 9-resistant biotypes. Leaves will yellow in 2–4 days with complete control in 7–14 days.

REGISTERED AND SUPPORTED POST-EMERGENT TANK MIXES

Post-emergent Herbicides:

- Spring wheat: 2,4-D ESTER 700, MCPA ESTER 600, BISON® 400 L
- Winter wheat: 2,4-D ESTER 700, MCPA ESTER 600, BISON® 400 L
- Barley: 2,4-D ESTER 700, MCPA ESTER 600, BISON® 400 L
- Oats: MCPA ESTER 600
- Corn: Accent®, Atrazine, Banvel®(Dicamba), Ultim®
- Fall rye: MCPA ESTER 600

Pre-seed:

- Glyphosate

Fungicides:

- BUMPER® 432 EC

MIXING INSTRUCTIONS

1. Fill spray tank ½ full with water.
2. Add the required amount of BROMOTRIL®. Begin agitation.
3. If tank-mixing, add any tank-mix partner(s) to the spray tank, agitate first and then add BROMOTRIL® (unless otherwise directed by the BROMOTRIL® and/or tank-mix partner label).
4. Add the remaining amount of water while agitation continues.

CROP ROTATIONS

No restrictions

STORAGE

Do not freeze.

PRE-HARVEST INTERVAL

30 days

GRAZING RESTRICTIONS

- Do not use treated crops for grazing of livestock or green feed until 30 days after application.
- Do not cut treated crops for forage until 30 days after application.

QUICK TIPS:

Avoid spraying if temperatures are above 25° C. Leaf scorching may occur in corn and flax if applied during or after adverse growing conditions, such as cool and wet or hot (above 27° C) and humid weather. For best results, spray when weeds are in the seedling stage.



ADAMA

Always read and follow registered product label instructions. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.
1.855.264.6262 ADAMA.COM

®/™ BISON, BROMOTRIL, and BUMPER are registered trademarks of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies.

© 2023 ADAMA Agricultural Solutions Canada Ltd.