



ADAMA

2,4-D ESTER 700

Provides reliable post-emergent control of broadleaf weeds and great tank-mix flexibility in wheat, barley, rye and corn.



GROUP 4

Active Ingredient:

2,4-D Ester EH Ester
660 g/L = EC

Application Rates and Acres Treated:

Rate: 200 – 600 ml/ac

Acres Treated: 17 – 50 ac/jug;
1665 – 5000 ac/tote

Packaging:

Case: 2 × 10 L jugs
Tote: 1000 L

Water Volume:

Ground: 12.5 – 50 L/ac
(5 – 15 US gal/ac)

Aerial: Minimum 12 L/ac (3 US gal/ac)

Rainfastness:

2 hours

APPLICATION TIMING:

2,4-D ESTER 700 can be used in pre-plant, in-crop or post-harvest applications.

REGISTERED CROPS:

- Barley
- Field Corn
- Rye (spring, fall)
- Wheat (spring, winter)

See registered label for timing and rates for each crop.

WEEDS CONTROLLED:

Susceptible weeds:

- | | | |
|-----------------------------------|--|----------------------------|
| • Annual Sow-Thistle ¹ | • Lamb's-Quarters | • Shepherd's Purse |
| • Bluebur | • Mustards
(except dog and tansy) | • Stinging Nettle |
| • Burdock ¹ | • Narrow-Leaved
Hawk's-Beard ² | • Stinkweed |
| • Cocklebur | • Plantain | • Sweet Clover (seedling) |
| • Daisy Fleabane | • Prickly Lettuce | • Thyme-Leaved
Spurge |
| • False Flax | • Ragweed | • Volunteer Canola |
| • False Ragweed | • Redroot Pigweed | • Wild Radish |
| • Flixweed | • Russian Pigweed | • Wild (prairie) Sunflower |
| • Giant Ragweed | • Russian Thistle | |
| • Goat's-Beard | | |
| • Kochia | | |

Harder-to-control weeds:

- | | | |
|----------------------------|--|-----------------------------------|
| • Curled Dock ¹ | • Heal-All | • Prostrate Pigweed |
| • Dog Mustard | • Knotweed ¹ | • Purslane |
| • Field Peppergrass | • Narrow-Leaved
Hawk's-Beard ⁴ | • Sheep Sorrel |
| • Flixweed ⁴ | • Oak-Leaved
Goosefoot | • Tansy Mustard |
| • Groundsel | • Pineapple Weed | • Tumble Pigweed |
| • Hairy Galinsoga | | • Velvetleaf |
| • Hawkweed | | • Volunteer Canola ^{1,3} |

¹All types

²In the fall, and at the 1- to 2-leaf stage in spring

³4- to 6-leaf stage

⁴If treated before bolting in spring



2,4-D ESTER 700

Very-hard-to-control weeds:

- Biennial Wormwood
- Blue Lettuce
- Bull Thistle
- Burdock
- Buttercup
- Canada Thistle
- Chicory
- Curled Dock
- Dandelion
- Field Bindweed
- Field Chickweed²
- Field Horsetail²
- Gumweed
- Hedge Bindweed
- Hemp-Nettle² (if treated before the 4-leaf stage)
- Hoary Cress
- Lady's Thumb²
- Leafy Spurge
- Mouse-Eared Chickweed²
- Perennial Sow Thistle
- Russian Knapweed
- Scentless Mayweed
- Smartweed²
- Tartary Buckwheat
- Teasel
- Volunteer Sunflower
- Wild Buckwheat²
- Yellow Rocket¹

¹1- to 3-leaf stage

²Use highest listed rate for suppression

HOW IT WORKS:

Systemic activity hinders plant cell growth in newly forming stems and leaves promoting uncontrolled, unsustainable growth, causing stem curl-over, leaf withering and eventual plant death.

REGISTERED AND SUPPORTED TANK MIXES:

- BROMOTRIL®
- BUMPER® 432 EC
- Glyphosate
- INVOLVE® 50 WDG

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

MIXING INSTRUCTIONS:

1. Fill sprayer tank ½ full of water.
2. Start sprayer tank agitation.
3. Add the required amount of 2,4-D ESTER 700 into the sprayer.
4. Agitate until the herbicide is thoroughly mixed.
5. Continue agitation while adding any required adjuvants or surfactants for tank-mix partners.
6. Complete filling the tank with water as agitation continues.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

CROP ROTATIONS:

No restrictions.

RE-ENTRY INTERVAL:

Not applicable.

PRE-HARVEST INTERVAL:

90 days

STORAGE:

May be stored at any temperature. Shake well before using.

GRAZING RESTRICTIONS:

30 days



[CLICK HERE FOR FULL PRODUCT DETAILS.](#)



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

BROMOTRIL®, BUMPER® and INVOLVE® are registered trademarks of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2025 ADAMA Agricultural Solutions Canada Ltd.

QUICK TIPS

Avoid spray drift to any off-target vegetation. Coarse sprays are less likely to drift. Do not spray during periods of high winds.