



ADAMA

# FOLPAN® 80 WDG

A water-dispersible granular fungicide that delivers exceptional protection against the most common and economically damaging diseases in grapes and other fruit, vegetable and ornamental crops.



## GROUP M4

### Active Ingredients:

80% Folpet = WDG

### Application Rates and Acres Treated:

Rate: 0.5-2 kg/ac

Acres Treated: 2.5-10 ac/pack

### Packaging:

Case: 2 x 5 kg packs

### Water Volume:

Ground: 400-1200 L/ac  
(100-317 US gal/ac)

Aerial: Do not apply by air.

### Rainfastness:

N/A

[CLICK HERE FOR FULL PRODUCT DETAILS.](#)



### REGISTERED CROPS:

- Apples
- Azalea
- Carnation
- Chrysanthemum
- Crabapples
- Cranberries
- Field Cucumbers
- Field tomatoes
- Grapes
- Iris
- Marigold
- Melons
- Poinsettia (greenhouse)
- Pumpkins
- Snapdragon
- Squash
- Strawberries
- Zinnia

### KEY DISEASES CONTROLLED:

- Alternaria Leaf Spot
- Anthracnose
- Black Rot
- Brooks Spot
- Dead Arm
- Didymellina Leaf Spot
- Downy Mildew
- Fly Speck
- Fruit Rot
- Grey Mould
- Leaf Spot
- Phythium Root Rot
- Powdery Mildew
- Scab
- Septoria Leaf Spot
- Sooty Blotch
- Stem Rot

### HOW IT WORKS:

Multi-site-contact mode of action with protective ability. Use in a regularly scheduled maintenance program.

### REGISTERED AND SUPPORTED TANK MIXES:

#### Kumulus® DF

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.



# FOLPAN® 80 WDG

## APPLICATION TIMING AND CROP STAGING:

Crop	Diseases	Timing
Apples, Crabapples	Alternaria Leaf Spot, Black Rot, Brooks Spot, Fly Speck, Scab, Sooty Blotch	Apply in a regular protective schedule from green tip up until harvest. Do not apply more than 6 applications per season. See label for additional precautions.
Cranberries	Fruit Rot	Apply when 5% of blossoms are open. Repeat 10–14 days later when 50–75% of blossoms are open.
Grapes	Phomopsis	Apply when new shoots are 1–2 inches long and repeat when growth is 4–6 inches long.
	Black Rot, Downy Mildew	Apply just before bloom, just after bloom and in first cover spray. For downy mildew, an additional application 2–3 weeks later may be needed.
	Powdery Mildew	Apply in a tank mix with Kumulus® DF at first sign of mildew and repeat after 10 days.
Field Cucumbers, Melons, Pumpkins, Squash	Anthrachnose, Downy Mildew	Apply when first true leaves appear. Repeat at 7-day intervals until crop is harvested.
Field Tomatoes	Anthrachnose	Apply during first bloom and repeat at 7-day intervals until harvest.
Strawberries	Grey Mould, Fruit Rot, Leaf Spot	Apply before first infection and repeat at 7-day intervals to protect crop until harvest.
Carnation, Poinsettia, Azalea, Marigold, Zinnia, Chrysanthemum, Iris, Snapdragon	Blight, Phythium Root Rot, Stem Rot, Alternaria Leaf Spot, Septoria Leaf Spot, Didymellina Leaf Spot, Anthracnose (depending on ornamental)	Generally, apply when ornamental emerges and repeat at regular intervals. Consult label for timing on specific disease and ornamental.

### RE-ENTRY AND PRE-HARVEST INTERVALS:

- Apples, Crabapples: 1 day
- Cranberries: 30 days
- Field cucumbers, Melons, Pumpkins, Squash: 11 days
- Greenhouse ornamentals grown for cut flower production: 48 days
- Greenhouse ornamentals not grown for cut flower production: 12 hours
- Ornamentals grown for cut flower production: 16 days

### MIXING INSTRUCTIONS:

1. Fill spray tank nearly full.
2. Start sprayer agitation.
3. Pour recommended amount of FOLPAN® 80 WDG on surface of water. Fungicide can be premixed in a bucket 1/2 filled with water. Mix can be poured through screen into nearly filled spray tank.
4. Finish filling tank.
5. Keep agitator running during filling and spraying.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

### Mechanical harvesting\*:

- Field tomatoes, Grapes, Strawberries: 1 day

### Hand harvesting\*

- Grapes: 3 days
- Strawberries: 11 days
- Field tomatoes: 22 days

\* Crops listed have different mechanical/hand harvesting intervals.

NOTE: Refer to the label for pre-harvest intervals for all other registered crops.



Always read and follow label directions.  
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

FOLPAN® is a registered trademarks of an ADAMA Group Company.  
All other products are trademarks of their respective companies.  
© 2025 ADAMA Agricultural Solutions Canada Ltd.