



LEOPARD®

ADAMA's grass control product to control hard-to-kill grassy weeds in dry beans, soybeans and cucurbit vegetable crops.



GROUP 1

Active Ingredient:

Quizalofop-P-ethyl 100 g/L = EC

Application Rates and Acres Treated:

Rate: 150 – 290 ml/ac;
standard rate 195 ml/ac

Acres Treated: 30 – 50 ac/jug;
standard 40 ac/jug;
480 ac/drum;
2500 ac/tote

Packaging:

Case: 2 × 7.8 L jugs;
Bulk: 93.6 L drums;
Tote: 487.5 L tote

Water Volume:

Ground: 40 L/ac (10 US gal/ac)
Aerial: 10 L/ac (2.5 US gal/ac)

Rainfastness:

1 hour

KEY BENEFITS:

- Flexible tank-mix options
- One solution for grassy weed control in multiple crops, including canola, pulses, soybeans, flax and more
- Controls hard to kill winter annual and perennial grasses

REGISTERED CROPS

- Alfalfa, seed
- Beans (dry & snap)
- Canola
- Chickpeas
- Peas (field & processing)
- Flax
- Hemp (grown for fibre, seed & oil)
- Lentils
- Mustard (oriental, yellow & brown)
- Soybeans

WEEDS CONTROLLED:

- Barnyard Grass
- Downy Brome
- Fall Panicum
- Foxtail Barley
- Green Foxtail
- Japanese Brome
- Proso Millet
- Quackgrass (suppression)
- Quackgrass (control)
- Volunteer Cereals (wheat, barley, oats)
- Volunteer Corn
- Wild Oats
- Witchgrass
- Yellow Foxtail

REGISTERED AND SUPPORTED TANK MIXES:

- Basagran®
- ADAMA Glufosinate
- DAVAI® 80 SL
- Glyphosate
- PHANTOM® 240 SL
- PYTHON®

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.



**CLICK HERE
FOR FULL
PRODUCT
DETAILS.**

LEOPARD®

HOW IT WORKS:

LEOPARD® is a selective postemergence herbicide for the control of annual and perennial grasses. LEOPARD® is a systemic herbicide which is rapidly absorbed and readily translocated from the treated foliage to the root systems and growing points of the plant. Treated plants show a reduction in growth and a loss of competitiveness. An early yellowing or browning of the younger plant tissues is followed by a progressive collapse of the remaining foliage. These symptoms will generally be observed in one to three weeks depending on the grass species treated and the environmental conditions. This product does not control sedges or broadleaf weeds.

CROP STAGING:

- Most crops are tolerant at all stages, so target applications at the optimal weed stage
- Always adhere to the pre-harvest interval for each crop

MIXING INSTRUCTIONS:

1. Thoroughly clean the sprayer by flushing the system with water containing detergent.
2. Fill clean spray tank ½ full with water. Start agitation.
3. If tank mixing LEOPARD® with another pesticide, add tank-mix partner followed by adjuvant.
4. Ensure that the herbicide is completely mixed before proceeding to the next step.
5. Add the rest of the required water to the tank. Mix well before applying to the crop.

*When mixing with glufosinate: Glufosinate + LEOPARD® + Surfactant

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

On repeat tank loads, ensure that the amount of spray solution left in the tank from the previous load is less than 10% of volume about to be mixed. Do not mix, load or clean spray equipment where there is a potential to contaminate wells or aquatic systems.

PRE-HARVEST INTERVALS:

- Beans (dry & snap), Cucurbits, Rutabagas: 30 days
- Canola: 64 days
- Peas, Small Red Beans: 65 days
- Soybeans: 80 days
- Sugarbeets: 60-80 days

ADJUVANT RATES:

LEOPARD® is not packaged with but requires an adjuvant such as:

- Merge® @ 0.5–1.0% v/v
- LI 700® @ 0.25–0.5% v/v
- Liberate™ @ 0.5% v/v
- Companero® Adjuvant @ 0.5–1.0% v/v

Or other non-ionic or methylated seed oil adjuvants

GRAZING RESTRICTIONS:

Do not cut treated crops for hay.

RE-ENTRY INTERVAL:

12 hours

STORAGE:

Do not freeze.

CROP ROTATIONS:

No restrictions.



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

LEOPARD®, DAVAI®, PHANTOM® and PYTHON® are registered trademarks of ADAMA Group Company. All other products are trademarks of their respective companies.
© 2025 ADAMA Agricultural Solutions Canada Ltd.

QUICK TIPS:

Apply to wild oats before tillering for best results.

Use the higher adjuvant rate when targeting quackgrass or wild oats, or when conditions are not conducive to good growth.