

# **TOPNOTCH**

ADAMA

Broad-spectrum disease control in multiple crops including cereals, field peas, edible beans and soybeans.

### **GROUP 3 & 11**

# **Active Ingredient:**

Azoxystrobin 143 g/L and Propiconazole 124 g/L = SC

# Application Rates and Acres Treated:

Rate: 210 – 620 ml/ac
Acres Treated: 14 – 40 ac/jug

# Packaging:

Case: 2 × 8.6 L jugs

### **Water Volume:**

Ground: 40 L/ac

(10 US gal/ac)

Aerial: 20 L/ac (5 US gal/ac)

### Rainfastness:

Avoid applying when heavy rainfall is in the forecast.

### **KEY BENEFITS:**

- Multiple modes of action for resistance management
- Curative and preventative systemic action
- Can use on multiple crops, with exceptional crop safety
- · Registered for both ground and aerial spraying
- Excellent Crown Rust control in oats

### **REGISTERED CROPS:**

- Barley
- Edible Beans
- Field Peas
- Lentils
- Oats
- Rye

- Soybeans
- Triticale
- Wheat

### **KEY DISEASES CONTROLLED:**

- Anthracnose
- Ascochyta Blight
- Barley Leaf Rust
- Mycosphaerella Blight
- Net and Spot Blotches
- Powdery Mildew

- Scald
- Septoria Spot
- Stripe Rust
- Tan Spot
- Wheat Leaf Rust
- White Mould1

 $^{1} Suppression \\$ 

### APPLICATION TIMING AND CROP STAGING:

Crop	Diseases	Application Timing	Rate
Barley	Barley Net Blotch, Barley Scald, Septoria Leaf Spot, Stripe Rust, Barley Leaf Rust, Tan Spot	Apply once between stem elongation and half-head emergence (Growth stage 29 – 55).	210 ml/ac
Oats	Barley Net Blotch, Crown Rust, Septoria Leaf Spot		
Rye	Septoria Leaf Spot, Barley Scald, Tan Spot		
Triticale	Septoria Leaf Spot, Tan Spot		
Wheat	Septoria Leaf Spot, Tan Spot, Stripe Rust, Wheat Leaf Rust		
Beans, Field Peas, Lentils, Soybeans	Mycosphaerella Blight, Anthracnose	Make the first application at the first sign of disease. Apply the high rate only under conditions of high disease pressures. A second application 14 days later may be needed if conditions persist. Good spray coverage and canopy penetration are important for best results.	310 – 620 ml/ac
	Powdery Mildew, White Mould (suppression only)		310 ml/ac



# **TOPNOTCH™**

#### **HOW IT WORKS:**

Used as both a curative and preventative fungicide,  $TOPNOTCH^{\text{\tiny{M}}}$  has broad-spectrum, systemic and contact activity.

### **REGISTERED AND SUPPORTED TANK MIXES:**

- ARROW ALL IN®
- Coragen®
- Decis®

- Glufosinate
- LEOPARD®
- Poast Ultra®

- SILENCER® 120 EC
- Voliam Xpress®
- ZIVATA®

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

### **MIXING INSTRUCTIONS:**

- 1. Fill spray tank  $\frac{1}{2} \frac{2}{3}$  full with water.
- 2. With agitator running, add required amount of TOPNOTCH™ and continue agitating while adding remainder of the water.
- 3. Begin application after TOPNOTCH™ is completely dispersed into the mix water, and maintain agitation during spraying operation.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol

### **CROP ROTATIONS:**

Do not plant any other crop intended for food, grazing or any component of animal feed or bedding within 105 days of application.

### **PRE-HARVEST INTERVALS:**

- Beans, Field Peas, Lentils, Soybeans: 30 days
- Cereals: 45 days

### **GRAZING RESTRICTIONS:**

Do not graze pea vines. 30 days for all other crops.

### **RE-ENTRY INTERVAL:**

12 hours

### **STORAGE:**

Do not freeze.

CLICK HERE FOR FULL PRODUCT DETAILS.





Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

ARROW ALL IN®, LEOPARD®, SILENCER® and ZIVATA® are registered trademarks, and TOPNOTCH® is a trademark of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2025 ADAMA Agricultural Solutions Canada Ltd.

## **QUICK TIPS:**

Good spray coverage and canopy penetration are important to achieve the best results.