



ADAMA

BADGE®

Get proven control of almost 30 broadleaf weeds in cereals and corn with easy-to-use tank-mix options for one-shot weed control.



GROUP 4 & 6

Active Ingredients:

Bromoxynil 225 g/L and MCPA 2 EH Ester 600 225 g/L = EC

Application Rates and Acres Treated:

Rate: 500 ml/ac

Acres Treated: 20 ac/jug

Packaging:

Case: 2 × 10 L jugs

Water Volume:

Ground:

Cereals: 20-40 L/ac
(5-10 US gal/ac)

Corn: 80-120 L/ac
(20-30 US gal/ac)

Forages: 60 L/ac
(15 US gal/ac)

Aerial:

8-20 L/ac
(2-5 US gal/ac)

Rainfastness:

1 hour

REGISTERED CROPS:

- Barley
- Corn
- Fall Rye
- Oats
- Wheat (spring, winter)

WEEDS CONTROLLED:

- American Nightshade
- Ball Mustard
- Bluebur
- Canada Thistle¹
- Cocklebur
- Common Buckwheat
- Common Groundsel
- Common Ragweed
- Cow Cockle²
- Flixweed
- Green Smartweed
- Kochia³
- Lady's Thumb
- Lamb's-Quarters
- Night-Flowering Catchfly
- Pale Smartweed
- Perennial Sow Thistle¹
- Redroot Pigweed
- Russian Thistle³
- Scentless Chamomile⁴
- Shepherd's Purse
- Stinkweed
- Tartary Buckwheat
- Velvetleaf⁵
- Volunteer Canola (all types)
- Volunteer Sunflower
- Wild Buckwheat
- Wild Mustard
- Wild Tomato
- Wormseed Mustard

¹Top growth control.

²Up to 4-leaf stage.

³Spray before plants are 2 inches high.

⁴Spring annual only.

⁵Spray before plants are 3 inches high.



BADGE®

HOW IT WORKS:

A combination of systemic and contact activity with weeds yellowing within 2–4 days and exhibiting abnormal growth (twisting and cupping of leaves) in 2–10 days.

CROP STAGING:

Crop	Stage
Barley, Oats, Spring Wheat	2 leaf to early flag
Corn	4–6 leaf
Fall Rye	When growth commences in spring to early flag leaf.
Timothy (established for seed production)	Prior to shot blade in the seed production year.
Winter Wheat (includes underseeded with red clover)	2–4 leaf stage in the fall or after growth resumes up to early flag leaf.

REGISTERED AND SUPPORTED TANK MIXES:

- **Corn:** Atrazine
- **Oats:** MCPA Ester 600
- **Spring Wheat and Barley:** MCPA Ester 600, Refine® SG
- **Winter Wheat:** MCPA Ester 600 or Refine® SG

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

MIXING INSTRUCTIONS:

1. Fill sprayer tank ½ full of water.
2. Start sprayer tank agitation.
3. Add the required amount of BADGE®.
4. If necessary, add any required adjuvants or surfactants for tank-mix partners.
5. Complete filling the tank with water as agitation continues.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

CROP ROTATIONS:

No re-cropping restrictions the year after treatment.

PRE-HARVEST INTERVAL:

30 days for feed. Refer to tank-mix partner label.

STORAGE:

Do not freeze.

GRAZING RESTRICTIONS:

- Do not graze treated grain or established Timothy crops or cut for feed within 30 days of application.
- Do not graze meadow foxtail in the year of treatment.
- Do not graze other treated forage grasses within 56 days of treatment.

[CLICK HERE FOR FULL PRODUCT DETAILS.](#)



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

BADGE®, is a registered trademarks of an ADAMA Group Company.
All other products are trademarks of their respective companies.
© 2025 ADAMA Agricultural Solutions Canada Ltd.

QUICK TIPS:

BADGE® herbicide is well known for being gentle on the crop. Avoid spraying if temperatures are above 25°C.