



ADAMA

VANTANA™

A protective, broad-spectrum fungicide used in a variety of fruit and vegetable crops. This Group 29 fungicide is a very important resistance management tool with activity on the hardest-to-control fungal diseases.



GROUP 29

Active Ingredient:

Fluazinam 500 g/L = SC

Application Rates & Acres Treated:

Rate: 162 – 906 ml/ac

Acres Treated: 27 – 80 ac/jug

Packaging:

Case: 2 x 10 L jugs

Water Volume:

Ground: Varies with the crop.
Refer to the label for more details.

Aerial: Minimum of 18.2 L/ac

Rainfastness:

Avoid application when heavy rain is forecast.

KEY BENEFITS:

- Resistance Management Tool
- Can be used on multiple crops
- Broad spectrum disease control
- Multiple rates for precise control

REGISTERED AND SUPPORTED CROPS:

- Apples
- Blueberries (highbush, lowbush)
- Bulb Onions
- Carrots
- Currant
- Dry-Shelled Beans
- Elderberry
- Edible-Podded Legume Vegetables (except peas)
- Field Peppers
- Ginseng
- Gooseberry
- Huckleberry
- Head and Stem Brassica (Crop sub-group 5A)
- Leafy Brassica Greens (Crop sub-group 5B)
- Potatoes
- Soybeans

For a full list of registered crops, please refer to the label.

HOW IT WORKS:

VANTANA™ works by inhibiting fungal adenosine triphosphate (ATP) production in the mitochondria (impairing energy production).

Active on a wide range of diseases, VANTANA™ is a protective fungicide that inhibits the germination of fungal spores.

MIXING INSTRUCTIONS:

1. Fill sprayer tank ½ full of water.
2. Start sprayer tank agitation.
3. Add the required amount of VANTANA™.
4. If applicable, add any required adjuvants or surfactants for tank-mix partners.
5. Complete filling tank with water.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

[CLICK HERE FOR FULL PRODUCT DETAILS.](#)





VANTANA™

APPLICATION TIMING AND KEY DISEASES CONTROLLED BY CROP:

NOTE: For application timing by crop, please refer to the label.

Crop	Disease	Rate
Apples	Cedar Apple Rust, Apple Scab, Flyspeck, Sooty Blotch	304–405 ml/ac (0.75–1.0 L/ha)
	For Suppression of MITES: Two-spotted Spider Mite, European Red Mite and Apple Rust Mite	
	Black Rot*, Brooks Spot*, Quince Rust*	405 ml/ac (1.0 L/ha)
Blueberries (lowbush)	Valdensinia Leaf Spot	462–324 ml/ac (0.4–0.8 L/ha)
Blueberries (highbush, lowbush)	Mummyberry*, Phomopsis Fruit Rots* and Fruit Anthracnose*	907 ml/ac (2.24 L/ha)
Currant, Elderberry, Gooseberry, Huckleberry	Fruit Anthracnose*	
Carrots	Alternaria Leaf Blight	469 ml/ac (1.16 L/ha)
Bulb Onions	Purple Blotch, Botrytis Leaf Blight	469 ml/ac (1.16 L/ha) (584 g)
Dry-Shelled Beans	White Mould	243–405 ml/ac (0.6–1.0 L/ha)
	Anthracnose	
Edible-Podded Legume Vegetables (except pea)	White Mould	
Field Peppers	Phytophthora Blight	688 ml/ac (1.7 L/ha)
Ginseng	Rhizoctonia Root Rot	486 ml/ac (1.2 L/ha)
	Alternaria Blight, Botrytis Blight	
Potatoes	Late Blight	162–242 ml/ac (0.4–0.6 L/ha)
	White Mould	356–473 ml/ac (0.88–1.17 L/ha)
Soybeans	White Mould	356–473 ml/ac (0.88–1.17 L/ha)
	White Mould*	178 ml/ac (0.44 L/ha)
Leafy Brassica Greens (Crop sub-group 5B) Pre-transplant treatment	Clubroot	1.17 L in 202 L of water per acre (2.9 L in 500 L of water per ha)
Head and Stem Brassica (Crop sub-group 5A), Leafy Brassica Greens (Crop sub-group 5B) Transplant treatment		Mix 50 ml with water to make a 100 L solution.

* Suppression

CROP ROTATIONS:

Areas treated with VANTANA™ may be replanted with potatoes and dry shelled beans as soon as practical after the last application. Other root crops and leafy vegetables can be planted 30 days after the last application. All other crops can be planted 70 days after the last application.

GRAZING RESTRICTIONS:

Do not feed treated foliage to livestock.

RE-ENTRY INTERVALS:

- Bulb Onions: 3 days
- All other crops: 24 hours

REGISTERED AND SUPPORTED TANK MIXES:

None registered

STORAGE:

Do not freeze.

PRE-HARVEST INTERVALS:

- Bulb Onions, Carrots: 7 days
- Edible-Podded Legume Vegetables (except peas), Potatoes: 14 days
- Apples: 28 days
- Blueberries (highbush, lowbush), Currant, Dry-Shelled Beans, Elderberry, Field Peppers, Ginseng, Gooseberry, Huckleberry Leafy Brassica Greens (Crop Subgroup 5B): 30 days
- Head and Stem Brassica (Crop Subgroup 5A): 65 days
- Soybeans: DO NOT apply after growth stage R3

QUICK TIPS:

For optimal efficacy VANTANA™ requires good coverage, best achieved with higher water volumes. Actual water volumes will vary with the crop, please refer to the label for more specific details.



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

VANTANA™ is a trademark of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2025 ADAMA Agricultural Solutions Canada Ltd.