



INSECTICIDE



ADAMA

# CORMORAN®

A multi-mode of action foliar insecticide for rapid knockdown and extended residual control of all damaging life stages of pests in an easy-to-use formulated mixture.



## GROUP 4 & 15

### Active Ingredients:

Novaluron 100 g/L,  
Acetamiprid 80 g/L = EC

### Application Rates:

Rate: 440–750 ml/ha

### Packaging and Acres Treated:

Case: 2 × 10.08 L jugs

Acres Treated: 13–23 ac/jug

### Water Volume:

Ground: 200 L/ha (80 L/ac)

Aerial: Do not apply by air.

### Rainfastness:

Avoid application when heavy rain is forecast.

### HOW IT WORKS:

CORMORAN® kills insect eggs by contact and larvae by ingestion. Containing two modes of action, CORMORAN® provides both rapid knockdown and residual control of insect pests.

### KEY INSECTS CONTROLLED IN POTATOES:

- Aphids
- Armyworm
- Cabbage Looper
- Colorado Potato Beetle
- European Corn Borer

### CROP STAGING AND RATES:

The need and timing of an application should be based on the presence of pests at vulnerable developmental stages and significant populations, as determined by local monitoring.

Crop	Insects Controlled	Rate (ml/ha)*	Application Instructions
Potatoes	Colorado Potato Beetle (CPB)	440–700	Minimum re-application interval of 7 days. DO NOT apply more than once every 10 to 14 days.
	Aphids	490	
	European Corn Borer	650–750	
	Armyworm, Cabbage Looper	440–750	Do not apply more than 1.68 L/ha per season. For CPB, do not apply more than twice to a single generation and do not apply to successive generations.

\* Use higher rate for larger infestations.

NOTE: Apply in a minimum finished spray volume of 200 L/ha (80 L/ac) by ground.

### MIXING INSTRUCTIONS:

1. Fill sprayer tank ½ full of water.
2. Start sprayer tank agitation.
3. Add the required amount of CORMORAN®.
4. If necessary, add any required adjuvants or surfactants for tank-mix partners.
5. Complete filling tank with water.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.



FOR FULL PRODUCT DETAILS, SCAN THE QR CODE.



INSECTICIDE

# CORMORAN®

## PRE-HARVEST INTERVALS:

Potatoes: 7 days

## CROP ROTATIONS:


No restrictions.

## RE-ENTRY INTERVAL (REI):


Potatoes: 12 hours all activities

## CORMORAN® Research Results:

### Colorado Potato Beetle Control in Potatoes (Bright, ON – BlackCreek) 2025



### Colorado Potato Beetle Control in Potatoes (Bright, ON – BlackCreek) 2025



Always read and follow label directions.  
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

CORMORAN® is a registered trademark of an ADAMA Group Company. All other products are trademarks of their respective companies.  
© 2026 ADAMA Agricultural Solutions Canada Ltd.