




ADAMA

TOPNOTCH™

A new broad-spectrum, multi-mode of action option for control of cereal leaf diseases as well as dangerous diseases in dry beans and soybeans.



Active Ingredient:

Azoxystrobin and Propiconazole

Fungicide Groups:

Group 3 and Group 11

Formulation:

Azoxystrobin 143 g/L and Propiconazole 124 g/L = SC

Application Rates and Packaging:

- 0.21 – 0.62 ml/ac
- 135 – 410 ac/jug
- 2 x 8.6 L jugs/case

Registered Crops:

- Barley
- Edible beans
- Field peas
- Lentils
- Oats
- Rye
- Soybeans
- Triticale
- Wheat

KEY BENEFITS:

- Multiple modes of action for resistance management
- Curative and preventative systemic action
- Can be used on multiple crops, with exceptional crop safety
- Registered for both ground and aerial spraying
- Excellent crown rust control in oats

KEY DISEASES CONTROLLED:

- Anthracnose
- Ascochyta blight
- Barley leaf rust
- Mycosphaerella blight
- Net and spot blotches
- Powdery mildew
- Scald
- Septoria spot
- Stripe rust
- Tan spot
- Wheat leaf rust
- White mould¹

HOW IT WORKS:

Used as both a curative and preventative fungicide, TOPNOTCH™ has broad-spectrum, systemic and contact activity.

APPLICATION TIMING AND CROP STAGING:

Crop	Diseases	Application Timing	Rate
Barley	Barley net blotch, Barley scald, Septoria leaf spot, Stripe rust, Barley leaf rust, Tan spot	Apply once between stem elongation and half-head emergence (Growth stage 29 – 55).	0.21 L/ac
Beans, Field peas, Lentils, Soybeans	Mycosphaerella blight, Anthracnose	Make the first application at the first sign of disease. Apply the high rate only under conditions of high disease pressures. A second application 14 days later may be needed if conditions persist. Good spray coverage and canopy penetration are important for best results.	0.31 L/ac – 0.62 L/ac
	Powdery mildew, White mould (suppression only)		0.31 L/ac
Oats	Barley net blotch, Crown rust, Septoria leaf spot	Apply once between stem elongation and half-head emergence (Growth stage 29 – 55).	0.21 L/ac


¹Suppression only




TOPNOTCH™

Crop	Diseases	Application Timing	Rate
Rye	Septoria leaf spot, Barley scald, Tan spot	Apply once between stem elongation and half-head emergence (Growth stage 29 – 55).	0.21 L/ac
Triticale	Septoria leaf spot, Tan spot		
Wheat	Septoria leaf spot, Tan spot, Stripe rust, Wheat leaf rust		
Durum wheat	Septoria leaf spot, Tan spot, Stripe rust		

PEA GROWTH STAGES




SOYBEAN GROWTH STAGES




TOPNOTCH™

LENTILS GROWTH STAGES



CEREAL GROWTH STAGES



Timing based on using the product alone. Please refer to tank-mix partners label for crop timing.

Always read and follow label directions.



TOPNOTCH™

WATER VOLUME:

Ground: minimum 100 L/ha or 40 L/ac

Air: 45 L/ha or 20 L/ac

RAINFASTNESS:

Do not apply when rain is imminent. Leaf coverage and penetration is critical for use.

SUPPORTED TANK MIXES:

Insecticides:

- SILENCER® 120 EC, Matador®, Coragen®, Voliam XPress®, Decis®

Herbicides:

- Assure® II, ARROW® 240 EC, ARROW ALL IN™, LEOPARD®, Liberty®, Poast Ultra®

MIXING INSTRUCTIONS:

1. Fill spray tank $\frac{1}{2}$ – $\frac{3}{4}$ full with water.
2. With agitator running, add required amount of TOPNOTCH™ and continue agitating while adding remainder of the water.
3. Begin application after TOPNOTCH™ is completely dispersed into the mix water, and maintain agitation during spraying operation.

CROP ROTATIONS:

Do not plant any other crop intended for food, grazing or any component of animal feed or bedding within 105 days of application.

PRE-HARVEST INTERVALS:

Cereals: 45 days

Field peas, Beans, Soybeans: 15 days

Lentils: 30 days

GRAZING RESTRICTIONS:

No restrictions

STORAGE:

Do not store below 0°C.

Quick Tips

Good spray coverage and canopy penetration are important to achieve the best results.



ADAMA

Always read and follow label directions.
Toll-free: 1.855.264.6262
Website: ADAMA.COM/CANADA

®/™ BUMPER, CUSTODIA, ORIUS and SILENCER are registered trademarks, and TOPNOTCH is a trademark of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies. © 2021 ADAMA Agricultural Solutions Canada Ltd.