

# **Active Ingredient:**

FOLPAN® 80 WDG

**Folpet** 

## Fungicide Group:

Group M

#### Formulation:

80% Folpet = SC

# Application Rates and Packaging:

0.5 – 2 kg/acre 2.5 – 10 acres/bag 5 – 20 acres/case

2 x 5 kg bags/case

#### **REGISTERED CROPS:**

- Apples
- Azalea
- Carnation
- Chrysanthemum
- Crabapples
- Field cucumbers
- Field tomatoes
- Grapes
- Iris
- Marigold
- Melons
- Poinsettia (greenhouse)
- Pumpkins
- Snapdragon
- Squash
- Strawberries
- Zinnia

#### **KEY DISEASES CONTROLLED:**

- Alternaria leaf spot
- Anthracnose
- Black rot
- Brooks spot
- Dead arm
- Didymellina leaf spot
- Downy mildew
- Fly speck
- Fruit rot
- Grey mould
- Leaf spot
- Phythium root rot
- · Powdery mildew
- Scab
- Septoria leaf spot
- Sooty blotch
- Stem rot

#### **HOW IT WORKS:**

Multi-site-contact mode of action with protective ability. Use FOLPAN® 80 WDG in a regularly scheduled maintenance program.

#### **APPLICATION TIMING AND CROP STAGING:**

- Apples, Crabapples: Apply in a regular protective schedule from green tip up until harvest. Do not apply more than 6 applications per season. See label for additional precautions.
- Cranberries: Apply when 5% of blossoms are open. Repeat 10–14 days later when 50–75% of blossoms are open.
- Grapes:
  - **Dead arm:** Apply when new shoots are 1–2 inches long and repeat when growth is 4–6 inches long.
  - Black rot, Downy mildew: Apply just before bloom, just after bloom and in first cover spray. For downy mildew, an additional application 2–3 weeks later may be needed.
  - Powdery mildew: Apply in a tank mix with Kumulus® DF at first sign of mildew and repeat after 10 days.



# FOLPAN® 80 WDG

#### **APPLICATION TIMING AND CROP STAGING (CONTINUED):**

- Field cucumbers, Melons, Pumpkins, Squash: Apply when first true leaves appear. Repeat at 7-day intervals until crop is harvested.
- Field tomatoes: Apply during first bloom and repeat at 7-day intervals until harvest.
- Strawberries: Apply before first infection and repeat at 7-day intervals to protect crop until harvest.
- Carnation, Poinsettia, Azalea, Marigold, Zinnia, Chrysanthemum, Iris, Snapdragon: Generally, apply when
  ornamental emerges and repeat at regular intervals. Consult label for timing on specific disease
  and ornamental.

#### **WATER VOLUME:**

- Ground: 400 1200 L/ac
- · Do not apply by air.

#### **RAINFASTNESS:**

No restrictions

#### **REGISTERED AND SUPPORTED TANK MIXES:**

Most commonly used insecticides, adjuvants and fungicides. Check compatibilities before making tank mixes.

#### **MIXING INSTRUCTIONS:**

- 1. Fill spray tank nearly full and pour recommended amount of FOLPAN® 80 WDG on surface of water.
- 2. Fungicide can be premixed in a bucket ½ filled with water. Mix can be poured through screen into nearly filled spray tank.
- 3. Finish filling tank.
- 4. Keep agitator running during filling and spraying.

#### **CROP ROTATIONS:**

No restrictions

#### **GRAZING RESTRICTIONS:**

No restrictions

#### **STORAGE:**

No restrictions

#### **PRE-HARVEST INTERVAL:**

Cranberries: 30 days
Field cucumbers: 1 day
Field tomatoes: 1 day
Grapes: 1 day

Melons:1 dayPumpkins:1 daySquash:1 dayStrawberries:1 day

### **QUICK TIPS:**

Fungicidal activity is reduced if combined with strongly alkaline materials such as hydrated lime. Do not use in combination with or closely following an oil spray.

Always read and follow registered product label instructions. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.





®/TM FOLPAN is a registered trademark of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies.