

Active Ingredient:

Diuron

Herbicide Group:

Group 7

Formulation:

80% Diuron = DF

Packaging:

1 x 10 kg bag

Application Rates (actual area sprayed):

NOTE: Rates vary by soil type-light sandy soils (LS), clay soils (C) or sandy loam soils (LM)

Asparagus:

- 0.45 0.91 kg/ac (LS)
- 0.91-1.82 kg/ac (C)

Grapes:

- 0.91-1.32 kg/ac (LS)
- 1.32-2.71 kg/ac (C)

Non-cropland

(broadcast application):

- 4.5-11 kg/ac (LS/LM)
- 16-22 kg/ac (C)

Spot treatment:

• $115 \text{ g}/100 \text{ m}^2 = 4.55 \text{ kg/ac}$

Irrigation and drainage ditches (restricted use):

- 115 g/100 m² = 4.55 kg/ac
- Do not apply by air

REGISTERED CROPS:

- Asparagus
- Grapes
- · Non-cropland

WEEDS CONTROLLED:

Asparagus, grapes:

• Weed seedlings such as crabgrass, foxtail, pigweed, ragweed and lamb's quarters

Non-cropland:

• Many annual and perennial grasses and broadleaf weeds

Spot treatment:

• Many annual and perennial grasses and broadleaf weeds plus deeprooted perennials such as quackgrass and toadflax

HOW IT WORKS:

Apply as a spray to the surface of the ground for control of weeds. The weed control effects are slow-to-appear and will not become apparent until moisture carries the chemical into the root zone.



KARMEX[®]

CROP STAGING:

Asparaaus:

- Do not apply to newly planted asparagus, or to young plants during the first growing season after setting or on plants with exposed roots as severe injury may result.
- · Apply as a band or broadcast treatment.
- Two (2) applications may be used.

Grapes:

• Apply only to established vineyards (at least 3 years old) as a band application to grape rows.

WATER VOLUME:

• 100-160 L/ac of actual area sprayed

RAINFASTNESS:

Moisture is desirable and required to move the active ingredient into the rooting zone.

SUPPORTED TANK MIXES:

None registered

MIXING INSTRUCTIONS:

- 1. Accurately weigh the amount of KARMEX® required and mix into necessary volume of water.
- 2. Material must be kept in suspension at all times by agitation, preferably by continuous mechanical or hydraulic means in the spray tank.
- 3. Openings in screens should be equal to or larger than 50 mesh.

CROP ROTATIONS:

Asparagus:

- Do not replant treated areas to any crop within 2 years after last application as injury to subsequent crops may result.
- Do not use on dwarf or semi-dwarf plants.

PRE-HARVEST INTERVAL:

None

GRAZING RESTRICTIONS:

Do not graze the treated areas or cut for hay.

· Can be applied any time, except when ground

· Best results are obtained when applied shortly

Irrigation and drainage ditches (restricted use):

For irrigation ditches, apply during the non-crop

season, and when ditch is not in use.

before weed growth begins.

STORAGE:

Non-cropland:

is frozen.

- · Keep from contact with fertilizers, insecticides, fungicides and seeds.
- · Do not contaminate food or feed.

QUICK TIPS:

KARMEX® requires moisture in the form of rainfall or irrigation after treatment. Sandy and sandy loam soils require lower rates than clay and high organic soils. Use higher rates where perennial weeds are dominant on non-cropland. On non-cropland, regrowth of plantain, thistle or wild carrot will indicate when retreatment is necessary.

Always read and follow registered product label instructions. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. Toll-free: 1.855.264.6262





^{®/TM} KARMEX is a registered trademark of ADAMA Agricultural Solutions Canada Ltd. All other products mentioned are trademarks of their respective companies. © 2021 ADAMA Agricultural Solutions Canada Ltd.