



ADAMA

# BROMOTRIL®

Tough broadleaf weed control with tank-mix flexibility and excellent crop safety.



HERBICIDE

## Active Ingredient:

Bromoxynil

## Herbicide Group:

Group 6

## Formulation:

Bromoxynil Octanoate Ester  
235 g/L = EC

## Application Rates and Packaging:

0.49 – 0.57 L/ac or  
17 – 20 ac/9.7 L jug

2 × 9.7 L jugs/case

### Post-Emergent Rate:

0.49 – 0.57 L/ac or  
17 – 20 ac/9.7 L jug

### Pre-Seed:

BROMOTRIL® at 0.49 L/ac  
(20 ac/9.7 L jug) tank mixed  
with glyphosate at a rate of  
175 g a.i./ac

## REGISTERED CROPS:

| Crop                | Stage                             |
|---------------------|-----------------------------------|
| Barley, Oats, Wheat | Pre-seed burn-off with glyphosate |

| Crop   | Crop Leaf Stage  |
|--|--|
| Alfalfa (seedling)   | 2 – 6 trifoliate                                       |
| Alfalfa (established)  | Spring: before the crop begins to shield the weeds     |
| Barley, Oats, Triticale, Wheat (spring, durum)   | 2 leaf to early flag                                   |
| Winter wheat   | Fall: 2 – 4 leaf<br>Spring: first growth to early flag |
| Corn (field, sweet) with drop pipes  | Beyond 8 leaf  |
| Canary seed (seed production only)   | 3 – 5 leaf   |
| Fall rye   | Spring: from first growth to early flag                |
| Flax   | 2 – 4 inches in height                                 |
| Forage millet, Sorghum   | 4 leaf to 8 inches                                     |
| Seedling grasses (seed production only):<br>Bromegrass, Fescue (creeping red, meadow),<br>Orchard grass, Reed canarygrass, Russian<br>wildrye, Timothy, Wheatgrass (crested,<br>intermediate, slender, tall) | 2 – 4 leaf (year of establishment only)                |

## WEEDS CONTROLLED:

### Seedling up to 4-leaf stage:

- American nightshade
- Bluebur
- Cocklebur
- Common ragweed
- Cow cockle<sup>1</sup>
- Green smartweed
- Kochia<sup>2</sup>
- Lady's thumb
- Pale smartweed
- Pigweed<sup>1</sup>
- Russian thistle<sup>2</sup>
- Stinkweed<sup>1</sup>
- Velvetleaf<sup>3</sup>
- Wild mustard<sup>1</sup>

### Seedling up to 8-leaf stage:

- Common buckwheat
- Common groundsel
- Lamb's quarters
- Tartary buckwheat
- Wild buckwheat

<sup>1</sup> In normal conditions, it will be controlled up to 4-leaf stage. Plants beyond this stage are unlikely to be controlled; the higher rate generally gives better results.

<sup>2</sup> Spray before plants are 2 inches high.

<sup>3</sup> Spray before plants are 3 inches high.

# BROMOTRIL®

## HOW IT WORKS:

BROMOTRIL® is a contact herbicide which controls Group 2 and Group 9 resistant biotypes. Leaves will yellow in 2–4 days with complete control in 7–14 days.

## WATER VOLUME:

- Ground: 20–40 L/ac
- Aerial (wheat and barley only): 8–16 L/ac

## RAINFASTNESS:

30 minutes

## REGISTERED AND SUPPORTED POST-EMERGENT TANK MIXES:

### Post-emergent Herbicides:

- Spring wheat: 2,4-D Ester, LADDER ALL IN®, MCPA Ester, BISON® 400 L, Traxos®
- Winter wheat: 2,4-D Ester, MCPA Ester, BISON® 400 L, Traxos®
- Barley: 2,4-D Ester, MCPA Ester, BISON® 400 L
- Oats: MCPA Ester
- Corn: Accent®, Atrazine, Banvel®(Dicamba), Ultim®
- Fall rye: MCPA Ester
- Flax: MCPA Ester or MCPA-K
- Canary seed: MCPA Ester
- Seedling grasses: MCPA Ester

### Pre-seed:

- Glyphosate

### Fungicides:

- BUMPER® 432 EC

## MIXING INSTRUCTIONS:

1. Fill spray tank ½ full with water.
2. Add required amount of BROMOTRIL®. Begin agitation.
3. If tank mixing, add the recommended amount of the tank-mix partner product to the spray tank first, agitate and then add BROMOTRIL® (unless otherwise directed by the BROMOTRIL® and tank-mix partner label).
4. Add the remaining amount of water while agitation continues.

## CROP ROTATIONS:

No restrictions.

## STORAGE:

Avoid freezing.

## PRE-HARVEST INTERVAL:

30 days

## GRAZING RESTRICTIONS:

- Do not use treated crops for grazing of livestock or green feed until 30 days after application.
- Do not cut treated crops for forage until 30 days after application.

## QUICK TIPS:

Avoid spraying if temperatures are above 25° C. Leaf scorching may occur in corn and flax if applied during or after adverse growing conditions, such as cool and wet or hot (above 27° C) and humid weather. For best results, spray when weeds are in the seedling stage.



Always read and follow registered product label instructions. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.  
Toll-free: 1.855.264.6262  
Website: ADAMA.COM/CANADA

BISON®, BROMOTRIL®, BUMPER®, LADDER ALL IN® are registered trademarks of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies. © 2020 ADAMA Agricultural Solutions Canada Ltd.