

Active Ingredient

80% Diuron = DF

Packaging

1 x 10 kg bag

Application Rates & Acres Treated

Rate: 0.45 – 2.71 kg/ac
Acres Treated: 4 – 20 ac/bag

Asparagus:

- Light sandy soils: 0.45 – 0.91 kg/ac (actual area sprayed)
- Clay soils: 0.91–1.82 kg/ac (actual area sprayed)

Grapes:

- Light sandy soils: 0.91–1.32 kg/ac (actual area sprayed)
- Clay soils: 1.32 2.71 kg/ac (actual area sprayed)

Water Volume

Ground: 100 – 160 L/ac (26-42 US gal/ac)

Aerial: Do not apply by air.

Rainfastness

Rain is desirable and required to move product to the rooting zone.

REGISTERED CROPS

- Asparagus
- Grapes

WEEDS CONTROLLED

Weed seedlings such as crabgrass, foxtail, pigweed, ragweed and lamb's quarters. See the label for the full listing of weeds controlled.

HOW IT WORKS

Applied as a spray to the surface of the ground for control of weeds. The weed control effects are slow to appear and will not become apparent until moisture carries the chemical into the root zone.

CROP STAGING

Asparagus

- Do not replant treated areas to any crop within 2 years after last application as injury to subsequent crops may result.
- Do not use on dwarf or semi-dwarf plants.

Grapes:

Apply only to established vineyards (at least 3 years old) as a band application to grape rows.



KARMEX®

SUPPORTED TANK MIXES

None registered

MIXING INSTRUCTIONS

- 1. Accurately weigh the amount of KARMEX® required and mix into necessary volume of water.
- 2. Material must be kept in suspension at all times by agitation, preferably by continuous mechanical or hydraulic means in the spray tank.
- 3. Openings in screens should be equal to or larger than 50 mesh.

CROP ROTATIONS

Asparagus:

- Do not replant treated areas to any crop within 2 years after last application as injury to subsequent crops may result.
- Do not use on dwarf or semi-dwarf plants.

PRE-HARVEST INTERVAL

None

GRAZING RESTRICTIONS

Do not graze the treated areas or cut for hay.

STORAGE

May be stored at any temperature

QUICK TIPS:

KARMEX® requires moisture in the form of rainfall or irrigation after treatment. Sandy and sandy loam soils require lower rates than clay and high organic soils. Use higher rates where perennial weeds are dominant on non-cropland. On non-cropland, regrowth of plantain, thistle or wild carrot will indicate when retreatment is necessary.



Always read and follow registered product label instructions. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.

1.855.264.6262 ADAMA.COM

^{®/TM} KARMEX is a registered trademark of ADAMA Agricultural Solutions Canada Ltd. All other products mentioned are trademarks of their respective companies.

© 2022 ADAMA Agricultural Solutions Canada Ltd.