ARMORY[®] 240

Provides fast drydown of crops, protecting yield and grade and reducing disease transmission late in the season.



Active Ingredient Diquat 240 g/L = EC

Packaging

Case: 2 x 10 L jugs Bulk: 5 x 120 L drums Totes: 450 L; 1000 L

Application Rates & Acres Treated

Ground: 360 – 1860 ml/ac **Aerial:** 690 – 1090 ml/ac

Acres Treated:

- Jug: 7–28 acres
- Drum: 83 333 acres
- 450 L tote: 320–1250 acres
- 1000 L tote: 700 2775 acres

Water Volume

Ground: 90 – 200 L/ac (24 – 53 US gal/ac) Aerial: Minimum 18 L/ac (5 US gal/ac)

Rainfastness

30 minutes

GROUP 22

REGISTERED CROPS

- Alfalfa
- Beans (white and red kidney, adzuki)
- Birdsfoot trefoil
- Canola
- Chickpeas
- Lentils

USES AND WEEDS CONTROLLED

- Potato vines
- Corn spurry in oats
- Desiccation for beans, and legume forage seed crops
- Weeds in stale seedbeds (vegetables and field crops)
- Weeds in vegetables (inter-row directed)
- Suppression of perennial grasses under apple trees

APPLICATION RATES

Crop	Rate
Potatoes	 Ground: 510 – 1420 ml/ac Aerial: 690 – 930 ml/ac
Vegetables	 Ground: 930–1860 ml/ac Aerial: Do not apply by air.
Fruit	• Ground: 1860 ml/ac
Beans, canola, chickpeas, field peas, flax, lentils, mustard, sunflowers	 Ground: 500 – 690 ml/ac Aerial: 690 – 930 ml/ac
Legumes	 Ground: 690 – 1090 ml/ac Aerial: 690 – 1090 ml/ac
Oats	 Ground: 360 – 510 ml/ac Aerial: Do not apply by air.

- Peas (field and dry)
- Potatoes
- Red and white clover
- Soybeans
- Sunflowers

ARMORY[®] 240

HOW IT WORKS

ARMORY[®] 240 works on contact to disrupt plant cells and is rainfast in 30 minutes, leading to more rapid drydown of plants and weeds when compared to systemic herbicides. Harvesting can typically begin within 4–10 days, depending on crop and weather conditions.

REGISTERED AND SUPPORTED TANK MIXES

- Agral[®] 90, LI 700[®], Liberate[®], other non-ionic surfactants
- Carfentrazone

MIXING INSTRUCTIONS

- 1. Fill the spray tank ¾ full with water.
- 2. Add the required amount of ARMORY® 240 into the sprayer.
- 3. Agitate until the herbicide is thoroughly mixed.
- Continue agitation while adding the required amount of recommended registered surfactant at 0.10% v/v non-ionic surfactant (NIS) or 0.25% v/v LI 700[®].
- 5. Complete filling the tank to the desired level with water.

ADJUVANT RATE

- LI 700[®] @ 0.25% v/v
- Non-ionic surfactant (NIS)@ 0.10% v/v

GRAZING RESTRICTIONS

Crop waste remaining after harvest (e.g. pea vines, alfalfa stems) may be used as a feed supplement for livestock.

STORAGE

Do not freeze.

QUICK TIPS:

Best results under cloudy conditions or in the evening. Suggested conditions for aerial applications are a temperature below 25° C, humidity above 50% and wind speed below 9 km/hr at flying height.



Always read and follow registered product label instructions. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. 1.855.264.6262 ADAMA.COM

 ® ARMORY is a registered trademark of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies.

 © 2023 ADAMA Agricultural Solutions Canada Ltd.